

MA40254 DIFFERENTIAL AND GEOMETRIC ANALYSIS : EXERCISES 4

Hand in answers by 1:15pm on Wednesday 2 November for the Seminar of Thursday 3 November

Homepage: <http://moodle.bath.ac.uk/course/view.php?id=57709>

**0** (Warmup). Compute the tangent space  $T_pM$  to the 1-dimensional submanifold  $M := \{(x, y) \in \mathbb{R}^2 : y = x^2\}$  of  $\mathbb{R}^2$  at the point  $p = (t, t^2)$ .

**1.** (i) Let  $M \subseteq \mathbb{R}^s$  be a submanifold. Let  $P \subseteq \mathbb{R}^s$  be an open subset that contains  $M$ , and let  $f : P \rightarrow \mathbb{R}^m$  be a smooth function. Suppose that the restriction of  $f$  to  $M$  is constant. Show that  $T_pM \subseteq \ker Df_p \subseteq \mathbb{R}^s$  for any  $p \in M$ .

(ii) Let  $P \subseteq \mathbb{R}^s$  be an open subset,  $f : P \rightarrow \mathbb{R}^m$  a smooth function,  $q \in \mathbb{R}^m$  a regular value of  $f$ , and  $M := f^{-1}(q)$ . Show that  $T_pM = \ker Df_p \subseteq \mathbb{R}^s$  for any  $p \in M$ .

**2.** For points  $x, y \in \mathbb{R}^2$  with  $x \neq y$ , let  $S(x, y) = \{tx + (1-t)y : t \in (0, 1)\} \subset \mathbb{R}^2$ . For which  $x, y, x', y' \in \mathbb{R}^2$  is  $S(x, y) \cup S(x', y')$  a submanifold of  $\mathbb{R}^2$ ?

**3.** Let  $O(n) = \{A \in GL_n(\mathbb{R}) : A^T = A^{-1}\}$ . Show that  $O(n)$  is a submanifold of  $M_{n,n}(\mathbb{R})$ . What is  $T_I O(n) \subseteq M_{n,n}(\mathbb{R})$  (the tangent space of  $O(n)$  at the identity matrix  $I \in O(n)$ )?

**4.** Let  $V$  be a real vector space of dimension  $n$ , and  $\text{Alt}^2(V)$  the space of alternating 2-forms on  $V$ , that is bilinear maps  $\omega : V \times V \rightarrow \mathbb{R}$  such that  $\omega(v, v) = 0$  for any  $v \in V$ . What is the dimension of  $\text{Alt}^2(V)$ ?

**5.** For  $v_1, \dots, v_n \in \mathbb{R}^n$ , let  $\text{Det}(v_1, \dots, v_n) \in \mathbb{R}$  denote the determinant of the  $n \times n$  matrix with columns  $v_1, \dots, v_n$ .

(i) Show that  $\text{Det}$  spans  $\text{Alt}^n(\mathbb{R}^n)$ .

(ii) For any  $u, v \in \mathbb{R}^3$ , show that there is a unique  $u \times v \in \mathbb{R}^3$  such that for any  $w \in \mathbb{R}^3$ ,

$$\text{Det}(u, v, w) = (u \times v) \cdot w.$$

Here the right hand side is the Euclidean inner product of the vectors  $u \times v$  and  $w$ .