

MA40254 DIFFERENTIAL AND GEOMETRIC ANALYSIS : EXERCISES 6

Hand in answers by 1:15pm on Wednesday 15 November for the Seminar of Thursday 16 November  
 Homepage: <http://moodle.bath.ac.uk/course/view.php?id=57709>

0 (Warmup). Let  $\gamma = \frac{-x_2 dx_1 + x_1 dx_2}{x_1^2 + x_2^2} \in \Omega^1(\mathbb{R}^2 \setminus \{0\})$ . Show that  $d\gamma = 0$ .

1. Let  $U = \{p \in \mathbb{R}^4 : x_2(p) \neq 0\}$ , and  $\varphi = (x_1, x_2^2 x_3, x_4/x_2) : U \rightarrow \mathbb{R}^3$ , where  $x_1, x_2, x_3, x_4$  are the coordinate functions on  $U \subseteq \mathbb{R}^4$ . Let  $\alpha = y_1 dy_2 \wedge dy_3 \in \Omega^2(\mathbb{R}^3)$  where  $y_1, y_2, y_3 : \mathbb{R}^3 \rightarrow \mathbb{R}$  denote the coordinate functions.

(i) Express  $\varphi^* \alpha$  and  $\varphi^* d\alpha$  in standard form, i.e., as a sum of terms  $f dx_I$ .

(ii) Compute directly  $d(\varphi^* \alpha)$ . What do you observe?

2. For  $U$  open in  $\mathbb{R}^n$ ,  $\alpha \in \Omega^k(U)$ ,  $p \in U$ , and  $v_1, \dots, v_k \in \mathbb{R}^n$ , show that

$$d\alpha_p(v_0, \dots, v_k) = \sum_{i=0}^k (-1)^i D\alpha_p(v_i)(v_0, \dots, \widehat{v}_i, \dots, v_k)$$

where  $v_0, \dots, \widehat{v}_i, \dots, v_k$  denotes the list obtained from  $v_0, \dots, v_k$  by omitting  $v_i$ . Equivalently

$$d\alpha_p = \sum_{i=0}^k \text{sgn}(\sigma_i) \sigma_i \cdot D\alpha_p^\vee,$$

where  $\sigma_i = (0 \ 1 \ \dots \ i)^{-1} \in G := \text{Sym}(\{0, 1, \dots, k\}) \cong S_{k+1}$ , and for  $\sigma \in G$  and  $\beta \in \text{Alt}^{k+1}(\mathbb{R}^n)$ ,  $(\sigma \cdot \beta)(v_0, \dots, v_k) = \beta(v_{\sigma(0)}, \dots, v_{\sigma(k)})$  (which is  $\beta(v_i, v_0, \dots, \widehat{v}_i, \dots, v_k)$  when  $\sigma = \sigma_i$ ).

3 (Less essential). Let  $U \subseteq \mathbb{R}^3$  an open subset. Given a vector-valued function  $v = (v_1, v_2, v_3) : U \rightarrow \mathbb{R}^3$ , define  $v^\flat \in \Omega^1(U)$  and  $v \lrcorner \text{Det} \in \Omega^2(U)$  by applying the corresponding operations on vectors pointwise.

(i) Let  $\text{div}(v) : U \rightarrow \mathbb{R}$  be defined by

$$\text{div}(v) = \frac{\partial v_1}{\partial x_1} + \frac{\partial v_2}{\partial x_2} + \frac{\partial v_3}{\partial x_3}.$$

Show that

$$d(v \lrcorner \text{Det}) = \text{div}(v) \text{Det} \in \Omega^3(U).$$

(ii) Let  $\text{curl}(v) : U \rightarrow \mathbb{R}^3$  be defined by

$$\text{curl}(v) = \left( \frac{\partial v_3}{\partial x_2} - \frac{\partial v_2}{\partial x_3}, \frac{\partial v_1}{\partial x_3} - \frac{\partial v_3}{\partial x_1}, \frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \right).$$

Show that

$$d(v^\flat) = \text{curl}(v) \lrcorner \text{Det} \in \Omega^2(U).$$

4. Let  $x_1, x_2 : \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{R}$  be the coordinate functions. Which of the following elements of  $\Omega^1(\mathbb{R}^2 \setminus \{0\})$  are closed? Which are exact?

(i)  $\alpha = -2x_1 x_2 dx_1 + x_1^2 dx_2$

(ii)  $\beta = x_2 dx_1 + x_1 dx_2$

(iii)  $\gamma = \frac{-x_2 dx_1 + x_1 dx_2}{x_1^2 + x_2^2}$