

Spectral decompositions and nonnormality of boundary integral operators in acoustic scattering

T. BETCKE[†], J. PHILLIPS[‡], E. A. SPENCE[§]

[Received on 8 January 2012]

Understanding the spectral properties of boundary integral operators in acoustic scattering has important practical implications, such as for the analysis of the stability of boundary element discretisations or the convergence of iterative solvers as the wavenumber k grows. Yet little is known about spectral decompositions of the standard boundary integral operators in acoustic scattering. Theoretical results are mainly available on the unit disk, where these operators diagonalise in a simple Fourier basis. In this paper we investigate spectral decompositions for more general smooth domains. Based on the decomposition of the acoustic Green's function in elliptic coordinates we give spectral decompositions on ellipses. For general smooth domains we show that approximate spectral decompositions can be given in terms of circle Fourier modes transplanted onto the boundary of the domain. An important underlying question is whether or not the operators are normal. Based on previous numerical investigations it appears that the standard boundary integral operators are normal only when the domain is a ball and here we prove that this is indeed the case for the acoustic single layer potential. We show that the acoustic single, double and conjugate double layer potential are normal in a scaled inner product on the ellipse. On more general smooth domains the operators can be split into a normal component plus a smooth perturbation. Numerical computations of pseudospectra are presented to demonstrate the nonnormal behaviour on general domains.

Keywords: acoustic scattering; boundary integral operators; spectra; pseudospectra

1. Introduction

The study of spectral properties of partial differential operators has seen tremendous advances in recent decades. An important question is whether an operator is nonnormal and how this nonnormality influences analytical and numerical properties of the operator. The development of pseudospectra as a tool for understanding nonnormality has given great insight into diverse areas, such as existence of solutions to linear PDE problems, the behaviour of numerical solvers, or transient behaviour in dynamical systems. For a wonderful overview about this field we refer to the recent book by Trefethen & Embree (2005). However, whilst nonnormality has been studied extensively for partial differential operators, little is known for boundary integral operators. The aim of this paper is to study nonnormality in the context of boundary integral operators in acoustic scattering.

Let $\Omega \subset \mathbb{R}^2$ be a bounded domain and denote the boundary of Ω by Γ . For the majority of this paper we assume that Γ is an analytic curve. Denote by $\Omega^+ := \mathbb{R}^2 \setminus \overline{\Omega}$ the exterior of Ω . Furthermore, let $g : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{C}$ be the acoustic Green's function defined by $g(x, y) = \frac{i}{4} H_0^{(1)}(k|x - y|)$, where k is the wavenumber. We define the following operators, which are bounded operators on $L^2(\Gamma)$ (for mapping properties of these operators in Sobolev spaces see, for example, McLean (2000)).

- *The acoustic single layer potential*

$$[S\phi](x) = \int_{\Gamma} g(x, y)\phi(y)ds(y), \quad x \in \Gamma$$

[†]Department of Mathematics, University College London, UK. Corresponding author. Email: t.betcke@ucl.ac.uk. Timo Betcke is supported by Engineering and Physical Sciences Research Council (EPSRC) Grant EP/H004009/1.

[‡]Department of Mathematics, University College London, UK. Email: joel.phillips@ucl.ac.uk. Joel Phillips is supported by EPSRC Grant EP/H004009/1.

[§]Department of Mathematics, University of Bath, UK. Email: e.a.spence@bath.ac.uk. Euan Spence is supported by EPSRC Grant EP/I025995/1.

- *The acoustic double layer potential*

$$[K\phi](x) = \frac{\phi(x)}{2} + \int_{\Gamma} \frac{\partial}{\partial n(y)} g(x,y) \phi(y) ds(y), \quad x \in \Gamma$$

- *The acoustic conjugate double layer potential*

$$[T\phi](x) = -\frac{\phi(x)}{2} + \int_{\Gamma} \frac{\partial}{\partial n(x)} g(x,y) \phi(y) ds(y), \quad x \in \Gamma$$

Here, n denotes the exterior normal to Ω . The double layer and conjugate double layer potentials usually appear without the factors of $\pm 1/2$. Our reason for defining them like this is that if we denote the exterior trace operator by γ_0^+ and the exterior normal derivative by γ_1^+ , then the above boundary operators are obtained from the single and double layer potentials,

$$[\mathcal{S}\phi](x) = \int_{\Gamma} g(x,y) \phi(y) ds(y), \quad [\mathcal{K}\phi](x) = \int_{\Gamma} \frac{\partial}{\partial n(y)} g(x,y) \phi(y) ds(y), \quad x \in \Omega^+$$

by $S = \gamma_0^+ \mathcal{S}$, $K = \gamma_0^+ \mathcal{K}$, $T = \gamma_1^+ \mathcal{S}$. The fact that S, K , and T arise directly as the traces of \mathcal{S} and \mathcal{K} makes the algebra of the spectral decompositions considered below more convenient. We will also be interested in the combined potential operator

$$A_{\eta} := 2(I + T - i\eta S)$$

defined for $\eta \neq 0$. In Chandler-Wilde & Langdon (2007) it was shown that A_{η} is a bounded operator with bounded inverse for any $k > 0$, $\eta \neq 0$, mapping $L^2(\Gamma)$ into $L^2(\Gamma)$ for any Lipschitz boundary Γ .

The main motivation for studying the behaviour of these integral operators is that they can be used to solve scattering problems modelled by the Helmholtz equation, with the combined potential operator A_{η} giving rise to a uniquely-solvable integral equation for the solution of the Helmholtz equation with Dirichlet boundary conditions (see Section 6 for details). Another important boundary integral operator is the hypersingular operator. This is $\gamma_1^+ \mathcal{K}$ and arises when formulating boundary value problems for the Helmholtz equation with Neumann boundary conditions. In this paper we are mainly focused on the behaviour of the combined potential operator A_{η} and so we do not consider the hypersingular operator.

The spectral behaviour of the operators S, K, T and A_{η} is especially simple on the unit circle. We have the following well known result (see e.g. Kress & Spassov (1983); Kress (1985); Domínguez *et al.* (2007)).

THEOREM 1.1 On the unit circle it holds that

$$Su_n = \lambda_n^{(S)} u_n, \quad Ku_n = \lambda_n^{(K)} u_n, \quad Tu_n = \lambda_n^{(T)} u_n, \quad A_{\eta} u_n = \lambda_n^{(A)} u_n,$$

where in polar coordinates $u_n(\theta) = e^{in\theta}$ and

$$\begin{aligned} \lambda_n^{(S)} &= \frac{\pi i}{2} H_n^{(1)}(k) J_n(k), \\ \lambda_n^{(K)} &= \frac{k\pi i}{4} H_n^{(1)}(k) (J_{n-1}(k) - J_{n+1}(k)), \\ \lambda_n^{(T)} &= \frac{k\pi i}{4} (H_{n-1}^{(1)}(k) - H_{n+1}^{(1)}(k)) J_n(k) \\ \lambda_n^{(A)} &= 2(1 + \lambda_n^{(T)} - i\eta \lambda_n^{(S)}). \end{aligned} \tag{1.1}$$

Proof. The result is a consequence of the Graf addition formula (NIST Digital Library, eq. 10.23.7), stating that

$$H_0^{(1)}(k|re^{i\theta} - \rho e^{i\phi}|) = \sum_{n=-\infty}^{\infty} e^{in(\theta-\phi)} H_n^{(1)}(kr_{>}) J_n(kr_{<}), \tag{1.2}$$

where $r_{>} = \max(r, \rho)$ and $r_{<} = \min(r, \rho)$. \square

(It is well known that on the unit circle $K - \frac{1}{2}I = T + \frac{1}{2}I$ and this can be seen from the eigenvalues using the facts that $J_{n-1}(k) - J_{n+1}(k) = 2J'_n(k)$, $H_{n-1}^{(1)}(k) - H_{n+1}^{(1)}(k) = 2H_n^{(1)'}(k)$, and $W[J_n, H_n^{(1)}, k] = 2i/(\pi k)$, where W denotes the Wronskian.)

An important feature of the circle case is that the operators S , K , T and A_η are normal. since they diagonalize in a unitary basis and therefore commute with their adjoints. Recall that a bounded operator A acting on a Hilbert space \mathcal{H} is *normal* if and only

$$AA^* = A^*A,$$

where A^* is the adjoint of A . Normal operators are simple in the sense that they can be shown to be unitarily equivalent to multiplication operators (see for example (Conway, 1985, Theorem 4.6)). In particular, if A is compact and normal then there exists an orthonormal system $\{\phi_j\}$, such that

$$A = \sum_j \lambda_j \phi_j \langle \phi_j, \cdot \rangle,$$

where the λ_j are the eigenvalues of A and the ϕ_j are the associated eigenfunctions. Hence, the behavior of a normal operator can be completely determined by spectral information.

For more general domains than the circle, little is known about spectral decompositions, however understanding the spectral properties and normality of these boundary integral operators has both theoretical and practical implications. On the theoretical side, in Ramm (1973, 1980) Ramm asked when the eigen-system of the single layer potential operator in 3-d forms a complete basis in $L^2(\Gamma)$ and gave normality as a sufficient condition. On the practical side, whether an operator is normal or not affects the convergence of iterative solvers such as GMRES.

An important question, therefore, is whether the operators S , K , T and A_η are normal on any domains other than the circle (in 2-d, or sphere in 3-d). Numerical experiments in Betcke & Spence (2011) demonstrate that the combined potential operator A_η appears to be nonnormal on a wide range of 2-d domains, which leads to the question of whether the circle is the only 2-d domain on which these operators are normal. In this paper we prove for the single layer potential operator S that, amongst all sufficiently smooth 2-d and 3-d domains, balls are the only domains for which the operator is normal in the standard $L^2(\Gamma)$ inner product (see Theorem 3.1 below). The rest of this paper is, in some sense, an investigation into whether the desirable property of normality can be recovered, either in special cases or approximately.

If the Helmholtz equation is separable in a particular orthogonal coordinate system, then the Green's function can be expanded in terms of special functions ((1.2) is this expansion for polar coordinates), and this allows the derivation of eigenvalue decompositions of the operators S , K , and T for certain domains. We show that these decompositions lead to the operators being normal in a modified L^2 inner product, and demonstrate this in the case on an ellipse.

The explicit eigenvalue decompositions above are only possible for a very restrictive class of domains. Nevertheless, for arbitrary smooth domains we derive an approximate eigenvalue decomposition that relates eigenvalues on smooth domains to eigenvalues on the unit circle. In particular, it allows us to separate the spectrum into $\mathcal{O}(k)$ eigenvalues that are not well approximated by circle eigenvalues, and a tail of eigenvalues that are well-approximated by circle eigenvalues with associated approximate eigenfunctions that are orthogonal in a modified L^2 inner product. This result shows that nonnormality appears mainly to affect a finite number of $\mathcal{O}(k)$ eigenvalues.

The paper is organized as follows. In Section 2 we briefly introduce two related tools for the study of nonnormality, namely pseudospectra and numerical ranges. We then demonstrate the effect of nonnormality on norm bounds and on coercivity constants for boundary integral operators in acoustic scattering. In Section 3 we present a proof that, amongst all sufficiently smooth domains, circles (respectively balls in 3-d) are the only ones for which the acoustic single layer potential is normal. The proof is based on transforming the question of normality into a uniqueness problem of equilibrium potentials, which has been solved by Reichel (1997). In Section 4 we use the decomposition of the acoustic Green's function in elliptic coordinates to derive an eigenvalue decomposition of the operators S , K and T on the boundary of an ellipse. This allows one to prove that these operators are normal in a scaled L^2 inner product. In Section 5 approximate eigendecompositions are derived for general analytic domains. Applications of the results to the combined potential operator A_η are treated in Section 6. Numerical examples will be given for two domains. The paper finishes with conclusions in Section 7.

Notation in this paper is mostly standard. Unless otherwise stated we use $\|\cdot\|$ for the norm induced by the standard complex $L^2(\Gamma)$ inner product

$$(u, v) := \int_{\Gamma} u(s) \overline{v(s)} ds,$$

where Γ is the boundary of a given domain. For disambiguation we will also use from time to time the symbols $\|\cdot\|_{L^2(\Gamma)}$ and $(\cdot, \cdot)_{L^2(\Gamma)}$ to emphasise the standard $L^2(\Gamma)$ norm and inner product. Frequently we will make use of a scaled L^2 inner product defined by

$$(u, v)_{L^2(F^{-1}, \Gamma)} := \int_{\Gamma} (F^{-1}u)(y) \overline{v(y)} ds(y)$$

and the induced norm

$$\|v\|_{L^2(F^{-1}, \Gamma)} := \left(\int_{\Gamma} (F^{-1}v)(y) \overline{v(y)} ds(y) \right)^{1/2},$$

where F is a given multiplication operator.

2. Nonnormality and its consequences for the solution of boundary integral equations

In this section we first give a brief introduction to two widely used tools for the analysis of nonnormality, namely pseudospectra and numerical ranges. Then by looking at norms and coercivity constants we give examples of how nonnormality influences the behavior of boundary integral operators.

2.1 Pseudospectra and numerical ranges of operators

A widely used tool for investigating nonnormality is via the pseudospectrum of an operator. It can be characterised by the following three equivalent definitions (Trefethen & Embree, 2005).

DEFINITION 2.1 (Pseudospectrum of a bounded operator) Let A be a bounded operator on a Hilbert space \mathcal{H} and $\varepsilon > 0$ be arbitrary: The ε -pseudospectrum $\sigma_{\varepsilon}(A)$ of A is the set of $z \in \mathbb{C}$ defined equivalently by any of the conditions

1. $\|(z - A)^{-1}\| > \varepsilon^{-1}$.
2. $z \in \sigma(A + E)$ for some bounded operator E with $\|E\| < \varepsilon$.
3. $z \in \sigma(A)$ or $\|(z - A)u\| < \varepsilon$ for some $u \in \mathcal{H}$ with $\|u\| = 1$.

If $\|(z - A)u\| < \varepsilon$ with $\|u\| = 1$ then z is an ε -pseudoeigenvalue of A and u is a corresponding ε -pseudoeigenvector (or pseudoeigenfunction or pseudomode).

The definition in (Trefethen & Embree, 2005, Chapter 4) is stated in terms of closed operators acting on Banach spaces. However, in this paper we are only interested in bounded Hilbert space operators. For normal matrices the ε -pseudospectrum is just the union of open disks of radius ε around the spectral values. In general, if A is nonnormal the form of the pseudospectra can be very different.

A rougher but also very useful tool to investigate nonnormality is the numerical range of a bounded linear operator A acting on a Hilbert space \mathcal{H} . It is defined as

$$W(A) := \{ \langle Au, u \rangle, u \in \mathcal{H}, \|u\| = 1 \}.$$

The numerical range is always a convex set such that the spectrum $\sigma(A) \subset \overline{W(A)}$. Furthermore, the closure of the numerical range of a normal operator is the convex hull of its eigenvalues. There exists a simple relationship between the numerical range and the ε -pseudospectrum, namely

$$\sigma_{\varepsilon}(A) \subset W(A) + \Delta_{\varepsilon},$$




FIG. 1: A crescent domain parameterised by $z(t) = e^{it} - \frac{0.1}{e^{it}+0.9}$, $t \in [0, 2\pi]$.




FIG. 2: Contour lines of the ε -pseudospectrum of the operator A_η for $\eta = k = 10$ on the crescent domain, where $\varepsilon = 10^\tau$ with the values of τ given in the legend. The dashed line is the boundary of the numerical range $W(A_\eta)$.




FIG. 3: The norm of the operators $\|S\|$, $\|K\|$ and $\|T\|$ on the unit circle.

where Δ_ε is the open disk around the origin of radius ε (Trefethen & Embree, 2005, Chapter 17). In Figure 2 we plot the ε -pseudospectrum and the numerical range for the combined potential operator A_η in $L^2(\Gamma)$ with $\eta = k = 10$ for the crescent domain shown in Figure 1 (see Section 6 for more examples and details of the computation). The differently coloured contour lines are the boundaries of the ε -pseudospectrum for $\varepsilon = 10^\tau$, where τ are the values given in the legend. The dashed line is the boundary of the numerical range and the dots are the eigenvalues. It is clear from this plot that the operator is not normal. The numerical range is much larger than the convex hull of the eigenvalues. The ε -pseudospectrum was computed using `EigTool`, a Matlab software package developed by Wright (2002).

2.2 Norm estimates

If the compact operator A is normal then by diagonalisation in its eigenbasis it follows that

$$\|A\| = \sup_n |\lambda_n(A)|.$$

This was used in Domínguez *et al.* (2007) to give estimates for the k -dependence of the norm of combined boundary integral operators on the circle in 2-d and the sphere in 3-d (following similar investigations by Kress & Spassov (1983), Kress (1985)). From the analysis in (Domínguez *et al.*, 2007, Section 4) it follows that

$$\|S\| \leq Ck^{-2/3}, \quad \|K\| \leq Ck^{1/3}$$

for the circle. Numerical experiments in Betcke *et al.* (2011) indicated that a sharper bound on $\|K\|$ is

$$\|K\| \leq C.$$

Indeed, this bound was proved for the unit sphere in 3-d in Banjai & Sauter (2007), and since the eigenvalues for the unit sphere are very similar to the eigenvalues for the unit circle this bound also holds for the unit circle. In Figure 3 we show the k -dependence of the norm of the operators S , K and T on the unit circle. For the BEM discretisation we used roughly 30 elements per wavelength and quadratic polynomials as basis functions on each element. The results were computed by taking the maximum over the absolute values of the first 100 eigenvalues computed from the formulas in Theorem 1.1. These computation con-




FIG. 4: Absolute value of the largest eigenvalue $\lambda_{max}(S)$ and $\|S\|_{L^2(\Gamma)}$ of the single layer potential S on the boundary of the crescent domain from Figure 1 for growing values k .

firm both the $O(k^{-2/3})$ decay for the single layer potential and the fact that the norms of K and T behave like $O(1)$ independently of k .

If A is nonnormal we cannot use eigenvalues to bound the norm since in general we only have

$$\sup_n |\lambda_n(A)| \leq \|A\|.$$

In Figure 4 we demonstrate the difference of $|\lambda_{max}| := \sup_n |\lambda_n|$ and $\|S\|_{L^2(\Gamma)}$ for the single layer potential on the boundary Γ for the crescent domain from Figure 1. Numerically, for the eigenvalues we obtain a decay of approximately $O(k^{-0.77})$, while the norm decays much slower with a rate of about $O(k^{-1/2})$. This effect is a consequence of nonnormality. In Betcke *et al.* (2011) various norm estimates are summarised; for general Lipschitz domains the estimate $\|S\|_{L^2(\Gamma)} = O(k^{-1/2})$ holds.

There is an interesting relationship between the norm of an operator and its numerical range. Let $r(A) := \sup\{|z| : z \in W(A)\}$ be the *numerical radius* of A . Then it holds that $r(A) \leq \|A\| \leq 2r(A)$ (Gustafson & Rao, 1997). Hence, k -dependent norm bounds immediately are also bounds for the extent of the numerical range of the operators S , K and T .

2.3 Coercivity Constants

An operator A acting on a Hilbert space \mathcal{H} is coercive if there exists $\gamma > 0$, such that

$$\gamma \|u\|^2 \leq |\langle Au, u \rangle|$$

for all $u \in \mathcal{H}$. Consider the abstract variational problem

$$\text{Find } u \in \mathcal{H}, \text{ such that } \langle Au, v \rangle = f(v), \quad \forall v \in \mathcal{H}, \quad (2.1)$$

where $f \in \mathcal{H}'$. Let $\mathcal{V}^{(h)}$ be a finite dimensional subspace of \mathcal{H} and denote by $u^{(h)}$ the solution of the variational problem (2.1) restricted to $\mathcal{V}^{(h)}$. Then if A is coercive, by Céa's Lemma, (Cea, 1964), we have

$$\|u - u^{(h)}\| \leq C \min_{v \in \mathcal{V}^{(h)}} \|u^{(h)} - v\|, \quad (2.2)$$

where $C = \|A\|/\gamma$.

The analysis of coercivity constants for the combined potential operator A_η is the subject of current research (see Spence *et al.* (2011)), and the main motivation for this is the following. Recall that establishing coercivity means that the quasi-optimality estimate (2.2) holds for *any* finite dimensional subspace $\mathcal{V}^{(h)}$. Recently much research effort has gone into designing k -dependent subspaces that accurately approximate the solution to scattering problems where k is large, even for relatively small subspace dimension. (Recall that the standard piecewise polynomial subspaces require their dimension to grow like k^{d-1} to maintain accuracy as k increases.) An overview of these “hybrid numerical-asymptotic” methods can be found in the review by Chandler-Wilde & Graham (2009). Since these subspaces depend on k in a complicated way, the only known way of proving convergence of Galerkin methods using these subspaces is via coercivity.

A first coercivity proof for A_k in $L^2(\Gamma)$ was given in Domínguez *et al.* (2007) for the case that Γ is the unit circle or the unit sphere. By a careful analysis of the eigenvalues of A_k it was shown that $\gamma \geq 1$ for sufficiently large k . For general domains an eigenvalue analysis is not sufficient any more. This can be seen as follows. From the definition of the numerical range it follows that A_η is coercive if and only if $0 \notin \overline{W(A_\eta)}$. Furthermore, if A_k is coercive then γ is just the distance of 0 to $\overline{W(A_k)}$. In the case of the circle the combined potential operator is normal and the numerical range is just the convex hull of the eigenvalues. But this need not be the case for more general domains. However, we can still numerically study the distance of the origin to the boundary of the numerical range in order to estimate coercivity constants. This was recently done in Betcke & Spence (2011). As an example consider the numerical range plot in Figure 2 for A_k defined on the boundary of the crescent domain. From the plot it follows that A_k is coercive with a coercivity constant smaller than 0.5. It is instructive to compare the boundary of the numerical range with the eigenvalues. The boundary of $W(A_k)$ is much closer to 0 than the eigenvalues, an effect due to the nonnormality of the operator.

3. A nonnormality proof for the single layer potential operator

It was observed in (Betcke & Spence, 2011, Lemma 3.5) that A_η is normal in $L^2(\Gamma)$ if Γ is the boundary of a circle in 2-d or a sphere in 3-d. Using the same argument S , K and T are also normal on the circle and sphere. In this section we provide a partial answer to the question of whether there are any domains, other than the circle and sphere, for which the operators S , K , T and A_η are normal in the standard $L^2(\Gamma)$ inner product. By reformulating the problem of normality into one of equilibrium potentials we prove that among all bounded $C^{2,\alpha}$ domains balls are the only ones for which the operator S is normal. The main result is the following theorem.

THEOREM 3.1 Let $\Omega \subset \mathbb{R}^d$, $d = 2, 3$ be a bounded $C^{2,\alpha}$ domain with boundary Γ . Then the single layer potential S is normal if and only if Ω is a ball.

The proof that S is normal on a ball follows by diagonalisation in the Fourier-basis (2-d) and by diagonalisation in spherical harmonics (3-d), see (Betcke & Spence, 2011, Lemma 3.5) for details. To prove the converse we first show that S normal implies that the logarithmic single layer potential operator defined by

$$[\mathcal{S}_0\phi](x) := \begin{cases} -\frac{1}{2\pi} \int_\Gamma \log|x-y|\phi(y)ds(y), & \text{2-d} \\ \frac{1}{4\pi} \int_\Gamma \frac{1}{|x-y|} \phi(y)ds(y), & \text{3-d} \end{cases}$$

applied to the constant function 1 is constant inside Ω . We can then use the following result to show that Ω must be a ball.

THEOREM 3.2 (Theorem 2 and §3 of Reichel (1997)) If Ω is a $C^{2,\alpha}$ domain and $\mathcal{S}_0 1$ is constant in Ω then Ω is a ball.

This problem can be understood in terms of *equilibrium potentials* (see, e.g., (Ransford, 1995, §3.3)). Given a domain, the nontrivial measure that gives rise to a constant potential throughout the domain is called the equilibrium measure. The corresponding potential is the equilibrium potential. Theorem 3.2 states that the potential $\mathcal{S}_0 1$ is the equilibrium potential only for the case when Ω is a ball. The proof in Reichel (1997) considers the equivalent problem of showing that the following overdetermined boundary

value problem has a solution if and only if Ω is a ball:

$$\Delta w = 0 \quad \text{in } \Omega, \quad \gamma_1^+ w = -1, \quad \gamma_0^+ w = a,$$

where a is a constant and an appropriate condition at infinity is prescribed. This condition at infinity is $u \rightarrow 0$ for $d \geq 3$, and one involving layer potentials for $d = 2$ (since the single layer potential does not decay at infinity in this case).

These overdetermined problems in potential theory have a long history, beginning with the celebrated work of Serrin (1971), and are motivated by physical questions including stress in hydrodynamics.

A natural question is where the restriction that $C^{2,\alpha}$ comes from. In order to apply the appropriate theorem on the overdetermined problem, (Reichel, 1997, Theorem 1), Reichel requires that $\mathcal{S}_0 1$ be in $C^2(\overline{\Omega^+})$. Reichel uses a special case of a general regularity theorem (Gilbarg & Trudinger, 1998, Theorem 6.14) to show that if D is a $C^{2,\alpha}$ domain, $\Delta u = 0$ in D , and the boundary data is $C^{2,\alpha}$, then $u \in C^{2,\alpha}(\overline{D})$.

In order to proof Theorem 3.1 from Theorem 3.2 we need to show that if the acoustic single layer potential S is normal then $\mathcal{S}_0 1$ is constant in Ω . The next lemma is technical and gives a reformulation of the condition that S is normal.

LEMMA 3.1 Let the acoustic Green's function $g(x, y) = u(|x - y|) + iv(|x - y|)$ where u and v are real. S is normal if and only if

$$C(x, z) := \int_{\Gamma} u(|z - y|)v(|x - y|) - u(|x - y|)v(|z - y|) ds(y) = 0 \quad (3.1)$$

for all $x, z \in \Gamma$.

Proof. Given $f \in L^2(\Gamma)$

$$\begin{aligned} [(SS^* - S^*S)f](x) &= \int_{\Gamma} \int_{\Gamma} (g(x, y)\bar{g}(z, y) - \bar{g}(y, x)g(y, z))f(z) ds(z) ds(y) \\ &= \int_{\Gamma} \int_{\Gamma} 2i(u(|z - y|)v(|x - y|) - u(|x - y|)v(|z - y|))f(z) ds(z) ds(y) \\ &= 2i \int_{\Gamma} f(z)C(x, z) ds(z) \end{aligned} \quad (3.2)$$

where $x, z \in \Gamma$. Interchanging the order of integration is justified by Fubini's theorem since g is weakly singular.

Assume that S is normal, that is $SS^* - S^*S = 0$. Given any arbitrary $z' \in \Gamma$, the right hand side of (3.2) can be made arbitrarily close to $2iC(x, z')$ by choosing an appropriate $f(z)$ with support in a neighbourhood of z' . Thus if $(SS^* - S^*S)(f)(x)$ is equal to zero for all $x \in \Gamma$ and for all f then $C(x, z') = 0$ for all $x, z' \in \Gamma$. The converse is immediate. \square

We know that S is normal for Γ the circle or sphere, and we can check that $C(x, z) = 0$ in this case. Indeed, the change of variables $y = (x + z) - y'$ transforms the unit circle/sphere centered at 0 to one centred at $x + z$, and $ds(y) = ds(y')$. Making this change of variables in the second term in (3.1) we see that, since $|z - y| = |x - y'|$ and $|x - y| = |z - y'|$, the second term cancels with the first and thus $C(x, z) = 0$.

For the next lemma we note that for $d = 2$, $u(r) = -\frac{1}{4}Y_0(kr)$ and $v(r) = \frac{1}{4}J_0(kr)$, where J_ν and Y_ν are Bessel functions of the first and second kind respectively. For $d = 3$, $g(x, y) = e^{ik|x-y|}/(4\pi|x-y|)$ and thus $u(r) = (\cos kr)/(4\pi r)$ and $v(r) = (\sin kr)/(4\pi r)$.

We can now prove the following.

LEMMA 3.2 If $C(x, z) = 0$ for all $x, z \in \Gamma$ then $\mathcal{S}_0 1$ is constant in Ω .

Proof. Since $C(x, z) = 0$ for all $x, z \in \Gamma$, then of course for every fixed z $C(x, z) = 0$ for all $x \in \Gamma$. Thus, $\nabla_{\Gamma, x} C(x, z) = 0$ for all $x \in \Gamma$, where $\nabla_{\Gamma, x}$ is the surface gradient. An explicit expression for this in terms of a parametrisation of the boundary can be found in (Monk, 2003, §3.4) (Colton & Kress, 1983, §2.1) for smooth domains. Furthermore, if w is C^1 in a neighbourhood of Γ then

$$\nabla_{\Gamma, x} w(x) = \nabla w(x) - n(x) \frac{\partial w}{\partial n}(x), \quad x \in \Gamma. \quad (3.3)$$

Ideally we would like to take $\nabla_{\Gamma,x}$ under the integral sign in the definition of $C(x,z)$ (3.1) and obtain

$$\nabla_{\Gamma,x}C(x,z) = P.V. \int_{\Gamma} \left(\frac{x-y}{|x-y|} - \frac{(x-y) \cdot n(x)}{|x-y|} n(x) \right) (u(|z-y|)v'(|x-y|) - u'(|x-y|)v(|z-y|)) ds(y) \quad (3.4)$$

where we are using (3.3) to find the surface gradient of $|x-y|$ under the integral sign. However, since the integrand is weakly singular, this interchange of integration and differentiation requires some justification.

The key result is that, for $\phi \in L^2(\Gamma)$ and $k \geq 0$,

$$\nabla_{\Gamma,x}[S\phi](x) = [\nabla_{\Gamma}S\phi](x) \quad (3.5)$$

where the right hand side is the integral operator defined by

$$\begin{aligned} [\nabla_{\Gamma}S\phi](x) &= P.V. \int_{\Gamma} \nabla_{\Gamma,x}g(x,y)\phi(y)ds(y), \\ &= P.V. \int_{\Gamma} \left(\frac{x-y}{|x-y|} - \frac{(x-y) \cdot n(x)}{|x-y|} n(x) \right) (u'(|x-y|) + iv'(|x-y|))\phi(y)ds(y). \end{aligned} \quad (3.6)$$

On C^2 domains this follows from the results for Hölder continuous ϕ in (Colton & Kress, 1983, Theorem 2.17) (using the density of Hölder continuous functions in $L^2(\Gamma)$ when Γ is C^2), and this result is true even on Lipschitz domains by the harmonic analysis results surveyed in (Meyer & Coifman, 2000, Chapter 15).

The result (3.5) justifies (3.4) as follows. C , defined by (3.1), consists of two terms: the first, viewed as a function of x , is the imaginary part of the single layer potential with density $u(|z-y|)$, the second, again viewed as a function of x , is the real part of the single layer potential with density $v(|z-y|)$. Both $u(|z-y|)$ and $v(|z-y|)$ are in $L^2(\Gamma)$ and thus interchanging the differentiation and integration is justified by the imaginary and real parts respectively of (3.5).

Since $\nabla_{\Gamma,x}C(x,z) = 0$ for all $x \in \Gamma$ in particular this holds for $x = z$. Let $\chi(z) := \nabla_{\Gamma,x}C(x,z)|_{(x=z)}$. Then, for $d = 2$ the Wronskian $u(r)v'(r) - u'(r)v(r) = -1/(2\pi kr)$ and

$$\chi(z) = P.V. \int_{\Gamma} \left(\frac{z-y}{|z-y|} - \frac{(z-y) \cdot n(z)}{|z-y|} n(z) \right) \frac{-1}{2\pi k|z-y|} ds(y) = -\frac{1}{2\pi k} \nabla_{\Gamma,z} \int_{\Gamma} \log(|z-y|) ds(y), \quad d = 2,$$

(where taking the surface gradient out from under the integral sign is justified by (3.5) above with $k = 0$). For $d = 3$, the Wronskian $u(r)v'(r) - u'(r)v(r) = k/(4\pi r^2)$ and

$$\chi(z) = P.V. \int_{\Gamma} \left(\frac{z-y}{|z-y|} - \frac{(z-y) \cdot n(z)}{|z-y|} n(z) \right) \frac{k}{4\pi|z-y|^2} ds(y) = \frac{k}{4\pi} \nabla_{\Gamma,z} \int_{\Gamma} \frac{1}{|z-y|} ds(y), \quad d = 3.$$

Thus $C(x,z) = 0$ for all $x, z \in \Gamma$ implies that $\chi(z) = 0$ for all $z \in \Gamma$ and thus $[\mathcal{S}_0 1](z)$ is constant for all $z \in \Gamma$. Since $\mathcal{S}_0 1$ is harmonic, by the maximum principle it is constant in Ω . \square

By combining Lemma 3.1 and Lemma 3.2 we can apply Theorem 3.2 to show that if S is normal then Ω must be a ball. This concludes the missing direction of the proof of Theorem 3.1.

4. The spectrum on the boundary of an ellipse

4.1 Eigendecomposition in elliptic coordinates

The main ingredient for computing an eigenvalue expansion on the disk is the Graf addition formula that gives an expansion of the acoustic Green's function in a polar coordinate system. In this section we generalise these results to an ellipse by using an elliptic coordinate system and an expansion of the acoustic Green's function in elliptic coordinates. A significant difference to the disk case is that the natural parameterisation of the boundary of an ellipse in elliptic coordinates has a non-constant Jacobian. The effect is that instead of a simple eigenvalue expansion for the single layer potential S we will arrive at a generalised eigenvalue expansion of the form $SF^{-1}u_n = \lambda_n u_n$, where F is a simple scaling operator. For the case of boundary integral operators in harmonic potential theory eigenvalue expansions on the ellipse were derived

by Rodin & Steinbach (2003). For the Helmholtz case we need Mathieu functions, which we will briefly introduce here.

Let Ω be an ellipse with boundary Γ parameterised by $\gamma(v) := \begin{bmatrix} a \cosh \mu \cos v \\ a \sinh \mu \sin v \end{bmatrix}$. Define $q = \frac{1}{4}(ka)^2$. Separation of variables of the Helmholtz equation in elliptic variables leads to the standard Mathieu equation

$$N''(v) + (\lambda - 2q \cos 2v)N(v) = 0 \quad (4.1)$$

and the modified Mathieu equation

$$M''(\mu) - (\lambda - 2q \cosh 2\mu)M(\mu) = 0, \quad (4.2)$$

where λ is a separation constant. Together with suitable boundary conditions (4.1) is a Sturm-Liouville problem, where $\lambda = \lambda(q)$ is the eigenvalue parameter. Depending on the boundary conditions two sets of eigenvalues $a_n(q)$ and $b_n(q)$ are defined (NIST Digital Library, §28.2(v)). By $ce_n(v, q)$ we denote the even eigenfunctions associated with the eigenvalues $a_n(q)$ and by $se_n(v, q)$ the odd eigenfunctions associated with $b_n(q)$. (NIST Digital Library, §28.2(vi)). These form an orthogonal basis of $L^2[0, 2\pi]$. For the normalisation of the functions we use the standard choice

$$\int_0^{2\pi} ce_n^2(v, q)dv = \int_0^{2\pi} se_n^2(v, q)dv = \pi.$$

For the modified Mathieu equation (4.2) we introduce the two sets of solutions $Mc_n^{(1)}(\mu, q)$ and $Mc_n^{(2)}(\mu, q)$ associated with the eigenvalues $a_n(q)$. From these the modified Mathieu functions $Mc_n^{(3/4)}$ are defined by

$$Mc_n^{(3/4)}(\mu, q) = Mc_n^{(1)}(\mu, q) \pm iMc_n^{(2)}(\mu, q).$$

$Mc_n^{(1)}$ is the elliptic analogue of the Bessel function J_n and $Mc_n^{(2)}$ is the elliptic analogue of the Bessel function of the second kind Y_n . Indeed, it holds that (NIST Digital Library, eq. 28.20.10, 28.20.15-16)

$$Mc_n^{(3/4)}(\mu, q) \sim H_n^{(1/2)}(2q^{1/2} \cosh \mu) \text{ as } \mu \rightarrow \infty. \quad (4.3)$$

One defines the functions $Ms_n^{(1/2)}$ and $Ms_n^{(3/4)}$ associated with the eigenvalues $b_n(q)$ similarly.

Similar to the circle case, there exists an expansion of the acoustic Green's function in elliptic coordinates.

THEOREM 4.1 (Expansion of fundamental solution in Mathieu functions)

$$\begin{aligned} \frac{i}{4}H_0^{(1)}(k|x-y|) = \frac{i}{2} & \left(\sum_{m=0}^{\infty} ce_m(v, q)ce_m(v', q)Mc_m^{(1)}(\mu_{<}, q)Mc_m^{(3)}(\mu_{>}, q) \right. \\ & \left. + \sum_{m=1}^{\infty} se_m(v, q)se_m(v', q)Ms_m^{(1)}(\mu_{<}, q)Ms_m^{(3)}(\mu_{>}, q) \right) \end{aligned} \quad (4.4)$$

where $x = (a \cosh \mu \cos v, a \sinh \mu \sin v)$, $y = (a \cosh \mu' \cos v', a \sinh \mu' \sin v')$ and $\mu_{>} = \max(\mu, \mu')$, $\mu_{<} = \min(\mu, \mu')$

Proof. This formula can be found in (Morse & Feshbach, 1953a, §11.2). However, it is not derived directly, but is quoted as a special case of a general technique described in (Morse & Feshbach, 1953b, §7.2). Furthermore, the notation for Mathieu functions used in these references is not the (now fairly standard) notation used in the Digital Library of Mathematical Functions that we use here. Because of these complications we give a full derivation of this expansion in Appendix B. \square

We cannot quite proceed as in the case of the unit circle. On the ellipse the boundary measure ds is not identical to the angular measure dv since $|\gamma'(v)| \neq \text{const}$. We have to introduce an additional scaling operator F to take this into account.

THEOREM 4.2 Let Γ be the boundary of an ellipse and $x = \gamma(s)$ be its parameterisation in elliptic coordinates as defined above. Denote by μ the radius of the ellipse in elliptic coordinates associated with the parameter $q = \frac{1}{4}(ka)^2$. Define the multiplication operator $F : L^2(\Gamma) \rightarrow L^2(\Gamma)$ by $[F\phi](x) = |\gamma'(\gamma^{-1}(x))|\phi(x)$. The following eigenvalue decompositions hold.

- For the single layer potential S :

$$\begin{aligned} SF^{-1}\phi_j^{(c)} &= \lambda_j^{(S,c)}\phi_j^{(c)} \\ SF^{-1}\phi_j^{(s)} &= \lambda_j^{(S,s)}\phi_j^{(s)}, \end{aligned}$$

where

$$\lambda_j^{(S,c)} = \frac{i\pi}{2}Mc_j^{(1)}(\mu, q)Mc_j^{(3)}(\mu, q), \quad \lambda_j^{(S,s)} = \frac{i\pi}{2}Ms_j^{(1)}(\mu, q)Ms_j^{(3)}(\mu, q)$$

- For the double layer potential K :

$$\begin{aligned} K\phi_j^{(c)} &= \lambda_j^{(K,c)}\phi_j^{(c)} \\ K\phi_j^{(s)} &= \lambda_j^{(K,s)}\phi_j^{(s)}, \end{aligned}$$

where

$$\lambda_j^{(K,c)} = \frac{i\pi}{2} \left[\frac{\partial}{\partial \mu} Mc_j^{(1)}(\mu, q) \right] Mc_j^{(3)}(\mu, q), \quad \lambda_j^{(K,s)} = \frac{i\pi}{2} \left[\frac{\partial}{\partial \mu} Ms_j^{(1)}(\mu, q) \right] Ms_j^{(3)}(\mu, q).$$

- For the conjugate double layer potential T :

$$\begin{aligned} TTF^{-1}\phi_j^{(c)} &= \lambda_j^{(T,c)}\phi_j^{(c)} \\ TTF^{-1}\phi_j^{(s)} &= \lambda_j^{(T,s)}\phi_j^{(s)}, \end{aligned}$$

where

$$\lambda_j^{(T,c)} = \frac{i\pi}{2}Mc_j^{(1)}(\mu, q) \left[\frac{\partial}{\partial \mu} Mc_j^{(3)}(\mu, q) \right], \quad \lambda_j^{(T,s)} = \frac{i\pi}{2}Ms_j^{(1)}(\mu, q) \left[\frac{\partial}{\partial \mu} Ms_j^{(3)}(\mu, q) \right].$$

In all three cases the eigenfunctions $\phi_j^{(c)}$ and $\phi_j^{(s)}$ are defined as

$$\phi_j^{(c)}(x) = ce_j(\gamma^{-1}(x), q), \quad \phi_j^{(s)}(x) = se_j(\gamma^{-1}(x), q).$$

Proof. From the decomposition of the acoustic Green's function in Theorem 4.1 and the orthogonality of Mathieu functions with respect to the inner product $(\cdot, \cdot)_{L^2([0, 2\pi])}$ it follows for $x = (a \cosh \mu \cos \nu, a \sinh \mu \sin \nu)$ that

$$\begin{aligned} [SF^{-1}\phi_j^{(c)}](x) &= \int_{\Gamma} \frac{i}{4} H_0^{(1)}(k|x-y|) \frac{\phi_j^{(c)}(y)}{|\gamma'(\gamma^{-1}(y))|} ds(y) \\ &= \int_0^{2\pi} |\gamma'(t)| \frac{i}{4} H_0^{(1)}(k|x-\gamma(t)|) \frac{ce_j(t, q)}{|\gamma'(t)|} dt \\ &= \frac{i\pi}{2} Mc_j^{(1)}(\mu, q) Mc_j^{(3)}(\mu, q) ce_j(\nu, q) \\ &= \lambda_j^{(S,c)} \phi_j^{(c)}(x). \end{aligned}$$

For the double layer and conjugate double layer potential we first note that $\frac{\partial}{\partial n(x)} = \frac{1}{|\gamma'(\gamma^{-1}(x))|} \frac{\partial}{\partial \mu}$ for $x \in \Gamma$ since elliptic coordinates are an orthogonal coordinate system. Let $\tilde{x} \in \Omega^+$ have the representation $(\tilde{\mu}, \tilde{\nu})$ in elliptic coordinates. It follows that

$$\begin{aligned} [\mathcal{K}\phi_j^{(c)}](\tilde{x}) &= \int_{\Gamma} \frac{\partial}{\partial n(y)} \frac{i}{4} H_0^{(1)}(k|\tilde{x}-y|) ce_j(\gamma^{-1}(y), q) ds(y) \\ &= \frac{i\pi}{2} \left[\frac{\partial}{\partial \mu} Mc_j^{(1)}(\mu, q) \right] Mc_j^{(3)}(\tilde{\mu}, q) ce_j(\tilde{\nu}, q). \end{aligned} \tag{4.5}$$

Since $Ku = \gamma_0^+ \mathcal{K}$ for $u \in L^2(\Gamma)$ the result follows for $\tilde{\mu} \rightarrow \mu$. For the conjugate double layer potential we note that

$$[\mathcal{S}F^{-1}\phi_j^{(c)}](\tilde{x}) = \frac{i\pi}{2} Mc_j^{(1)}(\mu, q) Mc_j^{(3)}(\tilde{\mu}, q) ce_j(\nu, q),$$

giving

$$[FTF^{-1}\phi_j^{(c)}](x) = [F\gamma_1^+ \mathcal{S}F^{-1}\phi_j^{(c)}](x) = \frac{i\pi}{2} Mc_j^{(1)}(\mu, q) \left[\frac{\partial}{\partial \mu} Mc_j^{(3)}(\mu, q) \right] ce_j(\nu, q)$$

for $x \in \Gamma$. The calculations for $\phi_j^{(s)}$ are similar.

□

The operator F is positive definite with bounded inverse on $L^2(\Gamma)$. We can therefore define the inner product

$$(u, v)_{L^2(F^{-1}, \Gamma)} := \int_{\Gamma} (F^{-1}u)(y) \bar{v}(y) ds(y)$$

and associated norm

$$\|v\|_{L^2(F^{-1}, \Gamma)} := \left(\int_{\Gamma} (F^{-1}v)(y) \bar{v}(y) ds(y) \right)^{1/2}$$

for all v in $L^2(\Gamma)$. Since $a \sinh \mu \leq |\gamma'(t)| \leq a \cosh \mu$ for all $t \in \mathbb{R}$ it follows that

$$\frac{1}{\sqrt{a \cosh \mu}} \|\cdot\|_{L^2(\Gamma)} \leq \|\cdot\|_{L^2(F^{-1}, \Gamma)} \leq \frac{1}{\sqrt{a \sinh \mu}} \|\cdot\|_{L^2(\Gamma)}.$$

We can now proof a simple normality result with respect to this scaled inner product.

THEOREM 4.3 The operators SF^{-1} , K and FTF^{-1} are normal with respect to the inner product $(u, v)_{L^2(F^{-1}, \Gamma)}$.

Proof. For any two functions $u, v \in L^2(\Gamma)$ we have

$$(u, v)_{L^2(F^{-1}, \Gamma)} = \int_{\Gamma} (F^{-1}u) \bar{v} ds = \int_0^{2\pi} u(\gamma(t)) \overline{v(\gamma(t))} dt.$$

Hence, from the orthogonality of the Mathieu functions ce_j and se_j with respect to the standard $L^2([0, 2\pi])$ inner product it follows that the functions $\phi_j^{(c)}$ and $\phi_j^{(s)}$ are orthogonal with respect to the inner product $(\cdot, \cdot)_{L^2(F^{-1}, \Gamma)}$. Together with Theorem 4.2 it therefore follows that the operators SF^{-1} , K and FTF^{-1} diagonalize in a unitary basis in $L^2(F^{-1}, \Gamma)$ proving normality. □

For the single layer potential a particularly nice relationship follows immediately.

COROLLARY 4.1 It holds that $SF^{-1}S^* = S^*F^{-1}S$.

Proof. The adjoint \tilde{S}^{*F} of $\tilde{S} := SF^{-1}$ in $L^2(F^{-1}, \Gamma)$ is given by $\tilde{S}^{*F} := S^*F^{-1}$ since

$$(\tilde{S}f, g)_{L^2(F^{-1}, \Gamma)} = (F^{-1}SF^{-1}f, g) = (F^{-1}f, S^*F^{-1}g) = (f, \tilde{S}^{*F}g)_{L^2(F^{-1}, \Gamma)}$$

for all $f, g \in L^2(F^{-1}, \Gamma)$. From the normality of \tilde{S} in $L^2(F^{-1}, \Gamma)$ it follows that \tilde{S} commutes with its adjoint \tilde{S}^{*F} . Hence, $\tilde{S}\tilde{S}^{*F} = \tilde{S}^{*F}\tilde{S}$ and therefore

$$SF^{-1}S^*F^{-1} = S^*F^{-1}SF^{-1},$$

from which the result follows by multiplying from the right with F . □

With $\tilde{\phi}_j^{(\tau)} := F^{-1}\phi_j^{(\tau)}$, $\tau = c, s$ we can equivalently formulate the result of Theorem 4.2 as

$$S\tilde{\phi}_j^{(\tau)} = \lambda_j^{(S, \tau)} F\tilde{\phi}_j^{(\tau)}, K\phi_j^{(\tau)} = \lambda_j^{(K, \tau)} \phi_j^{(\tau)}, T\tilde{\phi}_j^{(\tau)} = \lambda_j^{(T, \tau)} \tilde{\phi}_j^{(\tau)}.$$

Hence, the single layer potential S admits a generalized eigenvalue decompositions with eigenfunctions $\tilde{\phi}_j^{(\tau)}$ that are orthogonal in the $L^2(F, \Gamma)$ inner product. The operator K admits a standard eigenvalue decomposition with eigenfunctions that are orthogonal in $L^2(F^{-1}, \Gamma)$ and T admits a standard eigenvalue decomposition with eigenfunctions that are orthogonal in $L^2(F, \Gamma)$. It follows immediately that K and T are normal operators in $L^2(F^{-1}, \Gamma)$ and $L^2(F, \Gamma)$, respectively while the generalized pencil (S, F) has the same spectrum as the normal operator SF^{-1} in $L^2(F^{-1}, \Gamma)$.

4.2 A numerical example

To illustrate the results of this section we compare the spectra and numerical ranges of the operators S , K and T in the standard $L^2(\Gamma)$ inner product on the ellipse with the spectra and numerical ranges of the operators SF^{-1} , K and FTF^{-1} in the $L^2(F^{-1}, \Gamma)$ inner product. The example domain is an ellipse with boundary defined by the parameters $a = 1$ and $\mu = 0.3$ in elliptic coordinates. The wavenumber is $k = 10$. The results are presented in Figure 5. It is clearly visible that the standard operators S , K and T are not normal in $L^2(\Gamma)$ since the numerical ranges are not the closed convex hulls of the spectra of the operators (left plots in Figure 5). However, after scaling and changing the inner product up to the accuracy of the plotting scale the numerical ranges are the closed convex hulls of the spectra of SF^{-1} , K and FTF^{-1} (right plots in Figure 5). Note that the eigenvalues of SF^{-1} are certainly not identical to those of S , while for K and S the rescaling from Theorem 4.2 does not change the eigenvalues.

5. General smooth domains

Since the technique used for the eigenvalue decomposition of the previous section (relying on the decomposition of the Green's function in an orthogonal coordinate system) is only applicable to a very restrictive class of domains, we derive in this section approximate decompositions for general smooth domains. The idea is to use the well-known result that the boundary integral operators S , K and T on the boundary of a sufficiently smooth domain can be represented as compact perturbations of the corresponding operators on the unit circle (Sloan, 1992). For the purpose of this section we will identify \mathbb{R}^2 with the complex plane \mathbb{C} . We also note that all the analysis of this section is for a fixed wavenumber k .

Let Ω be a bounded domain with analytic boundary Γ . Denote by $\gamma_c(z)$ a conformal map from a neighborhood of the unit circle $C \subset \mathbb{C}$ into a neighborhood of Γ , such that $\gamma_c(e^{i\theta}) \in \Gamma$ for $\theta \in [0, 2\pi]$. We first consider the case of the acoustic single layer potential. For $\phi \in L^2(\Gamma)$ we have

$$[S\phi](x) = \int_{\Gamma} g(x, y)\phi(y)ds(y).$$

with the Green's function $g(x, y) := \frac{i}{4}H_0^{(1)}(k|x - y|)$.

By a change of variables onto the unit circle C we obtain

$$[S\phi](x) = \int_C g(\gamma_c(w), \gamma_c(v))\phi(\gamma_c(v))|\gamma_c'(v)|ds(v),$$

where $x = \gamma_c(w)$ and $y = \gamma_c(v)$. Similar to Section 4 we remove the scaling $|\gamma_c'(v)|$ by defining the multiplication operator F by $[F\phi](x) := |\gamma_c'(w)|\phi(x)$. If we denote the single layer potential operator on the unit circle as S_C and let $\phi^{(c)}(v) := \phi(\gamma_c(v)) \in L^2(C)$ we obtain


$$[(SF^{-1} - S_C)\phi^{(c)}](w) = \int_C (g(\gamma_c(w), \gamma_c(v)) - g(w, v))\phi^{(c)}(v)ds(v).$$

Using (NIST Digital Library, eq. 10.4.3, 10.8.2) $g(x, y)$ has the representation


$$\begin{aligned} g(x, y) &= \frac{i}{4}J_0(k|x - y|) - \frac{1}{4}Y_0(k|x - y|) \\ &= -\frac{1}{2\pi} \log(k|x - y|)J_0(k|x - y|) + h(x, y), \end{aligned}$$

where


$$\begin{aligned} h(x, y) &:= \frac{i}{4}J_0(k|x - y|) - \frac{1}{2\pi}(\gamma - \log 2)J_0(k|x - y|) \\ &\quad - \frac{1}{2\pi} \left(\frac{\frac{(k|x - y|)^2}{4}}{(1!)^2} - \left(1 + \frac{1}{2}\right) \frac{\left(\frac{(k|x - y|)^2}{4}\right)^2}{(2!)^2} + \left(1 + \frac{1}{2} + \frac{1}{3}\right) \frac{\left(\frac{(k|x - y|)^2}{4}\right)^3}{(3!)^2} - \dots \right). \end{aligned}$$



(a) Single Layer Potential



(b) Double Layer Potential



(c) Conjugate Double Layer Potential

FIG. 5: Eigenvalues and numerical ranges of the operators S , K and T in $L^2(\Gamma)$ (left plots) and the corresponding eigenvalues and numerical ranges of the operators SF^{-1} , K and FTF^{-1} in $L^2(F^{-1}, \Gamma)$ (right plots). The domain is an ellipse with $a = 1$ and $\mu = 0.3$. The wavenumber is $k = 10$.

Here, γ is Euler's constant (NIST Digital Library, eq. 5.2.3). The function $J_n(z)$ has the expansion (NIST Digital Library, eq. 10.2.2)

$$J_n(z) = \left(\frac{z}{2}\right)^n \sum_{j=0}^{\infty} (-1)^j \frac{\left(\frac{z^2}{4}\right)^j}{j!(n+j)!}. \quad (5.1)$$

It follows that $h(x, y)$ is real analytic in x and y . We can now calculate

$$\begin{aligned} s(w, v) &:= g(\gamma_c(w), \gamma_c(v)) - g(w, v) \\ &= -\frac{1}{2\pi} (\log(k|\gamma_c(w) - \gamma_c(v)|)J_0(k|\gamma_c(w) - \gamma_c(v)|) - \log(k|w - v|)J_0(k|w - v|)) \\ &\quad + h(\gamma_c(w), \gamma_c(v)) - h(w, v). \end{aligned} \quad (5.2)$$

Together with (5.1) we obtain

$$\begin{aligned} s(w, v) &= -\frac{1}{2\pi} \sum_{j=0}^{\infty} \frac{\left(-\frac{k^2}{4}\right)^j}{(j!)^2} [|\gamma_c(w) - \gamma_c(v)|^{2j} \log(k|\gamma_c(w) - \gamma_c(v)|) - |w - v|^{2j} \log(k|w - v|)] \\ &\quad + h(\gamma_c(w), \gamma_c(v)) - h(w, v) \\ &=: -\frac{1}{2\pi} \sum_{j=0}^{\infty} \frac{\left(-\frac{k^2}{4}\right)^j}{(j!)^2} \ell_j(w, v) + h(\gamma_c(w), \gamma_c(v)) - h(w, v) \end{aligned}$$

Define the difference quotient $q(w, v) := \frac{|\gamma_c(w) - \gamma_c(v)|}{|w - v|}$. Since γ_c is holomorphic, q is real analytic with respect to w and v with a removable singularity at $w = v$. We can reformulate

$$\log(k|\gamma_c(w) - \gamma_c(v)|) = \log(k|w - v|) + \log q(w, v),$$

where $\log q(w, v)$ is real analytic with respect to w and v with limit value $\log |\gamma'(w)|$ for $w = v$ (note that $\gamma'(w) \neq 0$ for $w \in C$). We obtain

$$\begin{aligned} \ell_j(w, v) &= |w - v|^{2j} \log(|w - v|) [q(w, v)^{2j} - 1] + |\gamma_c(w) - \gamma_c(v)|^{2j} \log q(w, v) \\ &\quad + |w - v|^{2j} [q(w, v)^{2j} - 1] \log k \end{aligned} \quad (5.3)$$

For $j = 0$ the singular term in ℓ_0 disappears. For $j > 0$ the ℓ_j are continuous in w and v . We summarize the results in the following lemma.

LEMMA 5.1 Similar to Section 4 define the multiplication operator F by $[F\phi](x) = |\gamma'_c(w)|\phi(x)$, where $x = \gamma_c(w)$. Let S be the single layer potential operator defined in Section 1 on the boundary Γ , and let S_C be the corresponding operator defined on the unit circle Γ_C . Define $\phi^{(c)}(x) = \phi(\gamma_c(x))$ for $\phi \in L^2(\Gamma)$. Then it holds that

$$[SF^{-1}\phi](x) = [S_C\phi^{(c)}](w) + [\hat{S}\phi^{(c)}](w),$$

where

$$[\hat{S}\phi^{(c)}](w) = \int_{\Gamma_C} s(w, v)\phi^{(c)}(v)ds(v)$$

\hat{S} is a compact operator on $L^2(\Gamma)$ with continuous kernel $s(w, v)$.

For $x = \gamma_c(w)$ with $w = e^{i\theta}$ for some $\theta \in [0, 2\pi)$ consider the function $u_n(x) := u_n^{(c)}(w) \in L^2(\Gamma)$ with $u_n^{(c)}(w) = w^n$, which is the n th Fourier mode transplanted to Γ and let F be the multiplication operator defined in Lemma 5.1. Then together with Theorem 1.1 we have

$$[SF^{-1}u_n](x) = [S_Cu_n^{(c)}](w) + [\hat{S}u_n^{(c)}](w) = \lambda_n^{(S)}u_n(x) + [\hat{S}u_n^{(c)}](w).$$

We now estimate the remainder term $\hat{S}u_n^{(c)}$. It has the form

$$[\hat{S}u_n^{(c)}](w) = \int_0^{2\pi} s(w, v(\theta)) e^{in\theta} d\theta,$$

where $v(\theta) = [\cos \theta, \sin \theta]^T$ is a parameterisation of the unit circle. Hence, $\hat{S}u_n^{(c)}$ is the $-n$ th Fourier mode of $s(w, v(\theta))$ with respect to θ . The decay of the Fourier coefficients is determined by the smoothness of s .

LEMMA 5.2 The Fourier coefficients $\hat{s}_n(w)$ of $s(w, v(\theta))$ with respect to θ satisfy $\hat{s}_n(w) = \mathcal{O}(|n|^{-3})$ where the omitted constant is independent of $w \in C$, where C is the unit circle.

Proof. From (5.3) it follows that

$$s(w, v(\theta)) = \frac{1}{2\pi} \left(\frac{k^2}{4} \right) |w - v(\theta)|^2 \log(|w - v(\theta)|) [q(w, v(\theta))^2 - 1] + \dots, \quad (5.4)$$

where we have written down only the most singular term. Since w and $v(\theta)$ both are variables on the unit circle we have that

$$|w - v(\theta)|^2 = 2 - 2\cos(\theta - \theta_w) \quad (5.5)$$

where $w = e^{i\theta_w}$ in complex coordinates. Thus

$$s(w, v(\theta)) = \frac{1}{2\pi} \left(\frac{k^2}{4} \right) (1 - \cos(\theta - \theta_w)) \log(2 - 2\cos(\theta - \theta_w)) [q(w, v(\theta))^2 - 1] + \dots \quad (5.6)$$

Differentiating with respect to θ gives

$$\begin{aligned} \frac{\partial s(w, v(\theta))}{\partial \theta} &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) [\sin(\theta - \theta_w) \log(2 - 2\cos(\theta - \theta_w)) + \sin(\theta - \theta_w)] [q(w, v(\theta))^2 - 1] \\ &+ \dots \\ \frac{\partial^2 s(w, v(\theta))}{\partial \theta^2} &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) [\cos(\theta - \theta_w) \log(2 - 2\cos(\theta - \theta_w)) + 1 + 2\cos(\theta - \theta_w)] [q(w, v(\theta))^2 - 1] \\ &+ \dots \end{aligned}$$

Using the Taylor expansion of γ_c we have

$$|\gamma_c(w) - \gamma_c(v)|^2 = |\gamma'_c(w)(v - w) + \mathcal{O}(|v - w|^2)|^2 = |\gamma'_c(w)|^2 |v - w|^2 + \mathcal{O}(|v - w|^3)$$

and therefore $q^2(w, v) = |\gamma'_c(w)|^2 + \mathcal{O}(|v - w|)$ as $v \rightarrow w$. Using this and (5.5) it follows that

$$\frac{\partial^2 s(w, v(\theta))}{\partial \theta^2} = \frac{1}{\pi} \left(\frac{k^2}{4} \right) \log |w - v(\theta)| [|\gamma'_c(w)|^2 - 1] + \dots, \quad (5.7)$$

where we have omitted terms of order $\mathcal{O}(|w - v(\theta)|^j \log |w - v(\theta)|)$ for $j > 0$ and analytic terms. For the Fourier-coefficients \hat{s}_n we now have, using integration by parts,

$$\hat{s}_{-n}(w) = \left(\frac{i}{n} \right)^2 \int_0^{2\pi} \frac{\partial^2 s(w, v(\theta))}{\partial \theta^2} e^{in\theta} d\theta.$$

By (5.7) it follows that up to higher order terms with faster decaying Fourier coefficients the integrand is a multiple of the $-n$ th Fourier coefficient of the kernel of the Laplace single layer potential on the unit circle, which decays like $\mathcal{O}(|n|^{-1})$ independently of $w \in C$ (see Appendix A). It follows that $\hat{s}_n(w) = \mathcal{O}(|n|^{-3})$. \square

We can now prove the following Theorem.

THEOREM 5.1 Using the notation of Lemma 5.1, the scaled Single Layer Potential SF^{-1} approximately diagonalizes in a scaled Fourier basis $u_n(x) = w^n$, where $x = \gamma_c(w)$, such that

$$SF^{-1}u_n = \lambda_n^{(S)} u_n + \mathcal{O}(|n|^{-3}), \quad (5.8)$$

where the eigenvalues $\lambda_n^{(S)}$ are the circle eigenvalues defined in Theorem 1.1.

Proof. The proof is a direct consequence of Lemma 5.1 and Lemma 5.2 since

$$[\hat{S}u_n^{(c)}](w) = \hat{s}_{-n}(w).$$

□

We note that this result does not necessarily imply that the eigenvalues on general domains converge to the eigenvalues on the unit circle for $n \rightarrow \infty$. The reason is that for nonnormal operators a residual of size ε for an approximate eigenpair only implies that the corresponding approximate eigenvalue is in the ε -pseudospectrum (see the third characterisation of pseudospectra in Definition 2.1). But this can be a large set depending on the nonnormality of the operator. It is nevertheless useful to compare the result in Theorem 5.1 to the decay of the eigenvalues $\lambda_n^{(S)}$ on the unit circle. In Appendix A it is shown that $\lambda_n \sim \frac{1}{2|n|}$. Hence, the cubic decay of the residual in Theorem 5.1 is nontrivial.

For the double layer potential a change of variables gives

$$\begin{aligned} [K\phi](x) &= \frac{\phi(x)}{2} + \int_{\Gamma} \frac{\partial}{\partial n(y)} \frac{i}{4} H_0^{(1)}(k|x-y|) \phi(y) ds(y) \\ &= \frac{\phi^{(c)}(w)}{2} + \int_C \frac{\partial}{\partial n(v)} \frac{i}{4} H_0^{(1)}(k|\gamma_c(w) - \gamma_c(v)|) \phi^{(c)}(v) ds(v) \end{aligned}$$

since γ_c is a conformal map and therefore $\frac{\partial}{\partial n(y)} = \frac{1}{|\gamma'(v)|} \frac{\partial}{\partial n(v)}$. Here, $x = \gamma_c(w)$, $y = \gamma_c(v)$ and $\phi^{(c)}(w) = \phi(\gamma_c(w))$. Let K_C be the double layer potential operator on the unit circle as defined in Section 1. We have

$$[K\phi](x) - [K_C\phi^{(c)}](w) = \int_C \frac{\partial}{\partial n(v)} s(w, v) \phi^{(c)}(v) ds(v),$$

where s is defined in (5.2).

From (5.4) we have

$$\begin{aligned} \frac{\partial}{\partial n(v)} s(w, v(\theta)) &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) 2(v(\theta) - w) \cdot n(\theta) \log(|w - v(\theta)|) [q(w, v(\theta))^2 - 1] \\ &+ \frac{1}{2\pi} \left(\frac{k^2}{4} \right) |w - v(\theta)|^2 \log(|w - v(\theta)|) \frac{\partial}{\partial n(v)} [q(w, v(\theta))^2] + \dots \\ &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) [1 - \cos(\theta - \theta_w)] \log(2 - 2\cos(\theta - \theta_w)) [q(w, v(\theta))^2 - 1] \\ &+ \frac{1}{2\pi} \left(\frac{k^2}{4} \right) [1 - \cos(\theta - \theta_w)] \log(2 - 2\cos(\theta - \theta_w)) \frac{\partial}{\partial n(v)} [q(w, v(\theta))^2] + \dots \\ &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) [1 - \cos(\theta - \theta_w)] \log(2 - 2\cos(\theta - \theta_w)) \\ &\times \left[q(w, v(\theta))^2 - 1 + \frac{\partial}{\partial n(v)} [q(w, v(\theta))^2] \right] + \dots, \end{aligned} \tag{5.9}$$

where $n(\theta) = v(\theta) = [\cos \theta, \sin \theta]^T$. Again, we have only written down the most singular term.

The singular factor in the leading term is identical to that of (5.6). Hence, we can proceed as in the proof of the single layer potential case and obtain

$$Ku_n = \lambda_n^{(K)} u_n + \mathcal{O}(|n|^{-3}), \tag{5.10}$$

In Appendix A we show that the eigenvalues of the double layer potential on the unit disk, $\lambda_n^{(K)} \sim \frac{1}{2} + \left(\frac{k^2}{4} \right) |n|^{-3}$. Hence, the approximate decomposition for the double layer potential in (5.10) does not seem to be as strong as the corresponding statement for the single layer potential since the eigenvalues for the double layer potential on the circle also converge cubically.

For the conjugate double layer potential a change of variables gives

$$[FTF^{-1}\phi](x) = \frac{\phi^{(c)}(w)}{2} + \int_C \frac{\partial}{\partial n(w)} \frac{i}{4} H_0^{(1)}(k|\gamma_c(w) - \gamma_c(v)|) \phi^{(c)}(v) ds(v)$$

since $\frac{\partial}{\partial n(x)} = \frac{1}{|\gamma_c'(w)|} \frac{\partial}{\partial n(w)}$ for $x = \gamma_c(w)$. Note that on the circle the normal direction n_w along w is just $n_w := w$. Proceeding as for the double layer potential we need to consider derivatives with respect to θ of

$$\begin{aligned} \frac{\partial}{\partial n(w)} s(w, v(\theta)) &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) 2(w - v(\theta)) \cdot n_w \log(|w - v(\theta)|) [q(w, v(\theta))^2 - 1] \\ &+ \frac{1}{2\pi} \left(\frac{k^2}{4} \right) |w - v(\theta)|^2 \log(|w - v(\theta)|) \frac{\partial}{\partial n(w)} [q(w, v(\theta))^2] + \dots \\ &= \frac{1}{2\pi} \left(\frac{k^2}{4} \right) [1 - \cos(\theta - \theta_w)] \log(2 - 2\cos(\theta - \theta_w)) \\ &\times \left[q(w, v(\theta))^2 - 1 + \frac{\partial}{\partial n(w)} [q(w, v(\theta))^2] \right] + \dots \end{aligned}$$

The singular factor in the leading order term is identical to the previous two cases. Hence, we can proceed as before and obtain

$$FTF^{-1}u_n = \lambda_n^{(T)} u_n + \mathcal{O}(|n|^{-3}). \quad (5.11)$$

Similar to the case of the double layer potential, the decay of the eigenvalues for the conjugate double layer potential on the unit circle is $\lambda_n^{(T)} \sim -\frac{1}{2} + \left(\frac{k^2}{4}\right) |n|^{-3}$. Hence, the decay of the residual is no faster than the decay of the eigenvalues on the unit circle.

We conclude this section with a simple numerical example. The domain is an ellipse defined by $\mu = 0.3$ and $a = 1$. We discretise the operators SF^{-1} , K , and FTF^{-1} using a Galerkin discretisation in $L^2(F^{-1}, \Gamma)$; this is the appropriate space for the scaled Fourier modes to be orthogonal. After discretisation we obtain the finite dimensional eigenvalue problem $\mathbf{B}x = \lambda \mathbf{M}^{(F^{-1})}x$, where $\mathbf{M}^{(F^{-1})}$ is the mass matrix in the $L^2(F^{-1}, \Gamma)$ inner product and \mathbf{B} is the discretisation of either of SF^{-1} , K , or FTF^{-1} in this inner product. Let \mathbf{U} be a matrix of discretised Fourier modes in the given finite dimensional basis. Here, we choose Fourier modes from $n = -200$ to $n = 200$. According to the results of this section \mathbf{U} approximately diagonalises the pencil $(\mathbf{B}, \mathbf{M}^{(F^{-1})})$, in the sense that for the columns \mathbf{R}_n of

$$\mathbf{R} := \mathbf{U}^H \mathbf{B} \mathbf{U} - \mathbf{U}^H \mathbf{M}^{(F^{-1})} \mathbf{U} \Lambda,$$

we have

$$\mathbf{R}_n = \mathcal{O}(|n|^{-3}),$$

where Λ is the diagonal matrix of circle eigenvalues of either the single, double, or conjugate double layer potential associated with the Fourier modes in \mathbf{U} . The decay of the columns of \mathbf{R}_n for the wavenumber $k = 10$ is shown in Figure 6. It is interesting to note that the convergence initially stagnates for $|n| < 10$. It seems that the approximation results of this section only become sharp for the asymptotic regime as $|n| > k$.

Finally we remark that repeating the above analysis for the Laplace single, double, and conjugate double layer potentials shows exponential decay of the residual when approximating eigenpairs on general domains with eigenpairs on the circle (as opposed to the algebraic decay above for the Helmholtz case). The reason for this is that the difference between the Laplace Green's functions

$$\frac{-1}{2\pi} \log |\gamma_c(x) - \gamma_c(y)| - \left(-\frac{1}{2\pi} \right) \log |x - y| = -\frac{1}{2\pi} \log \frac{|\gamma_c(x) - \gamma_c(y)|}{|x - y|}$$

has a removable singularity for $x = y$ and is therefore analytic in x and y , giving exponential decay of the corresponding Fourier coefficients. (Recall that in the Helmholtz case the difference $s(w, v)$ is not analytic - see (5.4) - hence the algebraic decay.)




FIG. 6: Cubic decay of the residual for the single layer, double layer, and conjugate double layer potential.

6. Spectral decompositions and nonnormality of the combined potential operator

In Section 1 we introduced the combined potential operator $A_\eta : L^2(\Gamma) \rightarrow L^2(\Gamma)$. In this section we will demonstrate how this operator can be modified to also fit into the framework of the decompositions derived in Section 4 and 5 and present numerical examples that demonstrate the effect of these modifications on the normality of the operator. In order to derive the modified combined potential operator we briefly recap the derivation of the combined potential operator. Consider the problem of time-harmonic acoustic scattering from a sound-soft bounded obstacle $\Omega \subset \mathbb{R}^d$, ($d = 2, 3$) with Lipschitz boundary $\Gamma := \partial\Omega$. That is, we are looking for the solution u of the problem

$$\Delta u + k^2 u = 0 \quad \text{in } \mathbb{R}^d \setminus \overline{\Omega}, \quad (6.1)$$

$$u = 0 \quad \text{on } \Gamma, \quad (6.2)$$

$$\frac{\partial u_s}{\partial r} - iku_s = o(r^{-(d-1)/2}), \quad (6.3)$$

where $u = u_{inc} + u_s$ is the total field, u_{inc} is an entire solution of (6.1), such as an incident plane wave, u_s is the scattered field, r is the radial coordinate, and $k > 0$ is the wavenumber. With the standard free-space Green's function

$$g(x, y) = \frac{i}{4} H_0^{(1)}(k|x-y|)$$

for $x, y \in \mathbb{R}^2, x \neq y$, the solution u is given by

$$u(x) = u_{inc}(x) - \int_{\Gamma} g(x, y) \partial_n u(y) ds(y), \quad x \in \mathbb{R}^d \setminus \overline{\Omega}, \quad (6.4)$$

where $\partial_n u$ is the outward pointing normal derivative of u . Taking the normal trace γ_1^+ of (6.4) and recalling the definition of the boundary integral operator T from Section 1 gives

$$(I + T) \partial_n u = \frac{\partial}{\partial n} u_{inc}. \quad (6.5)$$

Furthermore, taking the trace γ_0^+ of (6.4) and using the boundary condition (6.2) it follows that

$$S \partial_n u = u_{inc}. \quad (6.6)$$

Multiplying (6.6) with $i\eta$ for $\eta \neq 0$ and subtracting from (6.5) gives the operator equation

$$(I + T - i\eta S) \partial_n u = \frac{\partial}{\partial n} u_{inc} - i\eta u_{inc}.$$

The standard combined operator is defined as $A_\eta := 2(I + T - i\eta S)$, where the factor 2 is frequently included so that A_η is a perturbation of the identity (as opposed to $\frac{1}{2}I$). To apply the results from Section 4 and 5 we use the multiplication operator F defined in Lemma 5.1. Multiplying (6.6) by $i\eta F^{-1}$ and subtracting from (6.5) gives

$$\frac{1}{2} A_\eta^{(F)} u_n := (I + T - i\eta F^{-1} S) u_n = \frac{\partial}{\partial n} u_{inc} - i\eta F^{-1} u_{inc}.$$

In Chandler-Wilde & Langdon (2007) it was shown that A_η is a bounded operator with bounded inverse on $L^2(\Gamma)$ for every $\eta \neq 0$ and $k > 0$. Since F is bounded with bounded inverse the same arguments show that $A_\eta^{(F)}$ is bounded with a bounded inverse on $L^2(\Gamma)$. However, by rescaling S with F^{-1} we can now give a simple approximate eigenvalue decomposition for a bounded domain Ω with analytic boundary Γ .

THEOREM 6.1 Define F as in Lemma 5.1 and let the boundary Γ be analytic. Denote by u_n the n th scaled Fourier mode as defined in Theorem 5.1. The operator $FA_\eta^{(F)} F^{-1}$ admits an approximate eigendecomposition of the form

$$FA_\eta^{(F)} F^{-1} u_n = \lambda_n^{(A)} u_n + \mathcal{O}(|n|^{-3}),$$

for any $n \neq 0$, where $\lambda_n^{(A)}$ is defined in Theorem 1.1.

Proof. We have

$$FA_\eta^{(F)}F^{-1} = 2(I + FTF^{-1} + SF^{-1}).$$

Applying (5.8) and (5.11) now gives the desired result. \square

The asymptotics in Appendix A show that the eigenvalues $\lambda_n^{(A)}$ behave asymptotically like $1 + \mathcal{O}(|n|^{-1})$. Thus, the statement of (6.1) is nontrivial as it states that the eigenpairs on the circle approximate the eigenpairs of the combined potential operator on general smooth domains up to a cubically small error.

Instead of the general approximate eigenvalue decomposition given in Theorem 6.1 we can also use the exact eigenvalue decompositions of the operators SF^{-1} and FTF^{-1} on the ellipse given in Theorem 4.2 to obtain an exact eigenvalue decomposition of the combined potential operator on ellipses; we omit the details.

We now visualise the approximate decomposition of the combined potential using two examples, an ellipse and the crescent domain from Section 2. We compare the exact and approximate eigenvalue decomposition of the combined potential operator $A_\eta^{(F)}$ for growing k . We always choose $\eta = k$, which is a standard choice for sufficiently large wavenumbers (Betcke *et al.*, 2011).

After a change of variables the eigenvalue problem considered in Theorem 6.1 becomes

$$A_\eta^{(F)}\tilde{u} = \lambda\tilde{u}, \tag{6.7}$$

where $\tilde{u} = F^{-1}u$. Theorem 5.1 showed that we have an approximate eigendecomposition if then u_n are scaled Fourier modes. These functions are orthogonal in $L^2(F^{-1}, \Gamma)$. After the change of variables $\tilde{u}_n = F^{-1}u_n$ the functions \tilde{u} are orthogonal in $L^2(F, \Gamma)$. The variational formulation of (6.7) associated with this inner product is to find nontrivial pairs $(\lambda, u) \in \mathbb{C} \times L^2(F, \Gamma)$, such that

$$(FA_\eta^{(F)}u, v)_{L^2(\Gamma)} = \lambda(Fu, v)_{L^2(\Gamma)} \quad \forall v \in L^2(F, \Gamma).$$

Let $\mathbf{A}_\eta^{(F)}x = \lambda\mathbf{M}^{(F)}x$ be the finite dimensional Galerkin discretisation in $L^2(F, \Gamma)$ associated with this problem. Let \mathbf{U} be the matrix whose columns are the coefficients of the Galerkin projections of the functions \tilde{u}_n in $L^2(F, \Gamma)$. Then from Theorem 6.1 we expect that $\mathbf{U}^H\mathbf{A}_\eta^{(F)}\mathbf{U}$ is approximately diagonalised. Also, by the $L^2(F, \Gamma)$ orthogonality of the basis functions, the matrix $\mathbf{U}^H\mathbf{M}^{(F)}\mathbf{U}$ is diagonal up to discretisation errors.

Figure 7 shows logarithmic plots of $|\mathbf{U}^H\mathbf{A}_\eta^{(F)}\mathbf{U}|$ for an ellipse with $a = 1$ and $\mu = .3$ (left plots) and the crescent domain from Section 2 (right plots). The top plots show the case $k = 10$ and the bottom plots the case $k = 50$. The color scale uses white for large values. For the ellipse the operator is almost perfectly diagonalised in a Fourier basis with a small error concentrated at the low-frequency Fourier modes. In the crescent case we still have a strong diagonal, but now the off-diagonal errors are larger than in the ellipse case. This is expected since the conformal map from the unit circle to the crescent has singularities close to the boundary of the domain.

In Figure 8 we plot the relative distance between the diagonal values $|\text{diag}(\mathbf{A}_\eta^{(F)})|/|\text{diag}(\mathbf{M}^{(F)})|$ and the associated circle eigenvalues $\lambda_n^{(A)}$ for $k = 10$ on the crescent domain. After an initial stagnation phase up to approximately $|n| = 10$ the diagonal values $|\text{diag}(\mathbf{A}_\eta^{(F)})|/|\text{diag}(\mathbf{M}^{(F)})|$ are well approximated by circle eigenvalues. Figure 9 shows the diagonal values of $|\text{diag}(\mathbf{A}_\eta^{(F)})|/|\text{diag}(\mathbf{M}^{(F)})|$ for the crescent and the absolute values of the circle eigenvalues $\lambda_n^{(A)}$ in the case $k = 10$. The eigenvalues are oscillatory for $|n| < 10$. After a short transition zone they enter the asymptotic regime for $|n| > 10$ in which the circle eigenvalues behave like $1 + \mathcal{O}(|n|^{-1})$. The numerical results indicate that the approximations via circle eigenvalues are valid in this asymptotic regime.

We already showed the pseudospectrum for the combined potential operator A_η in the standard $L^2(\Gamma)$ inner product for the crescent domain and wavenumber $k = 10$ in Figure 2. In Figure 10 we compare the




FIG. 7: Logarithmic plots of $|\mathbf{U}^H \mathbf{A}_n^{(F)} \mathbf{U}|$ for an ellipse with parameters $a = 1$, $\mu = 3$ (left) and the crescent domain from Section 2 (right). The upper plots are for $k = 10$ and the bottom plots for $k = 50$. In both cases we display results for scaled Fourier modes from $n = -200$ to $n = 200$.




FIG. 8: Relative distance of the diagonal values $|\text{diag}(\mathbf{A}_\eta^{(F)})|/|\text{diag}(\mathbf{M}^{(F)})|$ to the associated circle eigenvalues $\lambda_n^{(A)}$ on the crescent domain with wavenumber $k = 10$.




FIG. 9: Plot of $|\text{diag}(\mathbf{A}_\eta^{(F)})|/|\text{diag}(\mathbf{M}^{(F)})|$ and the absolute values of the circle eigenvalues $\lambda_n^{(A)}$ for $k = 10$.

pseudospectrum and numerical range of A_η in $L^2(\Gamma)$ (top figure) with that of $A_\eta^{(F)}$ in $L^2(F, \Gamma)$ (bottom figure) for the crescent domain and wavenumber $k = 50$. In both cases the dashed line denotes the boundary of the numerical range. The pseudospectra and numerical range of the operator $A_\eta^{(F)}$ in $L^2(F, \Gamma)$ are approximated by those of the matrix $\mathbf{C}^{-H} \mathbf{A}_\eta^{(F)} \mathbf{C}^{-1}$, where $\mathbf{C}^H \mathbf{C} = \mathbf{M}^{(F)}$ is the Cholesky decomposition of $\mathbf{M}^{(F)}$. This corresponds to changing to a unitary basis of the finite dimensional Galerkin subspace of $L^2(F, \Gamma)$.¹

Most of the eigenvalues in the pre-asymptotic regime cluster around the value 2 in the complex plane. For the scaled operator $A_\eta^{(F)}$ nonnormality seems manifest itself mostly around these eigenvalues; this can be seen, for example, in the large boundary of the 10^{-1} -pseudospectrum, and also in the fact that the 10^{-2} -pseudospectrum is visible close to 2 on this plotting scale. The large eigenvalues in the transitional regime have only small circular 10^{-1} -pseudospectra. They behave almost like eigenvalues of a normal operator. The picture looks different for the unscaled operator A_η , where the pseudospectral sets are much larger in the pre-asymptotic and transitional regime. However, in both cases the overall nonnormality seems still quite mild. Even for the pre-asymptotic eigenvalues in the cluster around 2 the eigenvalue condition number computed by `EIGTOOL` is only of the order 10 .²

Interestingly, the numerical range includes the origin for the case of $A_\eta^{(F)}$, but not for A_η . Thus, although rescaling A_η has meant that its spectrum has become closer to that of the circle (in the sense of Theorem 6.1), coercivity has been lost.

Finally we note that changing the coupling parameter η in the combined potential operator A_η from a constant to an operator has been proposed in several other contexts (and goes back at least to Panich (1965)). For example, Colton & Kress (1998) considered η as an operator in order to obtain an invertible combined potential operator for the Neumann problem on smooth domains, and Mitrea (1996) for the Dirichlet problem on Lipschitz domains. (Since both these authors considered *indirect* boundary integral equations, i.e. those not arising from Green's integral representation (6.4), their operator η appears on the right of S , whereas for *direct* boundary integral equations, if η is an operator it appears on the left, as above.) More recently Buffa & Hiptmair (2005), Engleder & Steinbach (2007), and Engleder & Steinbach (2008) considered η as various operators in order to prove quasi-optimality on Lipschitz domains of the resulting combined potential operator (with the operator acting in the trace spaces). Operator-valued η s have also been considered from a more practical point of view, with Anand *et al.* (2011) and Antoine & Darbas (2011) interested in obtaining combined potential operators that require a small number of iterations when solved with Krylov subspace methods.

7. Conclusions

The main results of this paper can be summarized as follows. In the first part of this paper we proved that among all sufficiently smooth domains (at least $C^{2,\alpha}$) the operator S is normal if and only if the domain is a ball. This shows that at least for the single layer potential the operator is in general nonnormal. We believe that similar results will hold for K , T and the combined operator A_η , but proofs for these cases are not yet available.

The next result was that on an ellipse an exact eigendecomposition can be derived for the operators S , K and T , which also implies that these operators are normal on the ellipse under a suitable scaling in a modified L^2 inner product. The main tool for this result was the decomposition of the Green's function using separation of variables in elliptical coordinates. There are eleven orthogonal coordinate systems in 3-d in which the Helmholtz equation is separable (Moon & Spencer, 1971, §1), (Morse & Feshbach, 1953b, Chapter 5), and analogous decompositions of the Green's function in each of these can be obtained using the general procedure in (Morse & Feshbach, 1953b, §7.2) (or following the essentially equivalent method

¹In the same way, to approximate the pseudospectrum and numerical range of A_η in $L^2(\Gamma)$ we use the matrix $\mathbf{C}^{-H} \mathbf{A}_\eta \mathbf{C}^{-1}$, where \mathbf{A}_η is the discretisation of A_η in $L^2(\Gamma)$ and \mathbf{C} is the Cholesky factor of \mathbf{M} , the mass matrix in the standard $L^2(\Gamma)$ inner product.

²Eigenvalue condition numbers measure the sensitivity of an eigenvalue to perturbations in a matrix A . A condition number of 10 means that a perturbation in A of size ε maximally perturbs the eigenvalue by around 10ε . A precise definition is for example given in (Demmel, 1997).




FIG. 10: Pseudospectral plots of A_η in $L^2(\Gamma)$ (top figure) and $A_\eta^{(F)}$ in $L^2(F, \Gamma)$ (bottom figure)

of Appendix B). Thus, for domains that are defined by one coordinate being constant in one of these eleven coordinate systems (such as the circle/sphere in polar coordinates and the ellipse in elliptical coordinates) it should be possible to obtain normality of S , K , and T under modified inner products.

The final result was an approximate decomposition using circle eigenvalues and scaled Fourier modes on general analytic domains. The numerical experiments in Section 6 indicate that this approximation becomes valid for scaled Fourier modes with number n , such that $|n| > k$. Furthermore, the bottom plot of Figure 10 indicates that under the correct scaling the nonnormality is mostly concentrated in the part of the spectrum associated with the first $\mathcal{O}(k)$ eigenvalues in the pre-asymptotic regime.

An interesting byproduct of the approximate decompositions in Section 6 concerns the order of the double layer potential considered as a pseudo-differential operator. It is well-known that the operator $K - \frac{1}{2}I$ in 3-d is order -1 (Hsiao & Wendland, 2008, p. 518). However, the eigenvalue asymptotics derived in Appendix A on the circle seem to imply that the order is -3 for the 2-d circle case. Equation (5.10) indicates that the order may even be -3 for general smooth domains in 2-d.³

The next step will be to investigate the behavior of these operators for iterative solvers such as GMRES (see Antoine & Darbas (2011) for a recent survey). For normal operators GMRES convergence is purely determined by the distribution of the eigenvalues. For nonnormal operators this is not sufficient and techniques such as the numerical range or pseudospectra become important for the analysis. Research into this question is currently ongoing.

Another open question is the high-frequency behavior. The approximate decompositions given in this paper for general domains are valid for sufficiently high Fourier-modes with $|n| > k$. However, for high-frequency scattering problems it is important to know what happens for k very large. The analysis presented in this paper does not provide much insight into this case and one would need to develop high-frequency approximations. Some previous work in this direction is that of Warnick & Chew (2001) who considered solving the boundary value problem of scattering by a crack using the single layer potential S . They decomposed S into a normal part and a non-normal part. The normal part is related to the geometric optics approximation (valid for large k), which can be understood as replacing the crack with an infinite line. This decomposition was obtained using the exact spectral decomposition of S for this particular domain, which in turn relied on the analogue of (1.2) in cartesian coordinates.

A. Eigenvalue decay of layer potentials on the unit circle

In this appendix we derive asymptotic expressions for the eigenvalues

$$\begin{aligned}\lambda_n^{(S)} &= \frac{\pi i}{2} H_n^{(1)}(k) J_n(k) \\ \lambda_n^{(K)} &= \frac{k\pi i}{4} H_n^{(1)}(k) (J_{n-1}(k) - J_{n+1}(k)) \\ \lambda_n^{(T)} &= \frac{k\pi i}{4} (H_{n-1}^{(1)}(k) - H_{n+1}^{(1)}(k)) J_n(k)\end{aligned}$$

of the operators S , K and T associated with the eigenfunctions $u_n = e^{in\theta}$. We first note that we can focus on the case $n > 0$ since $(-1)^n J_n(x) = J_{-n}(x)$, $(-1)^n Y_n(x) = Y_{-n}(x)$ and $(-1)^n H_n^{(1)}(x) = H_{-n}^{(1)}(x)$ (NIST Digital Library, eq. 10.4.1-2). We need the power series expressions for $J_n(z)$ and $Y_n(z)$ for fixed $z > 0$ as

³A proof of this statement was recently given to the authors by W. Wendland in a private communication: The fact that the double layer potential on the circle is of order -3 follows from the expansion (Hsiao & Wendland, 2008, (2.1.20)) (essentially the low frequency asymptotics); the result for C^∞ boundary Γ follows by mapping the circle to Γ using a C^∞ diffeomorphism.

$n \rightarrow \infty$ (NIST Digital Library, eq. 10.2.2,10.8.1).

$$J_n(z) = \left(\frac{z}{2}\right)^n \sum_{j=0}^{\infty} \frac{(-1)^j}{j!(n+j)!} \left(\frac{z^2}{4}\right)^j \quad (\text{A.1})$$

$$\begin{aligned} Y_n(z) &= -\frac{1}{\pi} \left(\frac{z}{2}\right)^{-n} \sum_{j=0}^{n-1} \frac{(n-j-1)!}{j!} \left(\frac{z}{2}\right)^{2j} + \frac{2}{\pi} \log\left(\frac{z}{2}\right) J_n(z) \\ &\quad - \frac{1}{\pi} \left(\frac{z}{2}\right)^n \sum_{j=0}^{\infty} (\psi(j+1) + \psi(n+j+1)) \frac{\left(-\frac{z^2}{4}\right)^j}{j!(n+j)!}, \end{aligned} \quad (\text{A.2})$$

where $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$. Taking the leading order term in (A.1) and using Stirling's approximation that $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ we arrive at

$$J_n(z) \sim \frac{1}{\sqrt{2\pi n}} \left(\frac{ez}{2n}\right)^n \quad (\text{A.3})$$

for large n (NIST Digital Library, 10.19.1). The function $\psi(z)$ has the asymptotic expansion $\psi(x) \sim \log z - \frac{1}{2z} - \sum_{j=1}^{\infty} \frac{B_{2j}}{2jz^{2j}}$ as $z \rightarrow \infty$ (NIST Digital Library, 5.11.2). By considering the leading order term in (A.2) It follows that for $n \rightarrow \infty$

$$Y_n(z) \sim -\sqrt{\frac{2}{\pi n}} \left(\frac{ez}{2n}\right)^{-n} \quad (\text{A.4})$$

(see also (NIST Digital Library, 10.19.2)). Since $H_n^{(1)}(z) = J_n(z) + iY_n(z)$ it follows that

$$H_n(z) \sim iY_n(z) \sim -i\sqrt{\frac{2}{\pi n}} \left(\frac{ez}{2n}\right)^{-n} \quad (\text{A.5})$$

as $n \rightarrow \infty$. Combining the asymptotic expansions for H_n and J_n we arrive at

$$\lambda_n^{(S)} \sim \frac{1}{2|n|}$$

as $|n| \rightarrow \infty$. To derive a sharp expression for $\lambda_n^{(K)}$ we also need to include the next higher leading terms in the expansions, that is

$$\begin{aligned} J_n(k) &\sim \left(\frac{k}{2}\right)^n \left[\frac{1}{n!} - \left(\frac{k^2}{4}\right) \frac{1}{(n+1)!} + \left(\frac{k^2}{4}\right)^2 \frac{1}{2(n+2)!} - \left(\frac{k^2}{4}\right)^3 \frac{1}{6(n+3)!} \right], \\ H_n^{(1)}(k) &\sim -\frac{i}{\pi} \left(\frac{k}{2}\right)^{-n} \left[(n-1)! + (n-2)! \left(\frac{k^2}{4}\right) + \frac{(n-3)!}{2} \left(\frac{k^2}{4}\right)^2 + \frac{(n-4)!}{6} \left(\frac{k^2}{4}\right)^3 \right]. \end{aligned}$$

We obtain

$$\begin{aligned} \lambda_n^{(K)} &= \frac{k\pi i}{4} H_n^{(1)}(k) (J_{n-1}(k) - J_{n+1}(k)) \\ &\sim \frac{1}{2} \left[[(n-1)! + (n-2)! \left(\frac{k^2}{4}\right) + \frac{(n-3)!}{2} \left(\frac{k^2}{4}\right)^2 + \frac{(n-4)!}{6} \left(\frac{k^2}{4}\right)^3] \right. \\ &\quad \times \left. \left[\frac{1}{(n-1)!} - \left(\frac{k^2}{4}\right) \frac{1}{(n)!} + \left(\frac{k^2}{4}\right)^2 \frac{1}{2(n+1)!} - \left(\frac{k^2}{4}\right)^3 \frac{1}{6(n+2)!} \right] \right. \\ &\quad \left. - \left(\frac{k^2}{4}\right) \left[\frac{1}{(n+1)!} - \left(\frac{k^2}{4}\right) \frac{1}{(n+2)!} + \left(\frac{k^2}{4}\right)^2 \frac{1}{2(n+3)!} - \left(\frac{k^2}{4}\right)^3 \frac{1}{6(n+4)!} \right] \right] \end{aligned}$$

Multiplying and collecting terms up to order n^{-3} gives

$$\lambda_n^{(K)} \sim \frac{1}{2} + \left(\frac{k^2}{4}\right) |n|^{-3} + \mathcal{O}(|n|^{-4}).$$

For the eigenvalues $\lambda_n^{(T)}$ of the conjugate double layer potential we immediately have

$$\lambda_n^{(T)} \sim -\frac{1}{2} + \left(\frac{k^2}{4}\right) |n|^{-3} + \mathcal{O}(|n|^{-4})$$

since it holds that $\lambda_n^{(K)} - \lambda_n^{(T)} = 1$ on the unit circle (since $K - \frac{1}{2}I = T + \frac{1}{2}I$, as noted in Section 1).

Combining the results for the single and conjugate double layer potential we obtain for the combined potential operator

$$\lambda_n^{(A)} \sim 1 + \mathcal{O}(|n|^{-1})$$

It is instructive to compare these results to those for the Laplace single and double layer potential. In polar coordinates the single layer potential taken over a circle with radius ρ and evaluated at a point (r, ϕ) with $r > \rho$ is given by

$$[S_0^{(r,\rho)} u](r, \phi) = -\frac{1}{2\pi} \int_0^{2\pi} \log |re^{i\phi} - \rho e^{i\theta}| \phi(\theta) d\theta. \quad (\text{A.6})$$

Noting that $\log |z| = \text{Re}\{\log z\}$ and using the Taylor series representation of $\log(1-z)$ we have

$$\log |r^{i\phi} - \rho e^{i\theta}| = \log r - \sum_{m \neq 0} \frac{\rho^{|m|} r^{-|m|}}{2|m|} e^{im(\phi-\theta)}. \quad (\text{A.7})$$

Substituting into (A.6) together with $\phi(\theta) = u_n(\theta) := e^{in\theta}$ gives

$$[S_0^{(r,\rho)} u_0](r, \phi) = -\log r, \quad [S_0^{(r,\rho)} u_n](r, \theta) = \frac{\rho^{|n|} r^{-|n|}}{2|n|} e^{in\phi}, \quad n \neq 0.$$

In the limit $r = \rho = 1$ we obtain the eigenvalues of the standard single layer potential operator $S_0 := S_0^{(1,1)}$ on the unit circle as

$$\lambda_0 = 0, \quad \lambda_n = \frac{1}{2|n|}, \quad n \neq 0.$$

The eigenvalues of the double layer potential operator

$$[K_0 u](\theta) = \frac{1}{2} \phi(\theta) - \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial}{\partial \rho} \log |re^{i\phi} - \rho e^{i\theta}| \phi(\theta) d\theta$$

are obtained by taking the derivative with respect to ρ in (A.7), substituting this in the above equation, and then evaluating the series for $r = \rho = 1$. This results in

$$\lambda_0 = 0, \quad \lambda_n = \frac{1}{2}, \quad n \neq 0.$$

(Note that this implies that the eigenvalues of the operator $K_0 - \frac{1}{2}I$, the more-common definition of the double layer potential, are $-\frac{1}{2}$ for the 0th order Fourier mode and 0 for all other modes.) These results are consistent with the results for Helmholtz: letting $k \rightarrow 0$ in the limiting form of $\lambda_n^{(K)}$ as $n \rightarrow \infty$ above we recover $\lambda_n^{(K)} \sim \frac{1}{2}$.

B. Expansion of the Green's function in elliptic coordinates

In this section we derive the expansion of the Green's function in Mathieu functions (4.4) because of the disadvantages of the formula (Morse & Feshbach, 1953a, eq. 11.2.93) discussed in the proof of Theorem 4.1.

In order to guide the reader through the technicalities involving the Mathieu functions, we first derive the analogous expansion in polar coordinates, i.e. the Graf addition formula (1.2).

In 2-d polar co-ordinates, the equation defining the Green's function is

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial g}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 g}{\partial \theta^2} + k^2 g = -\frac{\delta(r-r')\delta(\theta-\theta')}{r} \quad (\text{A.1})$$

with periodicity in the θ variable and the radiation condition in the r variable. We choose to solve this PDE via an expansion in the angular eigenfunctions. Spectral analysis of the ODE in the θ -variable (see e.g. (Stakgold, 1967, Chapter 7)) shows that the appropriate transform in this variable is the usual Fourier series expansion. Thus, taking the finite Fourier transform of the PDE (A.1) we obtain the ODE

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \hat{g}}{\partial r} \right) + \left(k^2 - \frac{n^2}{r^2} \right) \hat{g} = -\frac{\delta(r-r')e^{-in\theta}}{r} \quad (\text{A.2})$$

where

$$\hat{g}(r, n) = \int_0^{2\pi} e^{-in\theta} g(r, \theta) d\theta$$

When solving a PDE via separation of variables, some authors (such as Morse & Feshbach (1953b, p. 825)), instead of taking the appropriate transform, expand the solution in terms of the eigenfunctions and then substitute this expansion into the PDE. This is not justified apriori, since in general the eigenfunctions are complete only in a L^2 sense and the expansion of the solution may not be differentiable; thus in some cases (for example where inhomogeneous boundary conditions are prescribed) this naive substitution fails to yield a correct expression for the solution. In the present case, however, both methods yield the same, correct, solution.

The change of variables $t = kr$ yields the ODE

$$\frac{\partial}{\partial t} \left(t \frac{\partial \hat{g}}{\partial t} \right) + \left(t - \frac{n^2}{t} \right) \hat{g} = -\delta(t-\tau)e^{-in\theta}, \quad (\text{A.3})$$

where $\tau = kr'$. We now solve this ODE: there are two linearly independent solutions, and considering the ODE over the two ranges $0 < t < \tau$ and $\tau < t < \infty$ gives rise to four arbitrary constants. These constants are fixed by the four conditions: (i) boundness of \hat{g} at $t = 0$, (ii) \hat{g} satisfying the radiation condition as $t \rightarrow \infty$, (iii) continuity of \hat{g} at τ , (iv) the jump condition

$$\tau \left[\frac{\partial \hat{g}}{\partial r} \right]_{\tau^-}^{\tau^+} = -e^{-in\theta}.$$

Since (i) the solution of the homogeneous version of (A.3) that is bounded at $t = 0$ is $J_n(t)$, (ii) the solution of the homogeneous version of (A.3) that satisfies the radiation condition as $t \rightarrow \infty$ is $H_n^{(1)}(t)$, and, regarding (iv), the Wronskian of $J_n(t)$ and $H_n^{(1)}(t)$ equals $2i/(\pi t)$, the solution of (A.3) is given by

$$\hat{g}(t, n) = \frac{i\pi}{2} e^{in\theta} J_n(t_{<}) H_n^{(1)}(t_{>}).$$

Changing variables from t to r and using the inverse finite Fourier transform we find the familiar formula (1.2) (with both sides multiplied by $i/4$). Note that, splitting the sum into sums over positive and negative integers and using $n \mapsto -n$ in the latter, the formula (1.2) can be rewritten as

$$g(r, \theta; r', \theta') = \frac{i}{2} \left[\frac{1}{2} J_0(kr_{<}) H_0^{(1)}(kr_{>}) + \sum_{n=1}^{\infty} (\cos n\theta \cos n\theta' + \sin n\theta \sin n\theta') J_n(kr_{<}) H_n^{(1)}(kr_{>}) \right] \quad (\text{A.4})$$

In this form we will be better able to see the analogy with the ellipse. Note that we could have arrived at the formula (A.4) directly if, instead of taking the finite Fourier transform, we had taken the finite sine and cosine transforms, and then used the Fourier inversion formula expressed in terms of these at the end. This is exactly analogous to what we will do in elliptic coordinates below, since in this case there is no analogue of the formula $e^{i\theta} = \cos \theta + i \sin \theta$.

In 2-d elliptic co-ordinates (with notation as in §4) the PDE defining g is

$$\frac{\partial^2 g}{\partial \mu^2} + \frac{\partial^2 g}{\partial \nu^2} + 2q(\cosh 2\mu - \cos 2\nu)g = -\delta(\mu - \mu')\delta(\nu - \nu') \quad (\text{A.5})$$

(NIST Digital Library, eq. 28.32.3) (this follows from the fact that the scale factors are $h_\mu = h_\nu = a\sqrt{(\cosh 2\mu - \cos 2\nu)}/2$ (Morse & Feshbach, 1953b, p. 503), (Morse & Feshbach, 1953a, p. 1195)). As in the polar coordinate case, we choose to expand in the ‘angular’ eigenfunctions associated with the ν variable (i.e. the eigenfunctions of (4.1)). Spectral analysis of the ν -ODE shows that the appropriate transform in this variable is

$$f(\nu) = \frac{1}{\pi} \left(\sum_{n=0}^{\infty} \hat{f}_c(n, q) ce_n(\nu, q) + \sum_{n=1}^{\infty} \hat{f}_s(n, q) se_n(\nu, q) \right) \quad (\text{A.6})$$

where

$$\hat{f}_c(n, q) = \int_0^{2\pi} f(\nu) ce_n(\nu, q) d\nu, \quad \hat{f}_s(n, q) = \int_0^{2\pi} f(\nu) se_n(\nu, q) d\nu. \quad (\text{A.7})$$

Completeness of this transform is guaranteed either by the direct method of integration in the complex plane, see e.g. (Naimark, 1967, §4), or by using results about compact operators on Hilbert spaces, see e.g. (Naimark, 1967, §5) (note that the associated Sturm-Liouville problem is regular, i.e. the simplest possible case).

The expansion (A.6) is the analogue of the usual Fourier series in the polar coordinate case (except as noted above there is now no analogue of $e^{i\theta} = \cos \theta + i \sin \theta$). Taking the two transforms (A.7) of the PDE (A.5) we find

$$\frac{\partial^2 \hat{g}_c}{\partial \mu^2} - (\lambda - 2q \cosh 2\mu) \hat{g}_c = -\delta(\mu - \mu') ce_n(\nu', q) \quad (\text{A.8})$$

and a similar equation for \hat{g}_s except with ce_n replaced by se_n . In what follows we shall only give the details of how to solve the ODE involving \hat{g}_c ; the details for \hat{g}_s are very similar. Again, the solution of (A.8) is given in terms of four constants, and these are fixed by same conditions (i)-(iii) as in the polar case, and the jump condition

$$\left[\frac{\partial \hat{g}}{\partial \mu} \right]_{\mu^-}^{\mu^+} = -ce_n(\nu', q).$$

The solutions of the homogeneous form of (A.8) are defined so that they are equal to Bessel functions in the limit $\mu \rightarrow \infty$ (NIST Digital Library, §28.20), thus $Mc_n^{(3)}$ is, by definition, the appropriate solution that satisfies the radiation condition. From the Bessel function analogy we expect $Mc_n^{(1)}$ to be the solution that is bounded at $\mu = 0$, but this is not immediate from the definition (which concerns the $\mu \rightarrow \infty$ limit). However, the expansion of the $Mc_n^{(j)}$ in terms of Bessel functions (NIST Digital Library, §28.23) show that this is indeed the case. The relevant Wronskian is (NIST Digital Library, 28.20.21)

$$W[Mc_n^{(1)}, Mc_n^{(3)}] = \frac{2i}{\pi}.$$

Solving the ODE (A.8) using these facts yields

$$\hat{g}_c(\mu, q, n) = \frac{i\pi}{2} ce_n(\nu', q) Mc_n^{(1)}(\mu_<, q) Mc_n^{(3)}(\mu_>, q),$$

and thus, using the inverse transform (A.6), we find (4.4).

As a check that (4.4) is correct we recover (A.4) from (4.4) by taking the limit $\mu \rightarrow \infty$ with $a \cosh \mu = \mathcal{O}(1)$ (thus a and $q \rightarrow 0$). In this limit, curves of constant μ (i.e. ellipses) become curves of constant r in polar coordinates (i.e. circles) where $r = a \cosh \mu$. Using the asymptotics (4.3) and the analogous asymptotics for $Mc_n^{(1)}$ (NIST Digital Library, eq. 28.20.11, 28.20.12), the Mathieu functions in (4.4) become the Hankel functions in (A.4). Using the fact that $ce_n(v, 0) = \cos nv$, $se_n(v, 0) = \sin nv$, and $ce_0(v, 0) = 1/\sqrt{2}$, the expression (4.4) becomes (A.4).

Finally we explain how (4.4) is equivalent to (Morse & Feshbach, 1953a, equation 11.2.93, p. 1421). The expression in Morse and Feshbach is

$$\begin{aligned} \frac{i}{4} H_0^{(1)}(k|x-y|) = i \left(\sum_{m=0}^{\infty} \frac{Se_m(h, \cos \theta_0) Se_m(h, \cos \theta)}{M_m^e(h)} Je_m(h, \cosh \mu_{<}) He_m(h, \cosh \mu_{>}) \right. \\ \left. + \sum_{m=1}^{\infty} \frac{So_m(h, \cos \theta_0) So_m(h, \cos \theta)}{M_m^o(h)} Jo_m(h, \cosh \mu_{<}) Ho_m(h, \cosh \mu_{>}) \right) \quad (\text{A.9}) \end{aligned}$$

where $\cos \theta$ and $\cos \theta_0$ are equivalent to our v and v' respectively, their $\cosh \mu$ is equivalent to our μ , and $h^2 = 2q$. The Se_m and So_m are analogous to our ce_m and se_m , but are normalised in a different way (involving the functions $M_m^e(h)$ and $M_m^o(h)$). However, following the discussion in Morse & Feshbach (1953a, p. 1409) we find that

$$\pi \frac{Se_m(h, \cos \theta_0) Se_m(h, \cos \theta)}{M_m^e(h)} \equiv ce_m(v, q) ce_m(v', q) \quad (\text{A.10})$$

and similarly for So_m . The Je_m, Jo_m, He_m , and Ho_m are analogous to our $Mc^{(1)}, Ms^{(1)}, Mc^{(3)}$, and $Ms^{(3)}$, but, again, are normalised differently. The asymptotics on (Morse & Feshbach, 1953a, p. 1410, p. 1589) show that

$$Je_m(h, \cosh \mu) \equiv \sqrt{\frac{\pi}{2}} Mc^{(1)}(\mu, q),$$

and similarly for Jo_m, He_m , and Ho_m , thus

$$\frac{2}{\pi} Je_m(h, \cosh \mu_{<}) He_m(h, \cosh \mu_{>}) \equiv Mc_m^{(1)}(\mu_{<}, q) Mc_m^{(3)}(\mu_{>}, q). \quad (\text{A.11})$$

Using (A.10) and (A.11) in (A.9) we recover (4.4).

REFERENCES

- ANAND, A., OVALL, J. & TURC, C. (2011) Well conditioned boundary integral equations for two-dimensional sound-hard scattering problems in domains with corners. *Journal Integral Equations and Applications*.
- ANTOINE, X. & DARBAS, M. (2011) Integral equations and iterative schemes for acoustic scattering problems. *Technical Report. Technical Report*.
- BANJAI, L. & SAUTER, S. (2007) A refined Galerkin error and stability analysis for highly indefinite variational problems. *SIAM J. Numer. Anal.*, **45**, 37–53 (electronic).
- BETCKE, T., CHANDLER-WILDE, S. N., GRAHAM, I. G., LANGDON, S. & LINDNER, M. (2011) Condition number estimates for combined potential operators in acoustics and their boundary element discretisation. *Numerical Methods for PDEs, to appear*, **27**, 31–69.
- BETCKE, T. & SPENCE, E. A. (2011) Numerical estimation of coercivity constants for boundary integral operators in acoustic scattering. *SIAM J. Numer. Anal.*, **49**, 1572–1601.
- BUFFA, A. & HIPTMAIR, R. (2005) Regularized combined field integral equations. *Numerische Mathematik*, **100**, 1–19.
- CEA, J. (1964) Approximation variationnelle des problèmes aux limites. *Annales de L'Institut Fourier*, **14**, 345–444.
- CHANDLER-WILDE, S. N. & GRAHAM, I. G. (2009) Boundary integral methods in high frequency scattering. *Highly Oscillatory Problems: Computation, Theory and Applications* (B. Engquist, T. Fokas, E. Hairer & A. Iserles eds). Cambridge University Press.
- CHANDLER-WILDE, S. N. & LANGDON, S. (2007) A Galerkin Boundary Element Method for High Frequency Scattering by Convex Polygons. *SIAM J. Numer. Anal.*, **45**, 610–640.
- COLTON, D. L. & KRESS, R. (1983) *Integral equation methods in scattering theory*. Pure and Applied Mathematics (New York). New York: John Wiley & Sons Inc., pp. xii+271. A Wiley-Interscience Publication.

- COLTON, D. & KRESS, R. (1998) *Inverse acoustic and electromagnetic scattering theory*. Springer.
- CONWAY, J. B. (1985) *A course in functional analysis*. Graduate Texts in Mathematics, vol. 96. New York: Springer-Verlag.
- DEMME, J. W. (1997) *Applied numerical linear algebra*. Philadelphia, PA: Society for Industrial and Applied Mathematics (SIAM), pp. xii+419.
- DOMÍNGUEZ, V., GRAHAM, I. G. & SMYSHLYAEV, V. P. (2007) A hybrid numerical-asymptotic boundary integral method for high-frequency acoustic scattering. *Numer. Math.*, **106**, 471–510.
- ENGLEDER, S. & STEINBACH, O. (2007) Modified boundary integral formulations for the Helmholtz equation. *Journal of Mathematical Analysis and Applications*, **331**, 396–407.
- ENGLEDER, S. & STEINBACH, O. (2008) Stabilized boundary element methods for exterior Helmholtz problems. *Numerische Mathematik*, **110**, 145–160.
- GILBARG, D. & TRUDINGER, N. (1998) *Elliptic partial differential equations of second order*, 2nd edn. Springer Verlag.
- GUSTAFSON, K. E. & RAO, D. K. M. (1997) *Numerical range*. New York: Springer-Verlag. The field of values of linear operators and matrices.
- HENRICI, P. (1986) *Applied and computational complex analysis. Vol. 3*. Pure and Applied Mathematics (New York). New York: John Wiley & Sons Inc., pp. xvi+637. Discrete Fourier analysis—Cauchy integrals—construction of conformal maps—univalent functions, A Wiley-Interscience Publication.
- HSIAO, G. C. & WENDLAND, W. L. (2008) *Boundary integral equations*. Springer Verlag.
- KRESS, R. (1985) Minimizing the condition number of boundary integral operators in acoustic and electromagnetic scattering. *The Quarterly Journal of Mechanics and Applied Mathematics*, **38**, 323.
- KRESS, R. & SPASSOV, W. T. (1983) On the condition number of boundary integral operators for the exterior Dirichlet problem for the Helmholtz equation. *Numer. Math.*, **42**, 77–95.
- MCLEAN, W. (2000) *Strongly Elliptic Systems and Boundary Integral Equations*. Cambridge University Press.
- MEYER, Y. & COIFMAN, R. (2000) *Wavelets: Calderón-Zygmund and multilinear operators*. Cambridge Studies in Advanced Mathematics. Cambridge University Press.
- MITREA, M. (1996) Boundary value problems and Hardy spaces associated to the Helmholtz equation in Lipschitz domains. *Journal of Mathematical Analysis and Applications*, **202**, 819–842.
- MONK, P. (2003) *Finite element methods for Maxwell's equations*. Oxford University Press, USA.
- MOON, P. & SPENCER, D. E. (1971) *Field theory handbook: including coordinate systems, differential equations and their solutions*, 2nd edn. Springer-Verlag.
- MORSE, P. M. C. & FESHBACH, H. (1953a) *Methods of Theoretical Physics*, vol. 2. McGraw-Hill Science/Engineering/Math.
- MORSE, P. M. C. & FESHBACH, H. (1953b) *Methods of Theoretical Physics*, vol. 1. McGraw-Hill Science/Engineering/Math.
- NAIMARK, M. A. (1967) *Linear differential operators. Part I: Elementary theory of linear differential operators*. Frederick Ungar Publishing Co, New York.
- NIST (2011). Digital Library of Mathematical Functions, <http://dlmf.nist.gov/>.
- PANICH, O. I. (1965) On the question of the solvability of exterior boundary-value problems for the wave equation and for a system of Maxwell's equations [in Russian]. *Uspekhi Mat. Nauk*, **20:1(121)**, 221–226.
- RAMM, A. G. (1973) Eigenfunction Expansion Associated with Discrete Spectrum in Problems of Diffraction. *Radiotekh Elektron+*, **18**, 496–501.
- RAMM, A. G. (1980) Nonselfadjoint operators in diffraction and scattering. *Math. Methods Appl. Sci.*, **2**, 327–346.
- RANSFORD, T. (1995) *Potential theory in the complex plane*. Cambridge University Press.
- REICHEL, W. (1997) Radial symmetry for elliptic boundary-value problems on exterior domains. *Archive for Rational Mechanics and Analysis*, **137**, 381–394.
- RODIN, G. J. & STEINBACH, O. (2003) Boundary element preconditioners for problems defined on slender domains. *SIAM Journal on Scientific Computing*, **24**, 1450–1464 (electronic).
- SERRIN, J. (1971) A symmetry problem in potential theory. *Archive for Rational Mechanics and Analysis*, **43**, 304–318.
- SLOAN, I. H. (1992) Error analysis of boundary integral methods. *Acta Numer.* Cambridge: Cambridge Univ. Press, pp. 287–339.
- SPENCE, E. A., CHANDLER-WILDE, S. N., GRAHAM, I. G. & SMYSHLYAEV, V. P. (2011) A new frequency-uniform coercive boundary integral equation for acoustic scattering. *Communications on Pure and Applied Mathematics*, **64**, 1384–1415.
- STAKGOLD, I. (1967) *Boundary value problems of mathematical physics, Volume I*. New York: The Macmillan Company; London: Collier-Macmillan Ltd.
- TREFETHEN, L. N. & EMBREE, M. (2005) *Spectra and pseudospectra*. Princeton, NJ: Princeton University Press.

WARNICK, K. F. & CHEW, W. C. (2001) On the spectrum of the electric field integral equation and the convergence of the moment method. *Internat. J. Numer. Methods Engrg.*, **51**, 31–56.

WRIGHT, T. G. (2002). EigTool, <http://www.comlab.ox.ac.uk/pseudospectra/eigtool>.