

**Università degli Studi di Milano**

---

FACOLTÀ DI SCIENZE MATEMATICHE, FISICHE E NATURALI  
DIPARTIMENTO DI MATEMATICA  
CORSO DI DOTTORATO DI RICERCA IN MATEMATICA  
XVII CICLO

**The unirationality of the moduli  
space of étale double covers of  
genus five curves**

MAT/03

Tesi di Dottorato di:  
**Marco Lo Giudice**

Relatore:  
**Dr. G. K. Sankaran**  
Department of Mathematical Sciences  
University of Bath

Coordinatore del Dottorato:  
**Prof. Antonio Lanteri**  
Dipartimento di Matematica  
Università degli Studi di Milano

Anno Accademico 2004/2005

To my family, with love.

# Acknowledgements

My international experience was supported from England by the EPSRC (Engineering and Physical Sciences Research Council) and from Italy by the *Università degli studi di Milano*; I am very grateful to both organisations and to the University of Bath, who welcomed me in the last three years.

I wish to thank my supervisor, Dr G. K. Sankaran who guided me on my way towards Algebraic Geometry; Prof Antonio Lanteri in Milan, not only for a brilliant course on algebraic surfaces but also for his patience and advice on “how to get a PhD;” and Prof Elham Izadi, who provided the original plan that guided my research throughout.

Being the final outcome of a long journey, from Genoa to Perugia to Milan and then to Warwick and Bath, this thesis was made possible by a variety of people that I was really lucky to meet in the right place and at the right time. I will never forget the support and the encouragement received from Prof Mauro Beltrametti, or the good time spent in Prof Enrique Arrondo’s company in Perugia, where I also met Prof Ivan Graham. Prof Bert van Geemen from Milan was always aware of my progress, and in Bath Dr Alastair King provided accurate and useful feedback. I wish to express all my gratitude to Prof Miles Reid for the time spent “talking maths” with me, Prof Fran Burstall for being such a good teacher and lecturer, and Prof Edoardo Sernesi who kindly replied to my electronic request of help.

As strange as it may seem, I still have friends who don’t know what a variety is and for whom a *ring* is something to be worn; this is mostly due to the patience and tolerance of just one woman, Huang Min-Fang, met one night on a musical trip to Bristol and very soon become the focus of my life (I love you Olive). Many thanks therefore to Charbel Moussa, Edwige Sima-Ella, Emanuele Guglielmino, Mark Rizzo, Clarice Chan, Carla Ramos, Ma-

men Valor, Mark Price, Pamela Fan, Andy and Christina Mitchell, Ioannis and J.J. Ignatiadis, Ting Ye and Yun Yao, Hiroko Tamaki, Kumiko Tsutsui and Motoko Nakamura. Without forgetting those I left in Italy, Davide Odetti, Fabio Carbone, Tiziano Caverni, Fabio Negri, Daniele Assereto, Daniele Purrone, Elisa Sartor, Carla Novelli, Alessio Del Padrone, Antonella Bertero. Religion has acquired a particularly important meaning for me in the last few years and I wish to thank Father Christopher (Coventry), Father Bill McLoughlin OSM and Father Tom for their spiritual guidance.

I want to dedicate a special mention to my fellow PhD students, especially those whom I shared an office with, in Italy or in England. Alessandra Bernardi, Daniele Cassani, Elisabetta Preve, Andrea Tironi, Silvia Zamperlin, Neil Donaldson, Aurea Quintino, Eugene Varvaruca, Barrie Cooper, Zhi Geng, Vedad Pasic. My fellows in algebraic geometry Nathan Broomhead and Alexander Kasprzyk, with whom I shared a supervisor, and Elizabeth Baldwin. But also Gundula Behrens, James Coughlan, Matt Dorey, Melina Freitag, Peter Hancock, John Harris, Carl-Friedrich Kreiner, Simone Mandica, Susana Santos, James Stewart-Cox.

Many thanks also to Mr Filippo Malaspina in Milan, who managed to make Italian bureaucracy easy, to Mrs Mary Baines, Ms Jill Parker and Miss Sue Paddock, for making me feel at home at Bath, and to Mr Mark Willis, Mr Darrel Johnson and Mr Adam John Prowse for computer related issues, especially when the crucial computation didn't seem to run on my machine.

Infine un affettuoso saluto al professor Franco Scoglio, recentemente salito al terzo anello del "Luigi Ferraris." Fu il tecnico del primo Genoa che io ricordi e contribuì a costruire l'undici che sconfisse il Liverpool ad Anfield.

E qui chiudo.

# Contents

<b>Introduction</b>	<b>1</b>
<b>1 Basic Material</b>	<b>7</b>
1.1 Schemes . . . . .	7
1.2 Abelian Varieties . . . . .	13
1.3 Prym Varieties . . . . .	17
<b>2 Quartic Surfaces and Double Covers of Curves</b>	<b>19</b>
2.1 Quartic Surfaces . . . . .	19
2.2 Double Solids and Conic Bundles . . . . .	23
2.3 How genus five curves arise . . . . .	28
2.4 An Explicit Example . . . . .	31
<b>3 Building Families</b>	<b>34</b>
3.1 Moduli Problems and Functors . . . . .	34
3.2 The Moduli Space of Quartic Surfaces . . . . .	38
3.3 Families of Curves and their Dimension . . . . .	44
<b>4 Computations</b>	<b>51</b>
4.1 The Standard Test Example . . . . .	51
4.2 The dimension of the family of surfaces . . . . .	56
4.3 The Family of Plane Curves . . . . .	59
4.4 The Rank of a big Matrix . . . . .	62
<b>Bibliography</b>	<b>67</b>

# Introduction

有志者事竟成  
Nothing is impossible  
to a willing mind.

## Moduli Spaces and Unirationality

Let  $X$  be an algebraic variety over a field  $\mathbb{K}$ . We say that  $X$  is *unirational* if there exists a dominant rational map  $\mathbb{P}^n \dashrightarrow X$  from a projective space to  $X$ . As Kollár, Smith and Corti [29] point out, here the use of the prefix “uni” refers to the map from  $\mathbb{P}^n$  to the variety, defined in one direction only in contrast with the two mutually inverse rational maps required for rationality.

Already in 1915 Severi proved that the moduli space  $\mathcal{M}_g$  of curves of genus  $g$  is unirational for  $g \leq 10$ —see Arbarello and Sernesi, [2], for a modern rigorous proof—and conjectured the same to hold for every genus; but this was to remain open until the beginning of the 1980s, when Harris and Mumford [27] disproved it by showing that over the complex numbers,  $\mathcal{M}_g$  is of general type if  $g$  is odd and  $g \geq 25$ . Meanwhile Sernesi [48] proved the unirationality of  $\mathcal{M}_{12}$ , thus leaving a gap at genus 11 that was filled later by Chang and Ran [6], who also proved that  $\mathcal{M}_{13}$  is unirational. The question of what happens for large  $g$  was settled when Eisenbud and Harris [17] published their proof that  $\mathcal{M}_g$  is of general type for all  $g \geq 24$  and  $\mathcal{M}_{23}$  has Kodaira dimension at least 1.

A rather similar story unfolded for  $\mathcal{A}_g$ , the moduli space of principally polarised Abelian varieties of dimension  $g$ . Tai [52] proved that  $\mathcal{A}_g$  is of general type for  $g \geq 9$ : this was improved to  $g \geq 8$  by Freitag [19] and to  $g \geq 7$  by Mumford [41]. This was the state of knowledge when Donagi [12] proved that

$\mathcal{A}_5$  is unirational. His proof is related to work by Clemens [8], who proved that  $\mathcal{A}_4$  is unirational, using intermediate Jacobians. Very recently Grushevsky and Lehavi [25] have shown that  $\mathcal{A}_6$  is of general type.

However, Donagi proved something else: he also showed the unirationality of  $\mathcal{R}_6$ , the moduli space of étale double covers of curves of genus six. Other proofs of this were given by Verra [54] and by Mori and Mukai [38]. It follows from this that  $\mathcal{A}_5$  is unirational, because of the generic surjectivity (also shown by Donagi) of the *Prym map*.

Let  $\pi: \tilde{C} \rightarrow C$  be an étale double cover of a smooth complete algebraic curve  $C$  of genus  $g \geq 2$ , and let  $\text{Jac}(\tilde{C})$  and  $\text{Jac}(C)$  be the Jacobians. As Mumford [40] explains, the Jacobian  $\text{Jac}(\tilde{C})$  splits under the action of the involution  $\iota: \tilde{C} \rightarrow \tilde{C}$  interchanging the two sheets of the cover, into a positive part and a negative part  $P$ , the Prym variety. The principal polarisation of  $\text{Jac}(\tilde{C})$  induces twice a principal polarisation on  $P$  and it is a fact that the dimension of  $P$  is  $g - 1$ . By associating to a double cover  $\pi$  its Prym variety we define the *Prym map*, that is a morphism

$$p_g: \mathcal{R}_g \longrightarrow \mathcal{A}_{g-1}$$

where  $\mathcal{R}_g$  denotes the moduli space of étale double covers of curves of genus  $g$ , or equivalently the moduli space of curves  $C \in \mathcal{M}_g$  equipped with a (nonzero) line bundle  $\mathcal{L}$  whose square is trivial. Thus  $\mathcal{R}_g$  is also equipped with a morphism to  $\mathcal{M}_g$ , which is a finite cover of degree  $2^{2g} - 1$ .

It is known [see Beauville, 4] that for  $g \geq 7$  the Prym map is generically injective, while for  $g \leq 6$  it is generically surjective. Thus in proving that  $\mathcal{R}_6$  is unirational, Donagi [12] was able to show the unirationality of  $\mathcal{A}_5$  using the Prym map. At about the same time Catanese [5] proved that  $\mathcal{R}_4$  is rational. Since then there has been a gap between four and six, which this thesis proposes to fill; in fact the same result will provide another proof of the unirationality of  $\mathcal{A}_4$ , substantially different from the one in Clemens [8].

The Prym map has been described extensively by Donagi and Smith [14] both in genus 6 and in genus 5. Classically  $p_6$  was known to be generically finite, and quite famously they proved that its degree is 27. Later Donagi [13] also showed that the general fibre of  $p_6$  carries the structure of the configuration of lines on a cubic surface, and gave a complete description of the two-dimensional fibres of  $p_5$ .

## The Genus Five Case

Let  $X$  be a hypersurface in  $\mathbb{P}^3$  of degree four, whose only singularities are six isolated ordinary double points  $P_0, \dots, P_5$ . Working in coordinates and realising  $\mathbb{P}^3$  as the projective spectrum of the polynomial ring  $R = \mathbb{K}[x_0, \dots, x_3]$ , assume the point  $P_0$  to be  $[0 : 0 : 0 : 1]$  and  $X$  to be given by  $\text{Proj } R/(F)$  where  $F$  is a form of degree four contained in  $(x_0, x_1, x_2)^2$ . In particular  $F$  can be written as follows

$$F = u_2x_3^2 + 2u_3x_3 + u_4$$

where  $u_d$  is a form of degree  $d$  in  $\mathbb{K}[x_0, x_1, x_2]$ . To such a surface we can associate a plane curve  $C_X$  as follows. Realise  $\mathbb{P}^2$  as the set of all the lines through  $P_0$ , more precisely fix a hyperplane in  $\mathbb{P}^3$  not containing  $P_0$  and projecting define a morphism  $\mathbb{P}^3 \setminus \{P_0\} \rightarrow \mathbb{P}^2$  whose fibres (over the closed points) are all the lines through the point  $P_0$ . Composing with the closed immersion of  $X \setminus \{P_0\}$  in the punctured projective space defines a morphism  $\pi: X \setminus \{P_0\} \rightarrow \mathbb{P}^2$  with finite fibres (in fact, generically finite). The curve  $C_X$  will be the scheme of lines through  $P_0$  that are tangent to  $X$  away from  $P_0$ , which is a closed subscheme of the locus of points in  $\mathbb{P}^2$  whose fibre under  $\pi$  consists of one point only. More precisely, working in coordinates as above, it is given by the following projective spectrum

$$C_X = \text{Proj } \mathbb{K}[x_0, x_1, x_2]/(u_3^2 - u_2u_4)$$

Furthermore the generic surface  $X$  gives rise to a curve  $C_X$  whose only singularities come from singularities of  $X$ , in other words generically the curve  $C_X$  has precisely five double points  $P_1, \dots, P_5$ , and therefore its arithmetic genus is five.

Following Clemens [8] we can now associate to the surface  $X$  a (quartic) *double solid*, that is a threefold  $\Lambda_X$  endowed with a two-sheeted covering  $p: \Lambda_X \rightarrow \mathbb{P}^3$  branched along the surface  $X$ . This can be realised as the closed subscheme of the *weighted projective space*  $\mathbb{P}(1, 1, 1, 1, 2)$  given by the equation  $F - t^2$ , in other words

$$\Lambda_X = \text{Proj } \mathbb{K}[x_0, \dots, x_3, t]/(F - t^2)$$

where the degree of  $x_i$  is considered to be 1, and the degree of  $t$  to be 2. According to Kreussler [31]  $\Lambda_X$  is birational to a conic bundle over  $\mathbb{P}^2$ : more precisely the unique morphism of schemes  $f$  which makes the following diagram commute

$$\begin{array}{ccc} \mathrm{Bl}_{P_0}(\Lambda_X) & \longrightarrow & \Lambda_X \xrightarrow{p} \mathbb{P}^3 \\ & \searrow f & \downarrow \pi \\ & & \mathbb{P}^2 \end{array}$$

is a conic bundle over  $\mathbb{P}^2$  for which the curve  $C_X$  is the locus of points whose fibre is degenerate. Blowing up in the remaining five points we construct a conic bundle over a degree 4 del Pezzo surface containing the canonical model of the curve  $C_X$ , which is given by the locus of points whose fibre consists of two lines meeting transversally, and comes therefore naturally endowed with a double cover.

Let  $\mathcal{Q}$  be the space of quartic hypersurfaces in  $\mathbb{P}^3$  with six isolated ordinary double points, one of which is marked. To see that this is unirational fix five points  $P_1, \dots, P_5$  in  $\mathbb{P}^3$  and consider the space  $B_0$  of all quartic hypersurfaces with double points at  $P_1, \dots, P_5$  and another at  $P_0$ , where  $P_0$  is not fixed. Each fibre of the morphism from  $B_0$  to  $\mathbb{P}^3$  associating to each surface the sixth node  $P_0$  is just a linear space, therefore  $B_0$  is a projective vector bundle over an open subset of  $\mathbb{P}^3$  and hence is rational. This proves that the construction above defines a morphism from a unirational variety to  $\mathcal{R}_5$ , which is in turn endowed with a finite (in fact 1023-to-1) natural projection to  $\mathcal{M}_5$ , the moduli space of curves of genus five. Since  $\mathcal{R}_5$  is known to be irreducible, to prove the unirationality it is now enough to prove that the map to  $\mathcal{M}_5$  is generically surjective

We now have a family of canonically embedded curves of genus five, each one given by the intersection of three quadrics in  $\mathbb{P}^4$ . Following the notes on deformations of schemes by Sernesi [49], we fix a central fibre in this family and with *Macaulay 2* [21] we compute explicit equations for it, whose coefficients depend on 13 parameters. The computation of the Kodaira-Spencer map is now reduced to linear algebra, in particular we exploit the property of our curves of being canonically embedded, meaning that two of them will be isomorphic if and only if they are projectively equivalent. In other words we

are now looking at the Grassmannian  $\text{Gr}(3, 15)$  of 3-dimensional linear spaces of quadrics in  $\mathbb{P}^4$ , more precisely at its tangent space at the point described by our central fibre. Taking into account the action of  $\text{SL}(5)$  describing classes of isomorphism, our computation reduces to computing the rank of a big matrix and our result is that this rank is maximal.

It is likely that the need for direct computation could be reduced, but we took this approach for practical reasons. Note that Donagi [12] also relies upon computer calculations.

## The Structure of this Thesis

This thesis is divided into four chapters. Notation is fixed in the first one by reviewing some standard basic topics in algebraic geometry, in particular schemes. After a brief description of Abelian varieties, with some emphasis on Jacobians and Pryms, the chapter ends with an original argument for the unirationality of  $\mathcal{A}_4$ , the moduli space of principally polarised Abelian varieties of dimension four.

The second chapter is devoted to the description of the geometric configuration concerning quartic hypersurfaces  $X$  in  $\mathbb{P}^3$  with at least one ordinary double point  $P$ . After the definition of the associated plane sextic  $C_X$  we analyse its singularities, showing that a surface  $X$  which is generic in a very precise sense gives rise to a curve  $C_X$  whose singularities are in one-to-one correspondence with the singularities of  $X$  other than  $P$ . Quartic double solids are introduced, and a comparative discussion about equivalent possible definitions is provided, before we state Kreussler's theorem [31] about double solids and conic bundles. But the main result is the following.

**Proposition (2.8).** *Let  $X$  be a generic quartic hypersurface in  $\mathbb{P}^3$  with six isolated ordinary double points  $P_0, \dots, P_5$ , and let  $\widetilde{C}_X$  be the canonical model of the associated plane sextic with respect to  $P_0$ . Then there exists a natural unbranched double cover of the curve  $\widetilde{C}_X$ .*

The possibility that the double cover might be trivial, that is consisting of two copies of the curve  $\widetilde{C}_X$ , is not ruled out directly, but it is shown to be equivalent to the reducibility of the locus of degeneracy of a conic bundle  $W_X$ . This

leads to a description of an open condition over the moduli space of quartic surfaces in  $\mathbb{P}^3$ , which a direct computation shows to be nonempty.

Finally we describe a general method to compute an explicit example, providing some background to better explain the results obtained with *Macaulay 2* later in the thesis. In particular we write down a homogeneous polynomial of degree four providing the equation for our standard test surface  $\mathbb{X}$ .

Chapter 3 is all about moduli spaces. We review the definition of fine and coarse moduli space, briefly discussing the Hilbert scheme as an example of a fine moduli space, and describe how to define a morphism using natural transformations. We then show that the variety  $\mathcal{Q}$ , parametrising quartic hypersurfaces in  $\mathbb{P}^3$  with six isolated ordinary double points, one of which is marked, up to projective equivalence, is unirational and 13-dimensional; and define a morphism  $\mathcal{Q} \rightarrow \mathcal{R}_5$ . With the main result in mind we then look for sufficient conditions for this morphism to be generically surjective.

**Theorem (3.14).** *The moduli space  $\mathcal{R}_5$  of étale two-sheeted coverings of curves of genus five is unirational.*

The computation of the differential of the morphism  $\mathcal{Q} \rightarrow \mathcal{M}_5$ , obtained from the one above by composition with  $\mathcal{R}_5 \rightarrow \mathcal{M}_5$ , at one given point  $\mathbb{X}$  is reduced to the computation of the rank of a  $46 \times 45$  matrix, and the theorem follows if it is shown that this rank is maximal for a generic quartic surface. Again we claim that the test curve defined in Chapter 2 satisfies this condition.

To complete the proof of the theorem, and to show that all the open sets we have considered so far are non-empty, we provide in Chapter 4 the results obtained with *Macaulay 2* when performing the computations described earlier in the thesis.

# Chapter 1

## Basic Material

After a very brief introduction to schemes we describe Abelian varieties, with some emphasis on Jacobians and Pryms. The former is done to fix notation while the latter to provide some context for our main result. In fact the chapter will end with an original argument for the unirationality of  $\mathcal{A}_4$ , the moduli space of principally polarised Abelian varieties of dimension four.

### 1.1 Schemes

The concept of a *scheme* naturally arises from the observation that an affine variety and its coordinate ring are essentially the same object. In fact, more than being just a remark, this awareness is an intrinsic part of the theory from the very beginning, whether expressed in classical style, as for instance on Shafarevich [50] or in a more modern way as on the two undergraduate texts by Miles Reid [43, 44]. In great generality the *prime spectrum* of a ring  $A$  is defined to be the set  $\text{Spec } A$  of all its prime ideals: thus for example consider the spectrum of the ring of polynomials over a field  $\mathbb{K}$ , that is

$$\mathbb{A}_{\mathbb{K}}^n := \text{Spec } \mathbb{K}[X_1, \dots, X_n].$$

Famously Hilbert's Nullstellensatz states that  $\mathbb{K}$  is algebraically closed if and only if the set of maximal ideals in this spectrum is in bijection with  $n$ -tuples in  $\mathbb{K}^n$ , so it makes sense to call this abstract set *affine space*. The Zariski topology is introduced in the classic way, only in this context it is more natural; indeed

closed subsets—denoted  $\mathcal{V}(\mathfrak{a})$ —will consist of prime ideals lying *above* a given ideal  $\mathfrak{a} \subseteq A$ , and a base for the topology will be given by open sets cut out by principal ideals, in other words by sets  $D(\alpha)$  defined to be the complement of  $\mathcal{V}(\alpha)$ , for some element  $\alpha \in A$ . Finally the main ingredient is provided by sheaf theory, as it always is according to Tennison [53], who points out that several types of manifolds can be obtained by specialising the abstract concept of *geometric space*. In the case of schemes this is achieved by defining  $\mathcal{O}(D(\alpha))$  to be the localised ring  $A_\alpha$  for any distinguished open subset  $D(\alpha)$ . A *scheme* is then a geometric space locally isomorphic to the spectrum of a ring, in particular it is a pair  $(X, \mathcal{O}_X)$  consisting of a topological space  $X$  and a sheaf of rings over  $X$ , satisfying the following properties:

- there exists an open *affine* covering of  $X$ , consisting of open subsets  $U$  such that the pair  $(U, \mathcal{O}_X|_U)$  is isomorphic to  $\text{Spec } \mathcal{O}_X(U)$ ;
- for any point  $x \in X$ , the stalk  $\mathcal{O}_{X,x}$  is a local ring;
- for any open subset  $U \subseteq X$ , the pair  $(U, \mathcal{O}_X|_U)$  is again a scheme.

The natural identification between an affine variety and its coordinate ring becomes in this way a precise statement: the category of affine schemes is equivalent to the category of commutative rings with identity, with arrows reversed. This equivalence is given by the contravariant functor  $\text{Spec}$ : more precisely Eisenbud and Harris [18, Theorem I-40] prove it is a consequence of the adjunction property of  $\text{Spec}$ .

There are essentially two possible ways to define projective space  $\mathbb{P}_{\mathbb{K}}^n$ : one consists in gluing together  $n + 1$  copies of affine  $n$ -space and the other is to introduce an equivalence relation on  $\mathbb{A}_{\mathbb{K}}^{n+1}$ . Both these ideas are encoded in the theory of schemes: the former is a fundamental procedure sometimes called *gluing lemma* and described in detail by Shafarevich [51, §V.3.2], and the latter amounts to a general construction analogous to  $\text{Spec}$ , the *homogeneous spectrum of a graded ring*. Given a graded ring  $S = \bigoplus_{d \geq 0} S_d$ , Liu [33, §2.3.3] defines  $\text{Proj } S$  to be the set of all homogeneous prime ideals  $\mathfrak{p}$  which do not contain all of  $S_+ = \bigoplus_{d \geq 1} S_d$ , and introduces in this set the topology induced by the inclusion  $\text{Proj } S \subset \text{Spec } S$ . Thus a closed subset of  $\text{Proj } S$  will be given by the intersection  $\mathcal{V}_h(E) = \mathcal{V}(E) \cap \text{Proj } S$ , where  $E$  could be in principle any subset of

$S$  but in fact can be assumed to contain only homogeneous elements of positive degree. Again open sets  $D_h(\alpha)$  corresponding to sets  $E$  with just one element form a base for the Zariski topology. Observe also that  $D_h(\alpha)$  coincides with the intersection  $D(\alpha) \cap \text{Proj } S$ . Then, if  $S_{(\alpha)}$  denotes the subring of elements of degree zero in the localised ring  $S_\alpha$ ,  $\text{Proj } S$  can be endowed with a unique structure of a scheme such that for any homogeneous  $\alpha \in S_+$ , the open set  $D_h(\alpha)$  is affine and isomorphic to  $\text{Spec } S_{(\alpha)}$  [Liu, 33, Proposition 2.3.38]. Projective space is thus defined to be the homogeneous spectrum of the polynomial ring, that is

$$\mathbb{P}_{\mathbb{K}}^n := \text{Proj } \mathbb{K}[X_0, \dots, X_n]$$

and each of the  $n + 1$  copies of affine space that cover it is given by  $D_h(X_i)$ . In fact it is possible to state a general characterisation for a union of basic open sets to cover  $\text{Proj } S$ .

**Proposition 1.1.** *Let  $E$  be any subset of  $S$  consisting of homogeneous elements of positive degree, and let  $\mathfrak{a}$  be the homogeneous ideal generated by  $E$ . Then the following are equivalent:*

$$(a) \quad \bigcup_{\alpha \in E} D_h(\alpha) = \text{Proj } S; \quad (b) \quad \mathcal{V}_h(\mathfrak{a}) = \emptyset; \quad (c) \quad S_+ \subseteq \sqrt{\mathfrak{a}}.$$

*Proof.* Recall that the radical  $\sqrt{\mathfrak{a}}$  of a homogeneous ideal  $\mathfrak{a}$  is homogeneous, and that the complement of the closed subset  $\mathcal{V}_h(\mathfrak{a})$  is given by the union of  $D_h(\alpha)$  where  $\alpha$  runs through a set of generators of  $\mathfrak{a}$ .  $\square$

Projective space can in fact be constructed *over* any ring  $A$  by simply considering the projective spectrum of  $A[X_0, \dots, X_n]$ , and more generally over any scheme by means of a *fibred product*. In this case it is written  $\mathbb{P}_B^n$  where  $B$  is either the *base scheme* or the base ring. By fibred product we mean here the familiar notion from category theory—see Mac Lane [34, §III.4]—sometimes called also pull-back and described by a well known universal property. As Mumford writes [42, §II.2], “there is one exceedingly important and very elementary existence theorem in the category of (pre)schemes. This asserts that arbitrary fibred products exist.” When all the schemes involved are affine, say  $B = \text{Spec } A$  is the base and  $X = \text{Spec } R$ ,  $Y = \text{Spec } T$  are spectra of  $A$ -algebras,

the fibred product is also affine and is given by

$$X \times_S Y = \text{Spec}(R \otimes_A T)$$

In this context the word *projective* assumes a relative meaning: a scheme  $X$  is projective over  $B$  if there exists a *closed immersion*  $X \hookrightarrow \mathbb{P}_B^n$ . In the equivalence of categories between affine schemes and commutative rings, closed immersions correspond to surjective homomorphisms, and closed subschemes correspond to ideals; but since  $\text{Proj}$  is not even a functor some work is required to establish the next result, which we quote from Liu [33, Proposition V.1.30].

**Proposition 1.2.** *Let  $A$  be a ring. A scheme  $X$  over  $A$  is projective if and only if it is isomorphic to  $\text{Proj } S/I$ , where  $S = A[X_0, \dots, X_n]$  and  $I$  is a homogeneous ideal of  $S$  contained in  $S_+$ .*


In this context a *projective algebraic variety* will be a projective scheme over a field  $\mathbb{K}$ , usually assumed to be algebraically closed, defined by a homogeneous prime ideal  $\mathfrak{p}$  inside the polynomial ring  $\mathbb{K}[X_0, \dots, X_n]$ .

If  $f: X \rightarrow Y$  is any continuous map, we can factor it through the closed subset  $\overline{f(X)}$  of  $Y$  as a dominant morphism followed by a closed immersion. The same factorisation is possible in the set-up of schemes, although not in complete generality. For instance if  $X$  and  $Y$  are affine schemes,  $f$  is induced by a ring homomorphism  $\varphi: B \rightarrow A$  which will factor through the sub-ring of  $A$  given by the image of  $\varphi$  as follows

$$B \longrightarrow \text{Im } \varphi \longrightarrow A$$


where  $B \rightarrow \text{Im } \varphi$  is surjective, and  $\text{Im } \varphi \rightarrow A$  is injective. This purely algebraic configuration corresponds precisely to the geometric one, indeed injective homomorphisms induce dominant maps and surjective homomorphisms induce closed immersions—see Hartshorne [28, Exercise II.2.18]. We call  $\text{Spec Im } \varphi$  the *scheme-theoretic image of  $f$* , and we denote it  $\overline{f(X)}$  in analogy with the topological construction. The following result can be found in Hartshorne [28, Exercise II.3.11] or Eisenbud and Harris [18, Definition V-2] or also Liu [33, Exercise II.3.17].

**Theorem 1.3.** *Let  $f: X \rightarrow Y$  be a quasi-compact morphism of schemes. Then there exists a scheme  $\overline{f(X)}$ , endowed with a closed immersion  $j: \overline{f(X)} \rightarrow Y$  and a dominant morphism  $g: X \rightarrow \overline{f(X)}$  such that  $f = jg$  and with the following universal property*



for every other scheme  $W$  endowed with a closed immersion  $i$  and a morphism  $h$  such that  $f = ih$  there exists a unique closed immersion  $\theta$  such that  $i\theta = j$ .

Furthermore, for every open subset  $U \subseteq Y$  we have the following commutative diagram, where  $Z' = \overline{f(f^{-1}(U))}$  and vertical arrows are open immersions



That is, the scheme-theoretic image of the restricted morphism  $f|_U$  is naturally endowed with an open immersion in the scheme-theoretic image of  $f$ .

*Proof.* If  $f$  is a morphism of affine schemes the above argument completes the proof, the universal property being easily checked (note that  $\overline{f(X)}$  is necessarily affine). If  $Y = \text{Spec } A$  is affine, then  $f$  is adjoint to the homomorphism  $f_Y^\#: A \rightarrow \Gamma(X, \mathcal{O}_X)$  and if such a scheme exists it must be affine. Since  $f$  is quasi-compact,  $X$  can be covered by a finite number of open affine subsets, say  $V_i = \text{Spec } B_i$ ; the composition  $V_i \rightarrow X \rightarrow A$  is induced by the composition of ring homomorphisms  $r_i \circ f_Y^\#$  where  $r_i: \Gamma(X, \mathcal{O}_X) \rightarrow B_i$  is the restriction homomorphism of the sheaf  $\mathcal{O}_X$ , let  $\mathfrak{a}_i$  be its kernel and let  $\mathfrak{a}$  be the sum of the  $\mathfrak{a}_i$ . Now the affine scheme  $\text{Spec } A/\mathfrak{a}$  comes endowed with a closed immersion and a dominant morphism as required, and the universal property is easily

checked (again  $\overline{f(X)}$  is necessarily affine).

Let  $U = D(\alpha)$  be an open basic subset of  $Y$ , which we still assume to be affine, then as in Hartshorne [28, Exercise II.2.16]  $f^{-1}(U) = X_\alpha$  and for each  $i$   $X_\alpha \cap V_i$  is affine and given by  $\text{Spec } B_{i\alpha}$ . This proves that, when  $Y$  is affine and  $U$  is an open basic subset of  $Y$ , there exists an open immersion  $Z' \rightarrow \overline{f(X)}$  as in the statement. If now we let  $U$  to be any open affine subset of  $Y$ , we can cover  $U$  by basic open sets—which will be basic open sets for both  $Y$  and  $U$ —and construct a collection of open immersions for both  $\overline{f(X)}$  and  $Z'$ , indexed on the open covering of  $U$  and therefore covering  $Z'$ . Thus we construct the open immersion  $Z' \rightarrow \overline{f(X)}$  by a gluing argument.

To complete the proof we need another gluing argument. When  $X$  and  $Y$  are not affine, we take an open affine covering  $\{U_i\}$  of  $Y$  and construct the family of schemes  $\{(Z_i, g_i, c_i)\}$  given by the scheme-theoretic images of the restrictions  $f|^{U_i}$ , where  $g_i$  is a dominant morphism and  $c_i$  is a closed immersion. In order to apply the Gluing Lemma on any of these schemes  $Z_i$  we define for each  $j$  the open subscheme  $Z_{ij} \subseteq Z_i$  as  $c_i^{-1}(U_i \cap U_j)$ . For any open affine subset of  $U_i \cap U_j$  we construct the scheme-theoretic image of the restriction of  $f$  and obtain an open immersion in both  $Z_{ij}$  and  $Z_{ji}$ , defining eventually an isomorphism.  $\square$

Since any morphism of varieties is of finite type, and any morphism of finite type is quasi-compact, this construction makes perfect sense for any morphism between algebraic varieties. Quite surprisingly however we can prove something more, namely the image of a variety is again a variety.

**Corollary 1.4.** *Let  $f: X \rightarrow Y$  be a quasi-compact morphism of schemes. When  $X$  is reduced the scheme-theoretic image of  $f$  is a reduced scheme, while when  $X$  is an integral scheme it is integral.*

*Proof.* Let  $Z$  be the scheme-theoretic image of  $f$  as in the Theorem, and let  $\tilde{Z}$  be the reduced scheme associated to  $Z$  as in Hartshorne [28, Exercise II.2.3] with closed immersion  $r: \tilde{Z} \rightarrow Z$ . Since  $X$  is reduced, there exists a unique morphism  $h: X \rightarrow \tilde{Z}$  such that  $rh = g$ , and if we let  $i = jr$  we obtain  $ih = jrh = jg = f$ . Therefore by the universal property of  $Z$  there exists a unique  $\theta$  such that  $i\theta = j$ , which means  $jr\theta = j$ . Since closed immersions are monic this implies  $r\theta = \text{id}_Z$ . Viceversa we have  $jr\theta r = i\theta r = jr$  therefore  $r\theta r = r$  which

implies  $\theta r = \text{id}_{\tilde{Z}}$ . In conclusion  $Z$  is isomorphic to  $\tilde{Z}$ , hence it is a reduced scheme.

Assume now  $X$  is an integral scheme, then all the restriction homomorphisms of  $\mathcal{O}_X$  are injective. Therefore with reference to the proof of the Theorem when  $Y = \text{Spec } A$  is affine the scheme theoretic image of  $f$  is given by  $\text{Spec } A/\mathfrak{a}$  where  $\mathfrak{a}$  is the kernel of the morphism  $A \rightarrow \Gamma(X, \mathcal{O}_X)$  inducing  $f$ . The ring  $\Gamma(X, \mathcal{O}_X)$  is a domain, therefore the scheme theoretic image is an integral scheme.  $\square$

## 1.2 Abelian Varieties

Roughly speaking an *Abelian variety* is a projective algebraic variety  $V$  which is also a group. However to make this statement precise one needs to be careful because of the abstract nature of algebraic varieties. The starting point is the concept of *group variety*, which is an algebraic variety over an algebraically closed field  $\mathbb{K}$  whose set of  $\mathbb{K}$ -rational points is a group; but even this can be made more general: see Milne [36]. The group axioms will be phrased in terms of properties of certain morphisms via category theory, with the identity element being itself a morphism from  $\text{Spec } \mathbb{K}$  to  $V$ . Abelian varieties are then defined to be complete connected group varieties, and it is proved [ibid., Corollary 1.4 and §5] that they are projective and commutative.

Over the complex numbers an algebraic variety can be regarded as a particular kind of complex manifold (Kähler), and can be studied using the machinery of differential geometry. This leads to some extent to a dichotomy in the theory, perfectly exemplified by the case of Abelian varieties: over  $\mathbb{C}$  these are indeed *complex tori*, that is quotients  $\mathbb{C}^g/\Gamma$  for some full lattice  $\Gamma$  [Mumford, 39]. As a consequence it is possible to state and prove all the general results about Abelian varieties in terms of rather simple complex functions [theta functions, see Debarre 10] without developing the more abstract theory; however not every complex torus is isomorphic to a projective variety, and one has to look for a *polarization*.

**Theorem 1.5** (Riemann conditions). *A complex torus  $\mathbb{C}^g/\Gamma$  is isomorphic to a projective variety if and only if there exists a skew-symmetric form  $E: \Gamma \times \Gamma \rightarrow \mathbb{Z}$  such*

that

$$(a) \quad E_{\mathbb{R}}(iv, iw) = E_{\mathbb{R}}(v, w);$$

$$(b) \quad E_{\mathbb{R}}(v, iv) > 0, \text{ for every } v \neq 0.$$

where  $E_{\mathbb{R}}$  is the skew-symmetric  $\mathbb{R}$ -bilinear form over  $\mathbb{C}^g$  obtained from  $E$  by extension, since  $\Gamma \otimes \mathbb{R} = \mathbb{C}^g$ .

$E$  is usually called a *Riemann form*, to give one is equivalent to give a Kähler differential form on the complex torus  $\mathbb{C}^g/\Gamma$ , which in turn is equivalent to give an ample line bundle (note that this is not true for arbitrary Kähler manifolds.) A polarisation is then the datum of a Riemann form over the lattice  $\Gamma$ , or one of its equivalent.

According to Mumford<sup>1</sup> the study of complex algebraic curves is equivalent to the study of Riemann surfaces, and he doesn't fail to remark the perversity of algebraic geometers calling *curve* what analysts regard as a two dimensional object. In fact, the analytic point of view can sometimes be closer to intuition, as it is the case in what follows. To any compact Riemann surface it is possible to associate an Abelian variety, the *Jacobian variety*, the geometry of which closely reflects the geometry of the surface. "It is the existence of this auxiliary variety that makes the theory of compact manifolds of dimension one so much more beautiful and complete than the theory of complex manifolds of higher dimensions" [Clemens, 7]. The well known construction is based over two major theorems, due to Abel and Jacobi respectively, and deals with linear systems of divisors [see Lang, 32, for the abstract approach.]

Let  $C$  be a compact Riemann surface. Recall that  $C$  is covered by coordinate neighborhoods  $(U, z)$  where  $U$  can be identified with an open subset of  $\mathbb{C}$  and  $z$  is the complex variable; if  $(U_1, z_1)$  is a second open set, then  $z = u(z_1)$  and  $z_1 = v(z)$  with  $u$  and  $v$  holomorphic functions on  $U \cap U_1$ . To give a differential form  $\omega$  on  $C$ , one has to give an expression  $f(z)dz$  on each  $(U, z)$  such that, on  $U \cap U_1$ ,

$$f(u(z_1)) \cdot u'(z_1) \cdot dz_1 = f_1(z_1) \cdot dz_1$$

---

<sup>1</sup>See the monograph "Curves and their Jacobians," now available as appendix of the famous "red book" [42].

A differential form is *holomorphic* if each of the functions  $f(z)$  is holomorphic (rather than meromorphic). Let  $\omega$  be a differential on  $C$  and let  $\gamma$  be a path in  $U \cap U_1$ ; then

$$\int_{\gamma} f(z)dz = \int_{\gamma} f_1(z_1)dz_1$$

Thus, it makes sense to integrate  $\omega$  along any path in  $C$ . The sheaf of differential forms over  $C$  is called the *canonical sheaf* and it is denoted  $\Omega^1$ .

**Theorem 1.6.** *The set of global sections of  $\Omega^1$  forms a  $g$ -dimensional vector space where  $g$  is the genus of  $C$ .*

We denote this vector space by  $H^0(C, \Omega^1)$ . If  $\omega_1, \dots, \omega_g$  is a basis for the space, then every holomorphic differential is a linear combination,  $\omega = \sum a_i \omega_i$ , of the  $\omega_i$ , and  $\int_{\gamma} \omega = \sum a_i \int_{\gamma} \omega_i$ ; therefore it suffices to understand the finite set of integrals  $\left\{ \int_{\gamma} \omega_1, \dots, \int_{\gamma} \omega_g \right\}$ .

Recall (from topology) that  $C$  is a  $g$ -holed torus, and that  $H_1(C, \mathbb{Z})$  has a canonical basis  $\gamma_1, \dots, \gamma_{2g}$ —roughly speaking each  $\gamma_i$  goes once round one hole. The vectors

$$\pi_j = \begin{pmatrix} \int_{\gamma_j} \omega_1 \\ \vdots \\ \int_{\gamma_j} \omega_g \end{pmatrix} \in \mathbb{C}^g, j = 1, \dots, 2g$$

are called the *period vectors*.

**Theorem 1.7.** *The  $2g$  period vectors are linearly independent over  $\mathbb{R}$ .*

Let  $\Gamma$  denote the lattice generated by the period vectors,  $\Gamma = \bigoplus \mathbb{Z}\pi_i$ , then  $\mathbb{C}^g/\Gamma$  is a complex torus. We shall see that in fact it is an abelian variety (i.e., it has a Riemann form). Fix a point  $P_0$  on  $C$ . If  $P$  is a second point, and  $\gamma$  is a path from  $P_0$  to  $P$ , then  $\omega \mapsto \int_{\gamma} \omega$  is a linear map  $H^0(C, \Omega^1) \rightarrow \mathbb{C}$ . Note that if we replace  $\gamma$  with a different path  $\gamma'$  from  $P_0$  to  $P$ , then  $\gamma'$  differs from  $\gamma$  by a loop. If the loop is contractible, then  $\int_{\gamma} \omega = \int_{\gamma'} \omega$ ; otherwise the two integrals differ by a sum of periods.

**Theorem 1.8 (Jacobi Inversion Formula).** *Let  $\ell$  be a linear map  $H^0(C, \Omega^1) \rightarrow \mathbb{C}$ ; then there exist points  $P_1, \dots, P_g$  on  $C$  and paths  $\gamma_i$  from  $P_0$  to  $P_i$  such that*

$$\ell(\omega) = \sum_i \int_{\gamma_i} \omega$$

for all  $\omega \in H^0(C, \Omega^1)$ .

**Theorem 1.9** (Abel). *Let  $P_1, \dots, P_r$  and  $Q_1, \dots, Q_r$  be points on  $C$  (not necessarily distinct). Then there exists a meromorphic function  $f$  on  $C$  with poles exactly at the  $P_i$  and zeros exactly at the  $Q_i$  if and only if, for all paths  $\gamma_i$  from  $P_0$  to  $P_i$  and all paths  $\gamma_i'$  from  $P_0$  to  $Q_i$ , there exists an element  $\gamma \in H_1(C, \mathbb{Z})$  such that*

$$\sum_i \int_{\gamma_i} \omega - \sum_i \int_{\gamma_i'} \omega = \int_{\gamma} \omega$$

for all  $\omega \in H^0(C, \Omega^1)$ .

Let  $\gamma \in H_1(C, \mathbb{Z})$ ; then  $\omega \mapsto \int_{\gamma} \omega$  is a linear function on the vector space  $H^0(C, \Omega^1)$ , i.e. an element of  $H^0(C, \Omega^1)^*$ . Thus we have a map

$$H_1(C, \mathbb{Z}) \longrightarrow H^0(C, \Omega^1)^*$$

which is injective. Set

$$J(C) = H^0(C, \Omega^1)^* / H_1(C, \mathbb{Z})$$

The choice of a basis for  $H^0(C, \Omega^1)$  identifies  $J(C)$  with  $\mathbb{C}^g / \Gamma$ , thus  $J(C)$  is a complex torus.

**Theorem 1.10.** *The intersection product*

$$H_1(C, \mathbb{Z}) \times H_1(C, \mathbb{Z}) \longrightarrow \mathbb{Z}$$

is a Riemann form on  $J(C)$ . Hence  $J(C)$  is an abelian variety.

Fix a point  $P$  on  $C$ . As we noted above,  $\int_P^Q \omega$  doesn't make sense, because it depends on the choice of a path from  $P$  to  $Q$ . But two choices differ by a loop, and so  $\omega \mapsto \int_P^Q \omega$  is well-defined as an element of

$$H^0(C, \Omega^1)^* / H_1(C, \mathbb{Z})$$

Thus we have a canonical map  $\varphi: C \rightarrow J(C)$  sending  $P$  to 0.

Now consider the map

$$\begin{aligned} \text{Pic}^0 C &\longrightarrow J(C) \\ \sum n_i P_i &\longmapsto \sum n_i \int_P^{P_i} \bullet \end{aligned}$$

The Jacobi inversion formula shows that this map is surjective (in fact it proves more than that). Abel's theorem shows that the kernel of the map is precisely the group of principal divisors. Therefore, the theorems of Abel and Jacobi show precisely that the above map defines an isomorphism

$$\text{Pic}^0 C \longrightarrow J(C)$$

### 1.3 Prym Varieties

Let  $\pi: \tilde{C} \rightarrow C$  be an unramified double covering of a smooth complete algebraic curve  $C$  of positive genus  $g$ . Then by Riemann-Hurwitz relation we know that the genus of  $\tilde{C}$  is  $2g - 1$ . If we denote by  $K_C$  the canonical class of  $C$  the argument is the following: since the covering is unbranched we have  $\deg K_{\tilde{C}} = 2 \deg K_C$ , and for any curve  $C$  we have  $\deg K_C = 2g - 2$ . Let  $J(\tilde{C})$  and  $J(C)$  be the Jacobians of the two curves and let  $\iota: \tilde{C} \rightarrow \tilde{C}$  be the involution of  $\tilde{C}$  that exchanges the two sheets of the covering  $\pi$ . As Mumford [40] explains,  $\iota$  acts on the Jacobian  $J(\tilde{C})$  of  $\tilde{C}$ , and  $J(\tilde{C})$  splits under this action into a positive part and a negative part; we call the negative part  $P$ . A more analytical approach is described by Masiewicki [35]. The principal polarisation over  $J(\tilde{C})$  induces twice a principal polarisation on  $P$ , thus in particular  $P$  is an Abelian variety which we call the Prym variety of the double covering  $\pi$ . It's a fact that the positive part is given by  $\pi^*(J(C))$ , therefore is  $g$  dimensional, thus the Prym variety  $P$  has dimension  $g - 1$ .

Let  $\mathcal{R}_5$  denote the moduli space of étale 2-sheeted coverings  $\pi: \tilde{C} \rightarrow C$ , where  $C$  is a smooth complete curve of genus five. Later in the thesis (Theorem 3.14) we are going to prove that this space is *unirational*, meaning that there exists a dominant rational map  $\mathbb{P}^n \dashrightarrow \mathcal{R}_5$  from a projective space to  $\mathcal{R}_5$ . By associating to any such double covering its Prym variety we construct a

morphism

$$p_5: \mathcal{R}_5 \longrightarrow \mathcal{A}_4$$

where  $\mathcal{A}_4$  denotes the moduli space of Abelian varieties of dimension four, in a sense that will be made clear in §3.1. It is known [see Beauville, 4] that  $p_5$  is generically surjective, and Donagi [13] gave a complete description of its two-dimensional fibres. Thus taking into account Theorem 3.14 the following result is proved.

**Theorem 1.11.**  *$\mathcal{A}_4$  is unirational.*

This theorem is not original in that Clemens [8] proved it more than twenty years ago, but the argument is. Indeed Clemens didn't know that  $\mathcal{R}_5$  is unirational and for his proof he relied on intermediate Jacobians.

## Chapter 2

# Quartic Surfaces and Double Covers of Curves

Given a quartic hypersurface in  $\mathbb{P}^3$  with at least one ordinary double point, it is possible to construct a plane curve endowed with a double cover. The process involves building a quartic double solid, which turns out to be a conic bundle over  $\mathbb{P}^2$ ; the plane curve will be then given by the discriminant of this conic bundle. We will ultimately be interested in the case of six double points: then the plane curve will be a five-nodal sextic, hence of genus five. Throughout the chapter we will work over a fixed field  $\mathbb{K}$  of characteristic different from 2 and 3, which we will assume to be perfect.

### 2.1 Quartic Surfaces

Let  $X$  be a hypersurface in  $\mathbb{P}^3$  of degree four, or equivalently let  $X$  be the projective spectrum of the graded algebra  $S = R/(F)$ , where  $F$  is an irreducible homogeneous polynomial of degree four inside  $R = \mathbb{K}[x_0, \dots, x_3]$ . A point  $P \in X$  is *regular* if the ring  $\mathcal{O}_{X,P} = S_{(\mathfrak{p})}$  is a regular local ring, otherwise  $P$  is *singular*, but it can be singular in many different ways: see Reid [45]. Without loss of generality we can assume  $P$  to be the point  $[0 : 0 : 0 : 1]$  and restrict our attention to the affine open subset  $D_h(x_3)$  of  $X$ , where the surface is given by  $\text{Spec } \mathbb{K}[x, y, z]/(f)$  with  $f$  an irreducible polynomial of degree four, and the point corresponds to the maximal ideal  $\mathfrak{m} = (x, y, z)$ . Writing  $f$  as the sum of

its homogeneous components  $f_1 + \cdots + f_4$ , with subscripts indicating degree, properties of the polynomials  $f_d$  correspond to properties of the point  $P$ ; for instance  $P$  is singular if and only if  $f_1 = 0$ , and  $P$  is said to be a *double point* when  $f_1 = 0$  but  $f_2 \neq 0$ , which can be rephrased abstractly by saying that the local equation  $f$  inside the ring  $\mathcal{O}_{X,P}$  defines a non-zero element of  $\mathfrak{m}^2/\mathfrak{m}^3$ . But even among double points only several different situations can arise, and one needs to look at the *tangent cone* to  $X$  at  $P$  for a finer distinction. This amounts to constructing the quotient of  $\mathbb{K}[x, y, z]$  by the *leading form* of  $f$  (see Milne, [37]), which in this case is  $f_2$ , or to construct abstractly the associated graded ring of the local ring  $\mathcal{O}_{X,P}$ . When  $f_2$  is an irreducible polynomial, or in other words when the associated graded ring is integral, we say that  $P$  is an *ordinary double point*. In terms of the graded algebra  $R/(F)$  this situation is described by saying that  $F$  is a form of degree four contained in the ideal  $(x_0, x_1, x_2)^2$ , so that in particular  $F$  can be written as follows

$$F = u_2x_3^2 + 2u_3x_3 + u_4$$

where  $u_d$  is a form of degree  $d$  in  $\mathbb{K}[x_0, x_1, x_2]$ . The point  $P$  will then be an ordinary double point if the quadratic form  $u_2$  is non-degenerate.

Consider now a quartic surface  $X$  with an isolated ordinary double point  $P = [0 : 0 : 0 : 1]$ , and let  $\pi: X \setminus P \rightarrow \mathbb{P}^2$  be the projection with centre  $P$ . We can realise  $\pi$  abstractly as the morphism associated to the linear system over  $\mathbb{P}^3$  of all the lines through  $P$ , which is given by the 3-dimensional linear space  $H^0(\mathbb{P}^3, \mathcal{P}(1))$  where  $\mathcal{P}$  is the ideal sheaf of the point, or more concretely by fixing a hyperplane  $H$  not containing  $P$  and identifying lines through  $P$  with points in  $H$ ; either way we will construct the morphism of schemes induced by the homogeneous homomorphism

$$\mathbb{K}[y_0, y_1, y_2] \longrightarrow \mathbb{K}[x_0, \dots, x_3]/(F)$$

sending  $y_i$  to  $x_i$  for  $i = 0, 1, 2$ . Any line in  $\mathbb{P}^3$  through the point  $P$  intersects  $X$  in at most two other points therefore, according to standard terminology (see for example Liu, [33]),  $\pi$  is a quasi-finite morphism, meaning that each of its fibres is a finite set, but it is not finite. Nevertheless we can easily determine the maximal open subset over which it is finite.

**Lemma 2.1.** *The restriction of  $\pi$  to the open set  $D_h(u_2) \subseteq \mathbb{P}^2$  is a finite morphism of degree two, and this is the maximal open subset over which  $\pi$  is finite.*

*Proof.* First observe that  $X \setminus P$  is covered by the three affine open sets  $D_h(x_i)$  for  $i = 0, 1, 2$ , and that locally  $\pi$  is induced by the inclusion of  $\mathbb{K}[x, y]$  in the ring

$$A = \mathbb{K}[x, y][z] / (G_2z^2 + 2G_3z + G_4)$$

where  $G_d$  is the polynomial obtained from  $u_d$  by multiplication by  $1/x_i^d$ , in particular  $\pi$  is an affine morphism. Now it is enough to observe that for any multiplicatively closed subset  $S$  of  $\mathbb{K}[x, y]$  the algebra  $S^{-1}A$  is integral over  $S^{-1}\mathbb{K}[x, y]$  if and only if  $G_2 \in S$ , and that over the field  $\mathbb{K}(x, y)$  this is a vector space of dimension 2 generated by 1 and  $z$ .  $\square$

The locus of lines through  $P$  that are tangent to  $X$  away from  $P$  is the plane curve that we propose to study in this section. It is so closely related to the quartic surface that we will denote it by  $C_X$ . In fact what follows could be found already in Kreussler [30, Lemma 5.1].

**Proposition 2.2.**  *$C_X$  is a plane curve of degree six. More precisely, working in coordinates as above, it is given by the following projective spectrum*

$$C_X = \text{Proj } \mathbb{K}[x_0, x_1, x_2] / (u_3^2 - u_2u_4)$$

*Furthermore if  $Q \in X$  is a singular point (different from  $P$ ), then the point  $\pi(Q)$  is a singular point of the curve  $C_X$ .*

*Proof.* In order to compute the equation of the curve we need in the first place to compute the fibre of  $\pi$  over any point of  $\mathbb{P}^2$ , and since it is an affine morphism we can restrict our attention to an open affine set. So for  $i = 0, 1, 2$  consider the restriction of  $\pi$  to the open set  $D_h(x_i) \subseteq \mathbb{P}^2$ , which is induced by the inclusion of  $\mathbb{K}[x, y]$  in the algebra  $A$  as above. A point of  $\mathbb{P}^2$  is given over this open set by a maximal ideal  $(x - a, y - b) \subseteq \mathbb{K}[x, y]$  and the fibre over this point is given by the tensor product, over  $\mathbb{K}[x, y]$ , of the quotient  $\mathbb{K}[x, y] / (x - a, y - b)$  with  $A$ . In other words it is given by the spectrum of the ring  $\mathbb{K}[x, y, z] / \mathfrak{a}$  where the ideal  $\mathfrak{a}$  is generated by the polynomials  $x - a$ ,  $y - b$  and  $G_2z^2 + 2G_3z + G_4$ . This ring is naturally isomorphic to the quotient

$\mathbb{K}[z]/(g_2z^2 + 2g_3z + g_4)$  where  $g_d$  is the element of  $\mathbb{K}$  obtained by computing  $G_d(a, b)$ . Now it is clear that the fibre over the point  $(a, b)$  consists of only one point if and only if either  $g_2 = 0$  or  $g_3^2 - g_2g_4 = 0$ . But while in the former case we have one point with multiplicity one—the dimension of the vector space  $\mathbb{K}[z]/(2g_3z + g_4)$  is one—in the latter we have multiplicity two as expected.

If  $Q \in X$  is a double point (different from  $P$ ) we can assume without loss of generality the point  $Q$  to be  $[1 : 0 : 0 : 0]$ , and concentrate our attention on the open set  $D_h(x_0)$ . Then, with the same notation as above, the polynomial  $G_2z^2 + 2G_3z + G_4$  will be an element of  $(x, y, z)^2$ . This implies immediately that  $G_4 \in (x, y)^2$  and that  $G_3 \in (x, y)$ , therefore  $G_3^2 - G_2G_4 \in (x, y)^2$  which means  $\pi(Q)$  is a singular point.  $\square$

It is perhaps useful to remark that the locus of points in  $\mathbb{P}^2$  whose fibre under  $\pi$  consists of only one point is not irreducible. Indeed we have encountered in the proof above the two components given by the curve  $C_X$  and the conic  $u_2 = 0$ . We can say something more about this configuration, again by comparison with properties of the surface.

**Corollary 2.3.** *Let  $\text{Bl}_P(X) \rightarrow X$  be the blow-up of the surface  $X$  at the point  $P$ . Then the unique morphism of schemes  $\text{Bl}_P(X) \rightarrow \mathbb{P}^2$  that commutes with  $\pi$  is a double cover of  $\mathbb{P}^2$  branched along the curve  $C_X$ . Moreover the image of the exceptional divisor in  $\text{Bl}_P(X)$  is the contact conic of  $C_X$  defined by the equation  $u_2 = 0$ .*

*Proof.* It is essentially enough to observe that the tangent cone of  $X$  at  $P$  is defined by the equation  $u_2 = 0$  in  $\mathbb{P}^3$ . Indeed the blow-up of  $X$  in  $P$  is given by the proper transform of  $X$  in the blow-up of  $\mathbb{P}^3$  at  $P$  (Eisenbud and Harris, [18], Proposition IV.21), whose exceptional locus is in correspondence with all the lines through  $P$ . Then the  $(-1)$ -curve inside  $\text{Bl}_P(X)$  will correspond to the set of lines in the tangent cone. To see that the conic  $u_2 = 0$  is a contact conic of  $C_X$  simply look at the ideal  $(u_2, u_3^2 - u_2u_4) = (u_2, u_3^2)$  inside the polynomial ring  $\mathbb{K}[x_0, x_1, x_2]$ .  $\square$

Let  $Y_d = \text{Proj } R/(u_d)$  be the cone over  $P$  defined by the form  $u_d$ , and let  $Q \in X$  be any point different from  $P$  such that  $\pi(Q) \in C_X$ . Then we claim that the following holds.

**Claim.**  $\pi(Q)$  is a singular point for  $C_X$  if and only if either  $Q$  is singular for  $X$  or  $Q \in Y_2 \cap Y_3 \cap Y_4$ .

Indeed if we denote by  $\mathfrak{q}$  the homogeneous prime ideal inside  $R/(F)$  corresponding to the point, then  $Q \in X$  means  $F \in \mathfrak{q}$  and  $\pi(Q) \in C_X$  means  $u_3^2 - u_2u_4 \in \mathfrak{q}$ . Besides from the equality

$$u_2F = (u_2x_3 + u_3)^2 - (u_3^2 - u_2u_4)$$

we obtain immediately that also  $u_2x_3 + u_3$  is in  $\mathfrak{q}$ . Now  $\pi(Q)$  is a singular point if and only if  $u_3^2 - u_2u_4 \in \mathfrak{q}^2$ , and this happens if and only if  $u_2F \in \mathfrak{q}^2$ , which is equivalent to have either  $u_2 \in \mathfrak{q}$  or  $F \in \mathfrak{q}^2$ . In the latter case  $Q$  is a singular point of  $X$ , in the former we also have  $u_3 \in \mathfrak{q}$ , since  $u_2x_3 + u_3 \in \mathfrak{q}$ , and  $u_4 \in \mathfrak{q}$  since  $2u_3x_3 + u_4 \in \mathfrak{q}$ .

**Proposition 2.4.** *The generic quartic surface  $X$  with an isolated double point  $P$  gives rise to a curve  $C_X$  whose only singularities come from singularities of  $X$ . More precisely we require the equation  $F = u_2x_3^2 + 2u_3x_3 + u_4$  to satisfy the condition*

$$\sqrt{(u_2, u_3, u_4)} = (x_0, x_1, x_2)$$

*Proof.* It is enough to rule out generically the case of a point  $Q$  contained in the intersection  $Y_2 \cap Y_3 \cap Y_4$ . Because of each  $Y_d$  is a cone over the point  $P$  we see immediately that if a point is in this intersection then the whole line through  $P$  and  $Q$  is contained in there. Thus we must rule out generically the case of a point in  $\mathbb{P}^2$  contained in the intersection between a non-degenerate conic, a cubic and a quartic curve. Now observe that the intersection between the conic and a generic cubic consists of six points, and the generic quartic in the plane will not go through any of them.  $\square$

## 2.2 Double Solids and Conic Bundles

A *double solid* is a threefold that can be realised as a branched two-sheeted covering of  $\mathbb{P}^3$ . The definition was first introduced by Clemens [8] at a time when few specific examples of threefolds were known, in the attempt to develop

an analogy with curve theory and clearly inspired by hyperelliptic curves. More recently Grooten [23] defines a double solid to be a closed subscheme  $\Lambda$  of the *weighted projective space*  $\mathbb{P}(1, 1, 1, 1, 2)$ , which in turn is defined to be the projective spectrum of the graded ring  $\mathbb{K}[x_0, \dots, x_3, y]$  where the degree of  $x_i$  is considered to be one, and the degree of  $y$  to be two. As Reid [46] explains, “Weighted projective spaces have appeared implicitly in algebraic geometry since ancient times; the most basic example is a hyperelliptic curve  $y^2 = f_{2g+2}(x)$  viewed as a double cover of  $\mathbb{P}^1$ , that is a weighted hypersurface  $C_{2g+2} \subset \mathbb{P}(1, 1, g + 1)$ .” The emphasis is on the double cover of  $\mathbb{P}^3$ , which is induced by the following inclusion of graded rings

$$\mathbb{K}[x_0, \dots, x_3] \hookrightarrow \mathbb{K}[x_0, \dots, x_3, y]/(y^2 - F),$$

and is branched along a hypersurface  $X \subseteq \mathbb{P}^3$  of even degree, represented by the homogeneous polynomial  $F$  inside  $R = \mathbb{K}[x_0, \dots, x_3]$ . For this reason the double solid given by  $\text{Proj}(\mathbb{K}[x_0, \dots, x_3, y]/(y^2 - F))$  will be denoted here  $\Lambda_X$ . Observe that  $\Lambda_X$  is the union of the four open affine subsets  $D_h(x_i)$  (where  $i = 0, \dots, 3$ ), since  $D_h(y) = D_h(y^2) = D_h(F)$  is contained in the union of these, so that the local equation of the threefold is in fact of the form  $t^2 = f(x, y, z)$  near every point.

In the language of Clemens [8] the same construction looks very different and as such it deserves to be mentioned, but to do it we need first to recall some basic facts. It is a well known result that for any scheme  $X$  vector bundles over  $X$  and locally free sheaves on  $X$  are equivalent concepts, the equivalence being given by a natural one-to-one correspondence up to isomorphism. Indeed a *vector bundle* over  $X$  can be defined in this category simply as being a scheme  $E$ , endowed with a morphism  $\pi: E \rightarrow X$ , and isomorphic to  $\mathbf{Spec} \text{Sym } \mathcal{E}$  (as schemes over  $X$ ), for some locally free sheaf  $\mathcal{E}$  of rank  $n$ . Here  $\mathbf{Spec}$  denotes the contravariant functor from quasi-coherent sheaves of  $\mathcal{O}_X$ -algebras to schemes over  $X$  that Eisenbud and Harris [18] call *global spec*. A *section* of a vector bundle over an open set  $U$  is then a morphism  $s: U \rightarrow E$  such that  $f \circ s$  is the open immersion of  $U$  in  $X$ , and the space of all sections is in fact a sheaf over  $X$  usually denoted  $\mathcal{S}_{E/X}$ .

**Theorem 2.5.** *Let  $X$  be a scheme. Then the two functors defined by  $E \mapsto \mathcal{S}_{E/X}$  and*

by  $\mathcal{E} \mapsto \mathbf{Spec} \operatorname{Sym} \mathcal{E}^\vee$  establish an equivalence of categories between vector bundles of rank  $n$  over  $X$  and locally free sheaves of rank  $n$  on  $X$ .

*Proof.* It is enough to prove that the sheaf  $\mathcal{S}_{E/X}$  is naturally identified with the locally free sheaf  $\mathcal{E}^\vee$ . To this purpose note that sections of  $\pi$  over the open subset  $U$  are in one-to-one correspondence with morphisms of quasi-coherent algebras  $\operatorname{Sym} \mathcal{E}|_U \rightarrow \mathcal{O}_X|_U$ , moreover if the open set  $U$  is also affine then by the adjoint property of  $\mathbf{Spec}$  morphisms like these correspond to  $\mathcal{O}_X(U)$ -algebra homomorphisms  $\operatorname{Sym} \mathcal{E}(U) \rightarrow \mathcal{O}_X(U)$ . Now by the very definition of symmetric algebra there is another bijection with homomorphisms of  $\mathcal{O}_X(U)$ -modules between  $\mathcal{E}(U)$  and  $\mathcal{O}_X(U)$ .  $\square$

Let now  $X$  be a hypersurface in  $\mathbb{P}^3$  of even degree  $2d$ . Then  $X$  is defined by a global section  $s$  of the locally-free sheaf  $\mathcal{O}(2d)$  on  $\mathbb{P}^3$ , which in turn can be considered as a morphism  $s: \mathbb{P}^3 \rightarrow E_{2d}$  where  $E_{2d}$  is the line bundle  $\mathbf{Spec} \operatorname{Sym} \mathcal{O}(-2d)$ . Clemens [8] realises a double solid abstractly as the inverse image in  $E_d$  of the scheme theoretic image of  $s$  under the *squaring map*  $E_d \rightarrow E_{2d}$ . Over any open affine subset of the standard cover of  $\mathbb{P}^3$ , that is for any degree one monomial  $e \in \{x_0, \dots, x_3\}$ , this configuration is totally algebraic and corresponds to the following diagram

$$\begin{array}{ccc} S_{(e)}[\tau] & \xrightarrow[\tau \mapsto f]{\sigma} & S_{(e)} \\ \downarrow q \quad \tau \mapsto t^2 & & \\ S_{(e)}[t] & & \end{array}$$

where  $f$  is the local equation of the hypersurface  $X$ ,  $\tau = 1/e^{2d}$  is a generator of the free module of rank one  $\mathcal{O}_X(-2d)|_{D_h(e)}$ , and  $t = 1/e^d$  a generator for  $\mathcal{O}_X(-d)|_{D_h(e)}$ . The double solid is defined by the extension of  $\ker \sigma$  via the homomorphism  $q$ , which is precisely the ideal  $(t^2 - f)$  in accordance with the above discussion.

Observe that singularities in double solids occur only as singularities of the surface. Indeed among the partial derivatives of the local equation of  $\Lambda_X$  there is always the polynomial  $2t$  which defines  $X$  inside the threefold. We are

applying here the Jacobian criterion [Eisenbud, 15, §16.6] for which we need the base field to be a perfect field.

*Quartic double solids* correspond to hypersurfaces in  $\mathbb{P}^3$  of degree four and are described by Clemens in detail first in [8] and later in [9]. Such threefolds are among the main characters of this thesis essentially because the singular ones are related to conic bundles over  $\mathbb{P}^2$  or more generally over del Pezzo surfaces. So assume now  $X$  to be defined by a polynomial of degree four, and to contain at least one ordinary double point  $P_0$ . Denote by  $p$  the double cover of  $\mathbb{P}^3$  given by  $\Lambda_X$ , and recall from Section 2.1 how to associate to  $X$  a plane curve  $C_X$  of degree six, in particular recall that projection from  $P_0$  defines a morphism  $\pi: X \setminus \{P_0\} \rightarrow \mathbb{P}^2$ .

The main result of this section is that the blow-up  $\text{Bl}_{P_0}(\Lambda_X)$  of  $\Lambda_X$  at the point  $P_0$  is in fact a conic bundle.

**Proposition 2.6.** *Let  $X$  be a quartic hypersurface in  $\mathbb{P}^3$  with an ordinary double point  $P_0$ . Then the unique morphism of schemes  $f$  which makes the following diagram commute*

$$\begin{array}{ccc} \text{Bl}_{P_0}(\Lambda_X) & \longrightarrow & \Lambda_X \xrightarrow{p} \mathbb{P}^3 \\ & \searrow f & \downarrow \pi \\ & & \mathbb{P}^2 \end{array}$$

*is a conic bundle over  $\mathbb{P}^2$  for which the curve  $C_X$  is the locus of points whose fibre is a degenerate conic.*

This statement was already contained in Clemens [8, p.222], but Kreuzler [31, Theorem 2.1] improved it by giving an explicit description of all the morphisms involved. The argument starts with the construction of the conic bundle, which is realised as a divisor  $W_X$  inside a  $\mathbb{P}^2$ -bundle over  $\mathbb{P}^2$ . Let  $\mathcal{E}$  be the locally free sheaf of rank three  $\mathcal{O} \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-2)$  over  $\mathbb{P}^2$ , and construct  $\mathbb{P}(\mathcal{E})$  by taking global Proj of the symmetric powers of  $\mathcal{E}$

$$\mathbb{P}(\mathcal{E}) = \mathbf{Proj} \text{Sym } \mathcal{E}$$

as described for instance in Hartshorne [28, §II.7]. In order to define the divisor we are interested in we take constant non-zero global sections  $z_0 \in H^0(\mathbb{P}^2, \mathcal{E})$ ,  $z_1 \in H^0(\mathbb{P}^2, \mathcal{E}(1))$  and  $z_2 \in H^0(\mathbb{P}^2, \mathcal{E}(2))$ , so that for any open set  $U \subseteq \mathbb{P}^2$  any

element  $s \in \mathcal{E}(U)$  can be written in a unique way as a sum

$$s = z_0|_U s_0 + z_1|_U s_1 + z_2|_U s_2$$

with  $s_d \in H^0(U, \mathcal{O}(-d)|_U)$ . In particular over any open set of the standard cover of the projective plane,  $D_h(x_i) \subseteq \text{Proj } S$  with  $S = \mathbb{K}[x_0, x_1, x_2]$ , the restriction of  $\mathcal{E}$  is a coherent sheaf given by

$$\begin{aligned} \mathcal{E}|_{D_h(x_i)} &= \mathcal{O}|_{D_h(x_i)} \oplus \mathcal{O}(-1)|_{D_h(x_i)} \oplus \mathcal{O}(-2)|_{D_h(x_i)} \\ &= \text{Shf } S_{(x_i)} \oplus \text{Shf } \frac{1}{x_i} S_{(x_i)} \oplus \text{Shf } \frac{1}{x_i^2} S_{(x_i)} \\ &= \text{Shf} \left( S_{(x_i)} \oplus \frac{1}{x_i} S_{(x_i)} \oplus \frac{1}{x_i^2} S_{(x_i)} \right) \end{aligned}$$

thus a local base for the free  $S_{(x_i)}$ -module  $\mathcal{E}(D_h(x_i))$  is realised by

$$z_0, \frac{z_1}{x_i}, \frac{z_2}{x_i^2}$$

and taking symmetric powers is equivalent to forming the polynomial ring over  $S_{(x_i)}$  in these variables. We now define the divisor  $W_X$  inside  $\mathbb{P}(\mathcal{E})$  by means of the following polynomial expression inside  $H^0(\mathbb{P}^2, \mathcal{O}(4) \otimes \text{Sym}^2 \mathcal{E})$

$$-z_2^2 + z_1^2 u_2 + 2z_1 z_0 u_3 + z_0^2 u_4$$

By this we mean that on any  $D_h(x_i)$  the conic bundle is given by the projective spectrum of the following graded algebra over  $\mathbb{K}[x, y]$

$$\mathbb{K}[x, y][\zeta_0, \zeta_1, \zeta_2] / (-\zeta_2^2 + \zeta_1^2 G_2 + 2\zeta_1 \zeta_0 G_3 + \zeta_0^2 G_4)$$

where  $\zeta_0, \zeta_1, \zeta_2$  is a local base for  $\mathcal{E}$  and  $G_d$  is the polynomial obtained from  $u_d$  by multiplication by  $1/x_i^d$  as in Lemma 2.1.

To complete the argument Kreussler [30] defines a morphism  $\Phi: W_X \rightarrow \mathbb{P}^3$  and proves that it factors through  $p$ : then he observes that  $W_X$  endowed with this morphism to  $\Lambda_X$  is actually the blow-up of the double solid and shows the compatibility with  $\pi$ .

## 2.3 How genus five curves arise

As we did before we begin this section by considering a quartic hypersurface in  $\mathbb{P}^3$ , but here we will assume the singularities of  $X$  to consist precisely of six isolated ordinary double points,  $P_0, \dots, P_5$ . With respect to  $P_0$  we will also assume  $X$  to be generic, in the precise sense of Proposition 2.4, and construct the plane curve  $C_X$  as in Section 2.1. The first observation is that under these hypotheses the curve  $C_X$  is a plane sextic with precisely five nodes. Thus using the well known formula for plane curves  $g = (d - 1)(d - 2)/2 - \delta$  where  $\delta$  is the number of nodes [see Arbarello et al., 1], we conclude immediately that the genus of  $C_X$  is five. Next we resolve the singularities of the curve by blowing up the plane at the five points  $P_0, \dots, P_5$ .

**Proposition 2.7.** *Let  $C$  be a plane curve of degree six with five ordinary nodes and let  $\widetilde{\mathbb{P}^2}$  be the blow-up of the projective plane in the five points corresponding to nodes of  $C$ . Then the strict transform of  $C$  inside  $\widetilde{\mathbb{P}^2}$  is a canonically embedded non-singular curve.*

*Proof.* According to Griffiths and Harris [22, §4.4], the blow-up of  $\mathbb{P}^2$  in five points is a Del Pezzo surface embedded in  $\mathbb{P}^4$  as the complete intersection of two quadrics by the complete linear system  $|-K_{\widetilde{\mathbb{P}^2}}|$ , where  $K_{\widetilde{\mathbb{P}^2}}$  is a canonical divisor in  $\widetilde{\mathbb{P}^2}$ . Thus the strict transform  $\widetilde{C}$  of  $C$  will be embedded in  $\mathbb{P}^4$  by the restriction of this linear system. Now let  $H$  be a line in  $\mathbb{P}^2$ , let  $E_1, \dots, E_5$  be the exceptional divisors in  $S$  and let  $\sigma$  be the blow-up morphism.

$$\begin{aligned} |-K_{\widetilde{\mathbb{P}^2}}| &= \left| \sigma^*(3H) - \sum_{i=1}^5 E_i \right| = \left| \sigma^*(-3H) + \sigma^*(6H) - \sum_{i=1}^5 E_i \right| \\ &= \left| \sigma^*(-3H) + \sigma^*(C) - \sum_{i=1}^5 E_i \right| \\ &= \left| \sigma^*(-3H) + \widetilde{C} + 2 \sum_{i=1}^5 E_i - \sum_{i=1}^5 E_i \right| \\ &= \left| \sigma^*(-3H) + \sum_{i=1}^5 E_i + \widetilde{C} \right| \\ &= |K_{\widetilde{\mathbb{P}^2}} + \widetilde{C}|. \end{aligned}$$

Therefore the curve  $\tilde{C}$  is embedded in  $\mathbb{P}^4$  by the restriction of the linear system  $|K_{\tilde{\mathbb{P}^2} + \tilde{C}}|$ , but by the adjunction formula this is just the canonical linear system of the curve. Observe also that  $\tilde{C}$  is non-singular because the singularities of  $C$  were only ordinary nodes.  $\square$

By Proposition 2.6 the blow-up of the double solid  $\Lambda_X$  at the point  $P_0$  is a conic bundle over  $\mathbb{P}^2$  branched along the curve  $C_X$ . Exploiting the existence of this conic bundle we can now define in a natural way an unbranched double cover of the normalisation of  $C_X$ .

**Proposition 2.8.** *Let  $X$  be a generic quartic hypersurface in  $\mathbb{P}^3$  with six isolated ordinary double points  $P_0, \dots, P_5$ , and let  $\tilde{C}_X$  be the canonical model of the associated plane sextic with respect to  $P_0$ . Then there exists a natural unbranched double cover of the curve  $\tilde{C}_X$ .*

*Proof.* Let  $W_X$  be the blow-up of the double solid  $\Lambda_X$  at the point  $P_0$  viewed as a conic bundle over  $\mathbb{P}^2$ , and let  $\widetilde{W}_X$  be the pull-back of  $W_X$  along the blow-up of  $\mathbb{P}^2$  at the five points  $P_1, \dots, P_5$  corresponding to the singularities of the curve  $C_X$ . In other words consider the following fibred product diagram

$$\begin{array}{ccc} \widetilde{W}_X & \longrightarrow & \widetilde{\mathbb{P}^2} \\ \downarrow & & \downarrow \sigma \\ W_X & \xrightarrow{f} & \mathbb{P}^2 \end{array}$$

By Proposition 2.7, the strict transform of the curve  $C_X$  inside  $\widetilde{\mathbb{P}^2}$  is the canonical model  $\tilde{C}_X$ . Abstractly we can realise  $\tilde{C}_X$  by blowing up  $C_X$  alone, the process is described for instance in Hartshorne [28, §II.7] or in Eisenbud and Harris [18, Proposition IV-21]. Let  $\Sigma$  be the preimage of the curve  $C_X$  inside the threefold  $W_X$  and consider also the blow up of  $\Sigma$  along the preimage of the five points  $P_1, \dots, P_5$ , which we call  $S$ . The situation is summarised by the following commutative cubic diagram: the base is the fibred product defining  $\Sigma$  as a closed subscheme of  $W_X$ ; the three “visible” vertical arrows are blow-ups and the vertical face whose base if  $f$  is the fibred product above. Note incidentally the existence and uniqueness of the dotted arrow, guaranteed by the universal property of the fibred diagram just mentioned. It would be interesting to

compare the top face of this cube with a fibred product, or in other words to compare  $S$  with the preimage of  $\widetilde{C}_X$  inside  $\widetilde{W}_X$ .

$$\begin{array}{ccccc}
 S & \xrightarrow{\exists!} & \widetilde{W}_X & & \\
 \downarrow \tilde{\sigma} & \searrow \tilde{f} & \downarrow & \searrow & \\
 \Sigma & \xrightarrow{\quad} & \widetilde{C}_X & \xrightarrow{\quad} & \widetilde{\mathbb{P}^2} \\
 & & \downarrow & & \downarrow \sigma \\
 & & W_X & \xrightarrow{f} & \mathbb{P}^2 \\
 & & \downarrow & & \\
 & & C_X & \xrightarrow{\quad} & \mathbb{P}^2
 \end{array}$$

Let  $\nu: \widetilde{S} \rightarrow S$  be the normalisation of  $S$ , and consider the Stein factorisation [Hartshorne, 28, Corollary III.11.5] of the morphism  $\tilde{f} \circ \nu$  to  $\widetilde{C}_X$

$$\begin{array}{ccc}
 \widetilde{S} & \xrightarrow{\nu} & S \\
 f' \downarrow & & \downarrow \tilde{f} \\
 \Gamma & \xrightarrow{g} & \widetilde{C}_X
 \end{array}$$

where  $f'$  is a projective morphism with connected fibres and  $g$  is a finite morphism. Evidently the degree of  $g$  is two, as the degenerate conic consists of two lines.

The double cover  $g$  is unbranched simply because the restriction of the conic bundle  $W_X$  to the curve  $C_X$  consists of a fibration by pairs of distinct lines. To see this recall from the previous section the local equation for the conic bundle, and construct the preimage of the point  $P$  of  $\mathbb{P}^2$  given by the maximal ideal  $(x - a, y - b) \subseteq \mathbb{K}[x, y]$ , which will be given by the form

$$-\zeta_2^2 + \zeta_1^2 g_2 + 2\zeta_1 \zeta_0 g_3 + \zeta_0^2 g_4$$

where  $g_d$  is the element of  $\mathbb{K}$  obtained by computing  $G_d(a, b)$  as in Proposition 2.2. The associated matrix is then the following

$$\begin{pmatrix} g_4 & g_3 & 0 \\ g_3 & g_2 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

Taking into account that we are assuming  $g_2, g_3$  and  $g_4$  never to be simultane-

ously null, this matrix will always have rank two, thus the quadratic polynomial will always consist of two distinct lines.  $\square$

It is still possible that the double cover  $g$  might be trivial, that is, it might consist of two copies of the curve  $\tilde{C}_X$ . Observe that this will be the case if and only if  $\Gamma$  is a disconnected curve, therefore reducible; but if  $\Gamma$  is reducible then  $\tilde{S}$  is also reducible. Taking into account that  $\tilde{S}$  is obtained from  $S$  by normalisation we see now that for the double cover to be non-trivial it is enough to show that  $S$  is irreducible, and since  $S$  is constructed by blowing up  $\Sigma$  it will be irreducible if and only if  $\Sigma$  is.

Going back again to the local equations of the conic bundle we are now interested in the divisor inside  $W_X$  defined by the equation  $u_3^2 - u_2u_4$ , which over the open set  $D_h(x_i)$  inside  $\mathbb{P}^2$  is given by the following ideal in  $\mathbb{K}[x, y][\tilde{\xi}_0, \tilde{\xi}_1, \tilde{\xi}_2]$

$$(-\tilde{\xi}_2^2 + \tilde{\xi}_1^2 G_2 + 2\tilde{\xi}_1 \tilde{\xi}_0 G_3 + \tilde{\xi}_0^2 G_4, G_3^2 - G_2 G_4)$$

where  $G_d$  is the polynomial obtained from  $u_d$  by multiplication by  $1/x_i^d$ . Note that in order to prove that  $\Sigma$  is irreducible we need to show that the ideal above is prime over any of the three standard open sets that cover  $\mathbb{P}^2$ , in other words we have to show that any of three ideals is prime.

If we denote by  $\mathcal{H}$  the Hilbert scheme of hypersurfaces in  $\mathbb{P}^3$  of degree four, then these ideals will be prime for  $X$  contained in some open subset of  $\mathcal{H}$ . We need only to check that this is not empty, then we shall have constructed a connected unbranched double cover of  $C_X$  for a general quartic surface  $X$ . The relevant computation was carried out with *Macaulay 2* and details can be found in Section 4.1.

## 2.4 An Explicit Example

Our purpose is now to compute an explicit example of a quartic hypersurface  $\mathbb{X}$  in  $\mathbb{P}^3$  with six ordinary double points. We will require  $\mathbb{X}$  to be generic in the sense of Proposition 2.4, and perform over it all the constructions described so far. Furthermore if later in the thesis we again require to specialise our arguments we want this to be our standard test example.

For ease of computation we work over the finite field  $\mathbb{F}_{101}$  (the prime 101 has no special importance here.) This is harmless in that if a variety in characteristic 0 is reducible so is its reduction modulo a prime  $p$ . Observe also that we shall have no restrictions in performing usual operations such as for instance computing derivatives to define the singular locus. Indeed the Jacobian criterion [Eisenbud 15, §16.6] holds over any perfect field and according to Zariski and Samuel [55, §II.4] these include  $\mathbb{F}_p$  for any prime  $p$ , as well as any algebraically closed field. All the computations described in this thesis were performed with *Macaulay 2* [21].

As usual we realise  $\mathbb{P}^3$  as the projective spectrum of the polynomial ring  $R = \mathbb{K}[x_0, \dots, x_3]$  and we fix the points  $P_0, \dots, P_5$  corresponding to the following prime ideals

$$\begin{aligned} \mathfrak{p}_0 &= (x_0, x_1, x_2), & \mathfrak{p}_1 &= (x_0, x_1, x_3), & \mathfrak{p}_2 &= (x_0, x_2, x_3), \\ \mathfrak{p}_3 &= (x_1, x_2, x_3), & \mathfrak{p}_4 &= (x_0 - x_1, x_0 - x_2, x_0 - x_3), \\ \mathfrak{p}_5 &= (2x_0 - x_1, 3x_0 - x_2, 4x_0 - x_3) \end{aligned}$$

The surface  $\mathbb{X}$  is then defined by a single irreducible homogeneous polynomial  $F$  of degree four contained in the intersection  $I = \bigcap_{i=0}^5 \mathfrak{p}_i^2$ , and as such it can be randomly selected. Thus we instruct the program to compute a system of generators for the ideal  $I$ , which will be returned in the form of the matrix associated to the homogeneous linear morphism whose image is the ideal  $I$

$$a: \bigoplus_{i=1}^{11} R(-d_i) \longrightarrow R$$

where  $d_i$  are the degrees of the generators. So the next step is to compute a random matrix of homogeneous polynomials in  $R$  such that when multiplied with the above the result is a form of degree four, or in other words a linear morphism

$$b: R(-4) \longrightarrow \bigoplus_{i=1}^{11} R(-d_i)$$

selected randomly for the composition  $a \circ b$  to be defined by a single polynomial of degree four  $F$ .

The result must now be tested, firstly to ensure its singular locus amounts to just the six double points and then to ensure it verifies the generic condition described in Proposition 2.4. The first is easily done by computing the derivatives of the polynomial  $F$  and asking for the Hilbert polynomial of the variety defined by  $(F, \partial F/\partial x_0, \dots, \partial F/\partial x_3)$  to be constant and equal to 6. For the second we need to write  $F$  in the form  $u_2x_3^2 + 2u_3x_3 + u_4$  and compute the Hilbert polynomial of  $(u_2, u_3, u_4)$ . Since these varieties are cones over  $P_0$  the condition is satisfied if they meet in  $P_0$  only, so we expect the Hilbert polynomial to be constant (in fact equal to 24). After these tests it appeared the following homogeneous forms satisfy our requirements.

$$\begin{aligned} u_2 &= 19x_0^2 - 33x_0x_1 + 50x_1^2 - 13x_0x_2 + 50x_1x_2 - 15x_2^2 \\ u_3 &= -2x_0^2x_1 - 35x_0x_1^2 - 18x_0^2x_2 - 8x_0x_1x_2 - 36x_1^2x_2 - 4x_0x_2^2 + 45x_1x_2^2 \\ u_4 &= -38x_0^2x_1^2 - 32x_0^2x_1x_2 - 32x_0x_1^2x_2 - 6x_0^2x_2^2 - 38x_0x_1x_2^2 + 2x_1^2x_2^2 \end{aligned}$$

Now we are left with the plane curve  $u_3^2 - u_2u_4 = 0$ , which we know has five ordinary nodes, and we want to compute its normalisation. In other words we want to compute the blow-up of the plane in five points, and to do so we can follow Griffiths and Harris [22, §4.4] and consider the linear system of cubic curves passing through the points. Details of this computation are described in §4.1, here we are more interested in the result which is the ideal in  $\mathbb{P}^4$  generated by the following three polynomials.

$$\begin{aligned} F_1 &= y_1y_2 + y_2^2 - y_0y_3 - 34y_2y_3 + 33y_2y_4 \\ F_2 &= y_0y_2 - 2y_2^2 - 33y_2y_3 - y_1y_4 + 35y_2y_4 \\ F_3 &= y_0^2 - 48y_0y_1 + 5y_1^2 + 12y_2^2 + 48y_0y_3 + 31y_1y_3 + 4y_2y_3 - 20y_3^2 + 42y_0y_4 + \\ &\quad - 50y_1y_4 - 45y_2y_4 + 17y_3y_4 - 19y_4^2 \end{aligned}$$

Thus the normalisation of the curve  $C_X$  is given by the intersection of three quadrics in  $\mathbb{P}^4$ , and is therefore canonically embedded. Note that this also proves that the singularities of the original plane curve are at worst nodes: indeed they are resolved by a single process of blow-up.

# Chapter 3

## Building Families

We are now in a position to describe a rational map  $\varrho$  from the space  $\mathcal{Q}$  of quartic hypersurfaces in  $\mathbb{P}^3$  with six isolated singularities to  $\mathcal{R}_5$ , the space of étale double covers of curves of genus five. Essentially this amounts to defining a natural way to construct families of curves from families of surfaces, but to make this precise we will need to use the formalism of category theory. Besides we will also show that  $\mathcal{Q}$  is *rational*, meaning that there exists a dominant birational map from a projective space  $\mathbb{P}^n$  to  $\mathcal{Q}$ , thus paving the way for the following result from which this thesis takes its title

**Theorem (3.14).** *The moduli space  $\mathcal{R}_5$  of étale two-sheeted coverings of curves of genus five is unirational.*

Unirationality is a weaker condition than rationality, in that it requires only the existence of a dominant rational map from  $\mathbb{P}^n$ . Therefore, on the last step in the proof of the theorem it will be sufficient to prove that  $\varrho$  is a dominant map. For simplicity we will always assume our schemes to be of finite type over  $\mathbb{C}$ .

### 3.1 Moduli Problems and Functors

A well known and widespread point of view is to identify a scheme  $X$  with its *functor of points*, which is defined by associating to any scheme  $Y$  the set of morphisms  $h_X(Y) = \text{Mor}(Y, X)$  and named from the observation that  $h_X(\text{Spec } K)$  is the set of *K-rational points* of  $X$ , for any field  $K$ . The construction is explained

in detail by Eisenbud and Harris [18] and consists of defining a fully faithful functor

$$h: (\text{Schemes}) \longrightarrow \text{Fun}((\text{Schemes})^\circ, (\text{Sets}))$$

from the category of schemes to the category of (contravariant) functors between schemes and sets, whose arrows are natural transformations [see for instance Mac Lane, 34, §II.4]. Thus any morphism  $u: X \rightarrow X'$  gives rise by composition to a natural transformation  $h(u)$  between the functors  $h_X$  and  $h_{X'}$ , with naturality square given by the following diagram

$$\begin{array}{ccc} Z & h_X(Z) & \xrightarrow{h(u)_Z} & h_{X'}(Z) \\ \alpha \uparrow & h_X(\alpha) \downarrow & & \downarrow h_{X'}(\alpha) \\ Y & h_X(Y) & \xrightarrow{h(u)_Y} & h_{X'}(Y) \end{array}$$

In fact the construction of  $h$  makes sense in any category, not just  $(\text{Schemes})$ , and it is a basic result in category theory that it always gives rise to a fully faithful functor. The following is the precise statement according to Eisenbud and Harris [18, Lemma VI-1].

**Lemma 3.1** (Yoneda). *Let  $\mathcal{C}$  be a category and let  $X, X'$  be objects of  $\mathcal{C}$ .*

- (a) *If  $F$  is any contravariant functor from  $\mathcal{C}$  to the category of sets, the natural transformations from  $\text{Mor}(-, X)$  to  $F$  are in natural correspondence with the elements of  $F(X)$*

$$\text{Nat}(\text{Mor}(-, X), F) \longleftrightarrow F(X)$$

- (b) *If the functors  $\text{Mor}(-, X)$  and  $\text{Mor}(-, X')$  from  $\mathcal{C}$  to the category of sets are isomorphic, then  $X \cong X'$ . More generally, the maps of functors from  $\text{Mor}(-, X)$  to  $\text{Mor}(-, X')$  are the same as maps from  $X$  to  $X'$*

$$\text{Nat}(\text{Mor}(-, X), \text{Mor}(-, X')) \longleftrightarrow \text{Mor}(X, X'),$$

*that is, the functor  $h: \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^\circ, (\text{Sets}))$  sending  $X$  to  $h_X$  is fully faithful.*

Posing a moduli problem amounts to defining a (contravariant) functor  $F$  from the category of schemes to  $(\text{Sets})$  by specifying what *families* of objects to consider. In general, for any scheme  $B$ , a family will be a morphism  $f: \mathcal{X} \rightarrow B$

with some properties depending on the particular problem, but always with fibres required to be of a certain kind. To any morphism of schemes  $B \rightarrow B'$  a moduli functor will then associate an arrow defined by pulling-back families or in other words by taking fibred products. Given a moduli functor, or also given any  $F: (\text{Schemes})^\circ \rightarrow (\text{Sets})$ , it is now natural to ask whether it is of the form  $h_X$  for some scheme  $X$  or not, and if it is to give it a special name. In this case we say that  $F$  is *representable* by  $X$  or that  $X$  *represents*  $F$ .

**Definition.** If  $F$  is representable by  $\mathcal{M}$ , then we say that the scheme  $\mathcal{M}$  is a *fine moduli space* for the moduli problem  $F$ .

The classic example of a fine moduli space is the Hilbert scheme, parametrising closed subschemes of  $\mathbb{P}^r$  with fixed Hilbert polynomial  $P = P(m)$ . The moduli problem is first presented formally by defining the functor  $H_{P,r}$ , whose value on  $B$  is the set of proper flat families

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{i} & \mathbb{P}^r \times B \\ & \searrow \varphi & \downarrow \\ & & B \end{array}$$

with  $\mathcal{X}$  having Hilbert polynomial  $P$ , and  $i$  being a closed immersion. The main result, due to Grothendieck [24], is then the following.

**Theorem 3.2** (Grothendieck).  $H_{P,r}$  is representable by a projective scheme  $\mathcal{H}_{P,r}$ .

Fine moduli spaces are characterised by the existence of a *universal family*, that is a family  $\mathcal{U} \rightarrow \mathcal{M}$  from which every other one can be obtained. To see this observe that the functor  $F$  is representable by  $\mathcal{M}$  if and only if morphisms in  $h_{\mathcal{M}}(B)$  are in bijection with families in  $F(B)$ . Thus a family over  $B$  is the same as a morphism  $\alpha \in h_{\mathcal{M}}(B)$ , which in turn is given by the image of the identity under  $h_{\mathcal{M}}(\alpha)$ .

$$\begin{array}{ccc} \mathcal{M} & F(\mathcal{M}) \longleftrightarrow & h_{\mathcal{M}}(\mathcal{M}) \\ \alpha \uparrow & F(\alpha) \downarrow & \downarrow h_{\mathcal{M}}(\alpha) \\ B & F(B) \longleftrightarrow & h_{\mathcal{M}}(B) \end{array}$$

The universal family is then the unique element in  $F(\mathcal{M})$  corresponding to the identity of  $\mathcal{M}$ .

As an example consider the moduli space of hypersurfaces in  $\mathbb{P}^r$  of degree  $d$ . If  $X$  is a projective variety contained in  $\mathbb{P}^r$  then it is well known (see for instance Harris and Morrison [26], §1.B) that  $X$  is a hypersurface of degree  $d$  if and only if its Hilbert polynomial is the following

$$P_X(m) = \binom{r+m}{r} - \binom{r+m-d}{r}$$

Thus this moduli space is in fact the Hilbert scheme  $\mathcal{H}_{P,r}$ , and it is given by the projective space  $\mathbb{P}(V)$ , where  $V$  is the vector space of homogeneous polynomials of degree  $d$  in  $r+1$  variables. The universal family is the closed subset of the product  $\mathbb{P}(V) \times \mathbb{P}^r$  defined by the equations  $\{(F, P) \mid F(P) = 0\}$ .

Unfortunately, the existence of a universal family is a requirement too restrictive for many natural moduli problems; in other words there are examples of moduli functors which are not representable but nevertheless describe interesting classes of objects. Perhaps the most famous of these is the functor describing families of smooth projective curves of genus  $g$ . One way of getting around this problem is to look for the existence of a scheme  $\mathcal{M}$  that “best represents” the functor, satisfying a weaker universal property.

**Definition.** A scheme  $\mathcal{M}$  and a natural transformation  $\Phi$  from  $F$  to the functor of points of  $\mathcal{M}$  are a *coarse moduli space* for the moduli problem  $F$  if

- 1) The map  $\Phi_{\mathbb{K}}: F(\text{Spec } \mathbb{K}) \rightarrow h_{\mathcal{M}}(\text{Spec } \mathbb{K})$  is a set bijection;
- 2) For any scheme  $\mathcal{M}'$  and any natural transformation  $\Psi: F \rightarrow h_{\mathcal{M}'}$ , there is a unique morphism  $u: \mathcal{M} \rightarrow \mathcal{M}'$  such that  $\Psi = h(u) \circ \Phi$ .

The moduli functor  $\mathbf{m}_g$  is defined by considering smooth proper morphisms  $\mathcal{X} \rightarrow B$  whose fibres are nonsingular irreducible projective curves of genus  $g$ .

**Theorem 3.3.** *Let  $g \geq 2$ . The functor  $\mathbf{m}_g$  is coarsely represented by a quasi-projective scheme  $\mathcal{M}_g$ , irreducible and  $(3g-3)$ -dimensional.*

The problem was first considered in the complex case by Riemann [47], who proved that the space of non isomorphic complex structures (named by

him “moduli”) over a compact connected surface of genus  $g \geq 2$  has complex dimension  $3g - 3$ . However in this generality the result is explained in detail by, and mostly due to, Deligne and Mumford [11].

**Lemma 3.4.** *Let  $\mathcal{F}$  be a fine moduli space for the moduli problem  $\mathbf{F}$ , and let  $\mathcal{M}$  be a coarse moduli space for the moduli problem  $\mathbf{m}$ . Then to give a morphism of schemes  $\mathcal{F} \rightarrow \mathcal{M}$  it is enough to give a natural transformation  $\mathbf{F} \rightarrow \mathbf{m}$ .*

*Proof.* This is just an application of Yoneda’s Lemma (3.1). Indeed by definition there are natural transformations  $\mathbf{m} \rightarrow h_{\mathcal{M}}$  and  $\mathbf{F} \rightarrow h_{\mathcal{F}}$ , the last of which is assumed to be invertible, so composition with another one from  $\mathbf{F}$  to  $\mathbf{m}$  will define a natural transformation between the functors  $h_{\mathcal{F}}$  and  $h_{\mathcal{M}}$ .  $\square$

This lemma is used in practice to define rational maps between moduli spaces. Indeed what generally happens is that there exists a Zariski open subset of a coarse moduli space over which the moduli functor will become representable. This is the case for instance of curves, where the open set is given by the locus of curves without automorphisms.

Consider now the functor  $\mathbf{r}_g$  ( $g \geq 2$ ) defined over the scheme  $B$  by considering families of smooth projective curves of genus  $g$  with a connected étale double cover. According to Beauville [3, §6] this functor is coarsely represented by an irreducible quasi-projective scheme  $\mathcal{R}_g$ . The dimension of this moduli space is  $3g - 3$ , the same as the dimension of  $\mathcal{M}_g$ . To see this recall that double covers of curves correspond one to one to non-trivial line bundles  $\mathcal{L}$  whose square  $\mathcal{L} \otimes \mathcal{L}$  is trivial (compare Hartshorne [28], Exercise IV.2.7), so the families considered by the functor  $\mathbf{r}_g$  are just families of curves with an extra structure. Such a line bundle on a smooth irreducible projective curve  $C$  of genus  $g$  corresponds to one of the  $2^{2g} - 1$  nontrivial 2-torsion points in the Jacobian of  $C$ . So forgetting about the extra information we clearly define a natural transformation between  $\mathbf{r}_g$  and  $\mathbf{m}_g$ , and applying Lemma 3.4 we then obtain a (at least rational) map  $\mathcal{R}_g \rightarrow \mathcal{M}_g$  with finite fibres.

### 3.2 The Moduli Space of Quartic Surfaces

Let  $\mathcal{Q}$  be the moduli space of “quartic hypersurfaces in  $\mathbb{P}^3$  with six isolated ordinary double points, one of which is marked, up to projective equivalence.”

As described in Section 3.1 this is equivalent to considering the moduli functor  $\mathcal{Q}$  whose value on the scheme  $B$  is the set of proper flat families

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{i} & \mathbb{P}^r \times B \\ & \searrow \varphi & \downarrow \\ & & B \end{array}$$

where  $i$  is a closed immersion and for every  $b \in B$  the fibre of  $\varphi$  over the point  $b$  is a quartic hypersurface as described above, but over the residue field  $K(b) = \mathcal{O}_{B,b}/\mathfrak{m}_b$ . Our first goal is to prove that this is in fact a fine moduli space, or in other words that this functor is representable by a scheme  $\mathcal{Q}$ .

**Proposition 3.5.**  *$\mathcal{Q}$  is a fine moduli space. More precisely it is a locally closed subscheme of the Hilbert scheme of quartic hypersurfaces in  $\mathbb{P}^3$ .*

*Proof.* To begin with consider the Hilbert scheme of quartic hypersurfaces in  $\mathbb{P}^3$ . This is the fine moduli space of projective varieties contained in  $\mathbb{P}^3$  with Hilbert polynomial

$$P_X(m) = \binom{3+m}{3} - \binom{m-1}{3} = 2m^2 + 2$$

and can be realised as  $\mathbb{P}(R_4)$ , where  $R$  is the polynomial ring  $\mathbb{K}[x_0, \dots, x_3]$ , or in other words as  $\mathbb{P}(H^0(\mathbb{P}^3, \mathcal{O}(4)))$ .

However we are interested in surfaces up to projective equivalence, so to take into account the action of  $PGL(4)$  we fix the five points in  $\mathbb{P}^3$  corresponding to the following prime ideals

$$\begin{aligned} \mathfrak{p}_0 &= (x_0, x_1, x_2), & \mathfrak{p}_1 &= (x_0, x_1, x_3), & \mathfrak{p}_2 &= (x_0, x_2, x_3), \\ \mathfrak{p}_3 &= (x_1, x_2, x_3), & \mathfrak{p}_4 &= (x_0 - x_1, x_0 - x_2, x_0 - x_3) \end{aligned}$$

So we are now looking at “quartic hypersurfaces in  $\mathbb{P}^3$  with five isolated ordinary double points, up to projective equivalence.” This is again a fine moduli space, in that it is the closed subscheme of the Hilbert scheme given by  $\mathbb{P}(I_4)$  where  $I = \bigcap_{i=0}^4 \mathfrak{p}_i^2$  is the ideal of the five double points. In other words we are looking at the linear space  $H^0(\mathbb{P}^3, \mathcal{I}(4))$ , where we denote  $\mathcal{I}$  the sheaf of ideals defined by  $I$ , inside  $H^0(\mathbb{P}^3, \mathcal{O}(4))$ , and then realising a projective space

with it. Observe that the dimension  $h^0(\mathbb{P}^3, \mathcal{I}(4))$  gives the value of the Hilbert function of  $\mathcal{I}$  in degree four, which a direct computation shows to be 15.

Going on, we ask now for a sixth double point. To realise this we can take the product  $\mathbb{P}(I_4) \times \mathbb{P}^3$  and consider the closed subscheme  $B_0$  given by the relations

$$\{(F, \mathfrak{p}) \mid F \in \mathfrak{p}^2\} = \left\{ (F, P) \mid F(P) = \frac{\partial F}{\partial x_i}(P) = 0 \right\}$$

Projecting onto  $\mathbb{P}(I_4)$  we consider now the scheme theoretic image of  $B_0$  (see §1.1) which is a closed subscheme  $B$  of  $\mathbb{P}(I_4)$ , and observe that in the universal factorisation

$$\begin{array}{ccc} B_0 & \xrightarrow{\beta} & B \\ & \searrow & \downarrow \\ & & \mathbb{P}(I_4) \end{array}$$

the dominant morphism  $\beta$  is also proper, because it is the external morphism of a composition which is proper [Hartshorne, 28, Exercise II.4.8]. As a consequence  $\beta$  is surjective and the scheme theoretic image of  $B_0$  coincides with the set theoretic image. Observe also that  $B$  is irreducible by Corollary 1.4. However  $B$  contains all the possible degenerations of a quartic hypersurface with six double points, while we are interested in those hypersurfaces with no other singularities. As this is clearly an open condition we have proved that  $\mathcal{Q}$  is an open subset of an irreducible closed subset of the Hilbert scheme, and thus it is a fine moduli space.  $\square$

The proof above in fact reveals something more about  $\mathcal{Q}$ , namely we now know its dimension and most of all we know that it is a *unirational* space. The last property in particular is of primary importance for the purposes of this thesis.

**Corollary 3.6.**  *$\mathcal{Q}$  is a 13-dimensional, unirational variety.*

*Proof.* We have seen in the proof of the Proposition above that there is a dominant rational morphism from a closed subscheme  $B_0$  of the product  $\mathbb{P}(I_4) \times \mathbb{P}^3$  to  $\mathcal{Q}$ , given by the projection onto  $\mathbb{P}(I_4)$ . Now focus on the other projection and observe that a generic point  $P \in \mathbb{P}^3$  defines four independent conditions on the linear space  $H^0(\mathbb{P}^3, \mathcal{I}(4))$ , thus there exists an open subset  $U$  of  $\mathbb{P}^3$

over which these conditions define a vector bundle  $E$  of rank 11. To see that  $U$  is not empty it is enough to fix a sixth point in  $\mathbb{P}^3$  and compute the Hilbert function of the ideal  $J = \bigcap_{i=0}^5 \mathfrak{p}_i^2$ , which we have already done in Section 2.4 with respect to the point  $\mathfrak{p}_5 = (2x_0 - x_1, 3x_0 - x_2, 4x_0 - x_3)$ . The projective space bundle  $B_1$  over  $U$  associated to  $E$  is a rational variety embedded in  $B_0$  as a dense open subset, and mapping dominantly onto  $\mathcal{Q}$ , which is therefore at least unirational.

Since  $\mathcal{Q}$  is irreducible, because we have seen in the theorem that it is an open subset of an irreducible variety, to complete the proof it is now enough to compute the differential of the projection at one general point and show that the dimension of  $\mathcal{Q}$  is 13. To this purpose the following procedure was applied to the standard test example of Section 2.4 (see Section 4.2 for details). Work over an affine subset of the product, where  $B_0$  is given by four equations  $F_1, \dots, F_4$  inside the polynomial ring  $\mathbb{K}[t_1, \dots, t_{17}]$ , and realise the tangent space of  $B_0$  at the point  $P$  as the linear space in  $\mathbb{A}^{17}$  corresponding to the ideal generated by the Jacobian equations  $\sum_j \partial F_i / \partial x_j|_P t_j$ . Then find the intersection of this ideal with the subalgebra  $\mathbb{K}[t_1, \dots, t_{14}]$ .  $\square$

Our goal is now to define a rational morphism from  $\mathcal{Q}$  to the moduli space  $\mathcal{R}_5$  of double covers of curves of genus five. Applying Lemma 3.4 it is enough to define a natural way of constructing families of curves from families of surfaces; natural in the precise sense of category theory, that is commuting with fibred products.

**Proposition 3.7.** *There exists a morphism of schemes  $\varrho: \mathcal{Q} \rightarrow \mathcal{R}_5$ , from  $\mathcal{Q}$  to the moduli space of double covers of curves of genus five.*

*Proof.* Recall the general constructions described earlier in the thesis about how to associate to any nodal quartic hypersurface in  $\mathbb{P}^3$  a nodal sextic plane curve with a double cover.

We now use a gluing argument, starting off assuming the base scheme to be the spectrum of a ring  $A$ . In this case we can proceed as we did before, associating to the scheme  $\mathcal{X} = \text{Proj } A[x_0, \dots, x_3] / (u_2 x_3^2 + 2u_3 x_3 + u_4)$  the plane curve over  $A$  defined by the equation  $u_3^2 - u_2 u_4$ . This association is also natural, in that it commutes with pull-backs. Indeed for any homomorphism of

rings  $A \rightarrow A'$  the pull-back of the family of curves just defined is given by the projective spectrum of the graded ring

$$(A \otimes A')[x_0, x_1, x_2]/(u_3^2 - u_2u_4)$$

which is the same graded ring one would obtain by first pulling back the family of surfaces and then applying the correspondence.  $\square$

We want to use the morphism described above to prove the unirationality of  $\mathcal{R}_5$ , in other words, in view of the other results, we want to prove that  $\varrho$  is dominant. To this purpose we now describe a first simplification of the problem. As we have seen in Section 3.1 the space  $\mathcal{R}_5$  maps dominantly onto  $\mathcal{M}_5$ , the moduli space of curves of genus five, so we can consider the composition of morphisms to get the commutative diagram

$$\begin{array}{ccc} \mathcal{Q} & \xrightarrow{\varrho} & \mathcal{R}_5 \\ & \searrow \theta & \downarrow \eta \\ & & \mathcal{M}_5 \end{array}$$

We now have the following result.

**Lemma 3.8.** *The morphism  $\varrho$  is dominant if and only if  $\theta$  is dominant.*


*Proof.* Given that  $\eta$  is a dominant morphism between irreducible spaces, if  $\varrho$  is also a dominant morphism the conclusion follows immediately. To see the converse it is probably more intuitive to proceed by contradiction. If  $\varrho$  was not dominant then the scheme theoretic image of  $\mathcal{Q}$  inside  $\mathcal{R}_5$  should be of lower dimension than  $\dim \mathcal{R}_5 = 12$ , because  $\mathcal{R}_5$  is irreducible (see Section 3.1). But in this case the scheme theoretic image of the composition  $\eta \circ \varrho$  would also be of lower dimension than 12. This is clearly not possible because assuming  $\theta$  to be dominant is equivalent to assuming the dimension of its image to be 12, because  $\mathcal{M}_5$  is also irreducible.  $\square$

To complete the argument we must now focus our attention to the morphism  $\theta$ . In particular we look for sufficient conditions for it to be dominant, i.e. generically surjective.

**Lemma 3.9.** *Let  $u: X \rightarrow X'$  be a morphism of algebraic schemes, with  $X'$  irreducible. Then the scheme theoretic image of  $u$  is the whole of  $X'$  if and only if there exists a point  $P \in X$  such that the differential  $du_P: T_{X,P} \rightarrow T_{X',u(P)}$  is surjective.*

*Proof.* Since  $X'$  is irreducible the scheme theoretic image of  $u$  will be the whole of  $X'$  if and only if the dimension of at least one of its irreducible components equals the dimension of  $X'$ . Now it is enough to recall that the dimension of the irreducible component of  $\overline{u(X)}$  containing a regular point  $u(P)$  is given by the rank of the differential.  $\square$

Following Harris and Morrison [26, §1.C] we can realise the tangent space to any scheme  $X$  at a closed point  $P$  as the set of maps from  $D = \text{Spec } \mathbb{K}[\varepsilon]/(\varepsilon^2)$  to  $X$  centred at  $P$  (that is, mapping the unique closed point  $\mathbf{0} \in D$  to  $P$ ). The linear operations on vectors are then given by means of fibred products, in that for any  $\alpha, \beta \in \mathbb{K}$  the vector  $\alpha\varphi + \beta\psi$  will be defined as



For any morphism  $u: X \rightarrow X'$  and any closed regular point  $P \in X$  the differential  $du_P: T_{X,P} \rightarrow T_{X',u(P)}$  will now be simply given by the composition of morphisms  $\varphi \mapsto u \circ \varphi$ .

**Lemma 3.10.** *Let  $u: X \rightarrow X'$  be a morphism of schemes, and let  $P \in X$  be a closed point. Then the differential of  $du$  of  $u$  is given by the restriction of the natural transformation  $h(u): h_X \rightarrow h_{X'}$  over  $D$  to the tangent space of  $X$  in the sense above.*

*Proof.* To see this we just need to unravel the definitions. In accordance with the previous discussion the tangent space of  $X$  at  $P$  is in the first place a subset of  $\text{Mor}(D, X)$ , and the differential operates on this subset precisely as does  $h(u)_D: \text{Mor}(D, X) \rightarrow \text{Mor}(D, X')$ .  $\square$

### 3.3 Families of Curves and their Dimension

Let  $C$  be a canonically embedded curve of genus five, which we assume to be given by the complete intersection of three quadrics in  $\mathbb{P}^4$ . According to *Petri's Theorem* (see Arbarello, Cornalba, Griffiths and Harris [1], §III.3) we are thus assuming  $C$  to be canonically embedded and not trigonal.

**Theorem 3.11** (Petri). *Let  $C$  be a canonical curve of genus 5. Then the ideal of  $C$  is generated by three quadrics, unless  $C$  is trigonal in which case it is generated by three quadrics and three cubics.*

Given a family  $\mathcal{F}$  of deformations of  $C$ , we want to know its dimension in terms of non-isomorphic curves, more precisely we look for a sufficient condition for its base to be generically surjective onto the moduli space  $\mathcal{M}_5$  of genus five curves. To do this we can follow Serresi [49] and realise  $\mathcal{F}$  as a fibred product diagram

$$\mathcal{F}: \begin{array}{ccc} C & \longrightarrow & \mathcal{C} \\ \downarrow & & \downarrow f \\ \text{Spec } \mathbb{K} & \xrightarrow{s} & S \end{array}$$

where  $f$  is flat and surjective, and  $S$  is the scheme of parameters, that is the base of the family. The question will then reduce to the computation of the *Kodaira-Spencer* map of  $\mathcal{F}$ , in that all the possible deformations of  $C$  are parametrised by the cohomology group  $H^1(C, T_C)$  and there exists a linear map  $\kappa_{\mathcal{F}}: T_{S,s} \rightarrow H^1(C, T_C)$  whose image consists precisely of those deformations contained in  $\mathcal{F}$ . Here  $H^1(C, T_C)$  should be thought of as the tangent space to  $\mathcal{M}_5$  at the point corresponding to  $C$ .

Observe that  $H^1(C, T_C)$  is isomorphic to the vector space  $H^0(C, K_C^{\otimes 2})^*$ , indeed the tangent sheaf  $T_C$  is dual to the canonical sheaf  $K_C$  and by Serre duality we have

$$H^1(C, \mathcal{F}) = H^0(C, \mathcal{F}^{\vee} \otimes K_C)^*$$

for any invertible sheaf  $\mathcal{F}$ . Since we are assuming the curve to be canonically embedded the sheaf  $K_C^{\otimes 2}$  will be in this case just the twisted sheaf  $\mathcal{O}_C(2)$ , thus if  $I$  is the ideal inside  $R = \mathbb{K}[x_0, \dots, x_4]$  defining the curve  $C$  the cohomology group above will be given by the quotient  $R_2/I_2$ .

We need to investigate necessary and sufficient conditions for two such curves to be isomorphic. First we observe that two canonically embedded curves are isomorphic if and only if they are projectively equivalent. Thus we are now reduced to deciding how many projectively equivalent curves there are among the fibres of the morphism  $f$ .

**Lemma 3.12.** *Two canonically embedded curves of genus  $g$  are isomorphic if and only if they are projectively equivalent.*

*Proof.* A consequence of Riemann-Roch is that the only linear system  $g_{2g-2}^{g-1}$  of degree  $2g - 2$  and dimension  $g - 1$  on a curve of genus  $g$  is the canonical series [Arbarello et al., 1, p.10]. Therefore if  $\varphi: C_1 \rightarrow C_2$  is an isomorphism, we can conclude that the pull-back of the canonical series of  $C_2$  is the canonical series of  $C_1$ . But then we can define an automorphism of  $\mathbb{P}^{g-1}$  inducing  $\varphi$ , concluding that  $C_1$  and  $C_2$  are projectively equivalent.  $\square$

In terms of projective equivalence we can realise a model containing all the possible canonical curves of genus five by considering the Grassmannian of dimension three subspaces of the vector space  $R_2$ . Projective equivalence is then given by the action of the group  $SL(5)$  and we are looking at the space of the orbits.

But the Grassmannian itself can be realised as an orbit space, this time under the action of  $GL(3)$ , as follows. Let  $V$  be the open set inside the 45-dimensional vector space  $R_2 \times R_2 \times R_2$  given by all the possible bases of dimension three subspaces in  $R_2$ , and consider the action of  $GL(3)$  whose orbits are all the possible bases for a given subspace. This is the obvious action

$$M(\underline{x}Q_1\underline{x}, \underline{x}Q_2\underline{x}, \underline{x}Q_3\underline{x}) = \left( \sum_{i=1}^3 m_{1i} \underline{x}Q_i\underline{x}, \sum_{i=1}^3 m_{2i} \underline{x}Q_i\underline{x}, \sum_{i=1}^3 m_{3i} \underline{x}Q_i\underline{x} \right),$$

where  $Q_i$  is a symmetric matrix and  $\underline{x}$  is the row vector  $(x_0, \dots, x_4)$ . We now consider the action of  $SL(5)$  whose orbits are all the possible projectively equivalent bases, which is the following

$$N(\underline{x}Q_1\underline{x}, \underline{x}Q_2\underline{x}, \underline{x}Q_3\underline{x}) = (\underline{x} \underline{N}Q_1 \underline{N}\underline{x}, \underline{x} \underline{N}Q_2 \underline{N}\underline{x}, \underline{x} \underline{N}Q_3 \underline{N}\underline{x})$$

It is important to observe that the two actions commute, thus they don't interfere with each other and we can regard one as acting on the orbit space of the other.

In order to investigate properties of a family of deformations of the curve  $C$  around its central fibre it is enough to consider the case in which the base scheme is affine and given by the spectrum of the ring  $A = \mathbb{K}[t_0, \dots, t_N]/\mathfrak{m}^2$ , where  $\mathfrak{m}$  is the maximal ideal generated by  $t_0, \dots, t_N$  which is assumed to correspond (as a point of  $\text{Spec } A$ ) to the curve  $C$ . In this situation the scheme  $\mathcal{C}$  will be defined by the projective spectrum of  $A[x_0, \dots, x_4]/(F_1, F_2, F_3)$ , where the coefficients of  $F_i$  depend linearly on the parameters  $t_i$ . In other words we can write every polynomial  $F_i$  as a sum

$$F_i = H_i + \sum_{j=0}^N t_j H_{ij},$$

where  $H_i$  and  $H_{ij}$  are quadric polynomials inside  $\mathbb{K}[x_0, \dots, x_4]$ . The  $N$  triplets of quadric polynomials  $(H_{1j}, H_{2j}, H_{3j})$  generate the linear space inside  $\mathbb{A}^{45}$  tangent to the family at the central point  $s = (H_1, H_2, H_3)$ . The problem is to determine whether or not this linear space fills the tangent space to  $\mathcal{M}_5$ ; the strategy we are going to adopt is to work inside the tangent space to  $V$  at  $s$ , constructing a basis for all the trivial deformations and then check how many of the above triplets lie inside this linear space; keeping in mind that the trivial deformations are those given by the group actions described above.

Around any point  $v = ({}^t x Q_1 x, {}^t x Q_2 x, {}^t x Q_3 x)$  in  $V$  the action of the two groups is linearised by the action of the corresponding Lie algebras, so a system of generators for the linear space tangent to the orbit passing through  $v$  is simply determined by applying a basis for the Lie algebra to it. Now the Lie algebra  $\mathfrak{gl}(3)$  is simply the whole space of three-by-three matrices and its action is the same as the one of  $GL(3)$ , thus we obtain a first set of trivial deformations given by the nine vectors

$$(H_1, 0, 0), (H_2, 0, 0), \dots, (0, 0, H_2), (0, 0, H_3)$$

The algebra  $\mathfrak{sl}(5)$  instead is given by the vector space of traceless  $5 \times 5$  ma-

trices [see Fulton and Harris, 20, §8.2] and its action is determined as follows

$$\begin{aligned} {}^t(N\underline{x})Q_i(N\underline{x}) &= {}^t\underline{x}{}^t(I + \varepsilon\Delta)Q_i(I + \varepsilon\Delta)\underline{x} \\ &= {}^t\underline{x}(Q_i + \varepsilon({}^t\Delta Q_i + Q_i\Delta))\underline{x} \\ &= {}^t\underline{x}Q_i\underline{x} + \varepsilon\underline{x}({}^t\Delta Q_i + Q_i\Delta)\underline{x} \end{aligned}$$

thus letting  $\Delta$  vary among a basis for  $\mathfrak{sl}(5)$  we get another set of trivial deformations given by the 24 vectors

$$({}^t\Delta Q_1 + Q_1\Delta, {}^t\Delta Q_2 + Q_2\Delta, {}^t\Delta Q_3 + Q_3\Delta).$$

Here one should first write the associated symmetric matrix  $Q_i$  to each of the polynomials  $H_i$ ; to keep this in mind we can write each of the entries above rather improperly as  ${}^t\Delta H_i + H_i\Delta$ .

Construct now the matrix  $M_{\mathcal{F}}$  as follows.

$$M_{\mathcal{F}} : \begin{pmatrix} H_{11} & H_{21} & H_{31} \\ H_{12} & H_{22} & H_{32} \\ \vdots & \vdots & \vdots \\ H_{1N} & H_{2N} & H_{3N} \\ H_1 & 0 & 0 \\ H_2 & 0 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & H_2 \\ 0 & 0 & H_3 \\ D_{21}H_1 + H_1D_{12} & D_{21}H_2 + H_2D_{12} & D_{21}H_3 + H_3D_{12} \\ D_{31}H_1 + H_1D_{13} & D_{31}H_2 + H_2D_{13} & D_{31}H_3 + H_3D_{13} \\ \vdots & \vdots & \vdots \\ D_{55}H_1 + H_1D_{55} & D_{55}H_2 + H_2D_{55} & D_{55}H_3 + H_3D_{55} \end{pmatrix}$$

The first  $N$  rows are given by the family  $\mathcal{F}$ , they are tangent vectors at the central point  $s = (H_1, H_2, H_3)$ . The second set of 9 rows is given by the tangent vectors to the orbit of the  $GL(3)$ -action. The last 24 rows are the tangent vectors to the orbits of the  $SL(5)$ -action described above. We have considered here the

basis for the vector space of traceless  $5 \times 5$  matrices given by the matrix  $D_{ij}$  with a unique nonzero entry (which will be a 1) in position  $ij$  for  $i \neq j$  and

$$D_{22} = \text{diag}(1, -1, 0, 0, 0)$$

$$D_{33} = \text{diag}(1, 0, -1, 0, 0)$$

$$D_{44} = \text{diag}(1, 0, 0, -1, 0)$$

$$D_{55} = \text{diag}(1, 0, 0, 0, -1)$$

The rows of the matrix  $M_{\mathcal{F}}$  generate a linear space, which is the span inside the tangent space to  $V$  of the three linear spaces tangent respectively to the given family and to each of the two orbits through  $s$ . To determine the dimension of this span we now need to compute the rank of  $M_{\mathcal{F}}$ .

**Proposition 3.13.** *Let  $C$  be the complete intersection of three quadrics in  $\mathbb{P}^4$ , and let  $\mathcal{F}$  be a family of deformations of  $C$  as above. Assume also that the dimension of  $\mathcal{F}$  is at least 12, in other words  $N \geq 12$ . If the rank of the matrix  $M_{\mathcal{F}}$  is maximal then the Kodaira-Spencer map of  $\mathcal{F}$  is surjective.*

*Proof.* First observe that in the matrix  $M_{\mathcal{F}}$  there are 45 columns, and under our assumptions there are at least 45 rows. When the rank of the matrix  $M_{\mathcal{F}}$  is maximal the span of the three vector spaces, the two corresponding to trivial deformations and the one given by the family, is the whole of the tangent space to  $V$  at the point  $s$ . Thus we are guaranteed the existence of enough linearly independent deformations, namely 12, to fill the tangent space to  $\mathcal{M}_5$ .  $\square$

Looking back now at the morphism  $\varrho: \mathcal{Q} \rightarrow \mathcal{R}_5$  defined in Proposition 3.7 we put everything together and show how the matrix  $M_{\mathcal{F}}$  described above can help us proving the unirationality of  $\mathcal{R}_5$ . We have already observed in Lemma 3.8 that it is enough to prove that the morphism  $\theta: \mathcal{Q} \rightarrow \mathcal{M}_5$  is generically surjective, and in Lemma 3.9 we have reduced the argument to showing that the differential of  $\theta$  at one point is surjective. By Lemma 3.4 the morphism  $\theta$  corresponds to a natural transformation

$$h(\theta): h_{\mathcal{Q}} \longrightarrow h_{\mathcal{M}_5}$$

that we actually have defined in Proposition 3.7.

**Theorem 3.14.** *The moduli space  $\mathcal{R}_5$  of étale two-sheeted coverings of curves of genus five is unirational.*

*Proof.* We fix a regular point  $\mathbb{X}$  inside  $\mathcal{Q}$ , and compute the differential of  $\theta$  at  $\mathbb{X}$  as in Lemma 3.10 by restricting the natural transformation over  $D$

$$h(\theta)_D: \mathbf{Q}(D) \longrightarrow \mathbf{m}_5(D)$$

To this purpose we start with a particular family of deformations of  $\mathbb{X}$ , the one we have defined in Proposition 3.5. Recall the definition of  $B_0 \subseteq \mathbb{P}^{14} \times \mathbb{P}^3$  and consider the family  $\mathcal{F}$  given by the following fibred product diagram

$$\mathcal{F}: \begin{array}{ccc} \mathbb{X} & \longrightarrow & \mathcal{S} \\ \downarrow & & \downarrow f \\ \text{Spec } \mathbb{K} & \xrightarrow{b} & B_0 \end{array}$$

where  $b$  is the point of  $B_0$  corresponding to  $\mathbb{X}$ . Pulling back  $\mathcal{F}$  along a basis of vectors  $\varphi_i: D \rightarrow B_0$  ( $i = 1, \dots, 13$ ) for the tangent space of  $B_0$  at  $b$  we obtain linearly independent elements of the tangent space  $T_{\mathcal{Q}, \mathbb{X}} \subset \mathbf{Q}(D)$ , which by the way will form a basis for  $T_{\mathcal{Q}, \mathbb{X}}$  because the dimension coincides.

Applying the natural transformation  $h(\theta)_D$  to any of these we construct 13 first order deformations of the canonically embedded curve  $\widetilde{C}_{\mathbb{X}}$  as in Section 2.3, which will give rise to as many rows of the matrix  $M_{\mathcal{F}}$  described above.

It is now enough to check if the rank of this matrix is maximal for a particular example, which may as well be the test curve described in Section 2.4. Indeed we performed the required computations on the test surface  $\mathbb{X}$ , and obtained a positive answer: the rank is maximal. In Section 4.4 we describe the details and explain the computer code used with *Macaulay 2*.  $\square$

In the 1980s Donagi [12] proved the unirationality of  $\mathcal{R}_6$  at about the same time as Catanese [5] proved that  $\mathcal{R}_4$  is in fact rational, but since then there has been a gap between four and six, which we have now filled with the above theorem. Besides, as we have seen at the end of Chapter 1 we also obtain as a consequence a new proof of a theorem of Clemens [8]

**Corollary 3.15.**  $\mathcal{A}_4$  is unirational.

This result follows in fact from the previous theorem because the Prym map  $p_5: \mathcal{R}_5 \rightarrow \mathcal{A}_4$  is generically surjective [see Beauville, 4]. The original proof of Clemens differs from the present in that it relied upon intermediate Jacobians.

# Chapter 4

## Computations

The last part of this thesis is devoted to the computations. Having already made clear in Section 2.4 the reasons why the results obtained here are relevant, in what follows we just describe how we were able to construct an explicit example, and to test it for all the properties needed. The computer algebra package which the code refers to is *Macaulay 2* [21], so in principle one should be able to copy it as it stands and obtain at least similar results; if the online documentation doesn't help there is also the book edited by Eisenbud, Grayson and Stillman [16]. When describing the code, input data will be preceded by `i`: while output data will be preceded by `o`:, for no particular reason at all the ground field is always  $\mathbb{F}_{101}$ .

### 4.1 The Standard Test Example

We fix the coordinate ring of  $\mathbb{P}^3$  and the six points  $P_0, \dots, P_5$ , which will be the double points of the surface.

```
i: R = ZZ/101[x_0..x_3];
i: P0 = ideal(x_0,x_1,x_2);
i: P1 = ideal(x_0,x_1,x_3);
i: P2 = ideal(x_0,x_2,x_3);
i: P3 = ideal(x_1,x_2,x_3);
i: P4 = ideal(x_0-x_1,x_0-x_2,x_0-x_3);
i: P5 = ideal(x_1-2*x_0,x_2-3*x_0,x_3-4*x_0);
```

The surface  $\mathbb{X}$  will be defined by a single irreducible homogeneous polynomial  $F$  of degree four contained in the intersection  $I = \bigcap_{i=0}^5 p_i^2$ , and as such it can be randomly selected. More precisely we ask *Macaulay 2* to compute a system of generators for the ideal  $I$ , which will be returned in the form of a matrix as follows.

```
i: I = intersect(P0^2,P1^2,P2^2,P3^2,P4^2,P5^2);
i: polys = gens I;
      1      11
o: Matrix R <--- R
```

Then we instruct the program to compute a random matrix of homogeneous polynomials in  $R$  such that when multiplied with the above the result is a form of degree four.

```
i: MM1 = random(source polys, R^{-4})
      11      1
o: Matrix R <--- R
i: F1mat = polys * MM1;
      1      1
o: Matrix R <--- R
```

Observe that  $F1mat$  is in fact a one-by-one matrix, so that the equation we are after will be the determinant  $F1 = \det F1mat$ . The result must now be tested, firstly to ensure its singular locus amounts to just the six double points and then to ensure it verifies the generic condition described in Proposition 2.4.

```
i: D1 = diff F1mat;
i: F1sing = ideal(D1);
i: hilbertPolynomial F1sing
o = 6*P
      0
o: ProjectiveHilbertPolynomial
```

The Hilbert polynomial ensures us that the singular locus consists of just the six points we fixed. Next we turn to the condition on the cones over  $P_0$ . The equation  $F1$  will be of the form  $u_2x_3^2 + 2u_3x_3 + u_4$  and we want to compute the forms  $u_i$  first. The command `coefficients` will do, but we must remember

that it returns a list of two lists, one containing the powers of  $x_3$  present in the expression and the other the actual coefficients.

```
i: CoeffListF1 = coefficients({3},F1mat);
i: CoeffF1 = CoeffListF1_{1};
i: F1cones = ideal(CoeffF1);
i: hilbertPolynomial F1cones
o = 24*P
      0
o: ProjectiveHilbertPolynomial
```

Again the Hilbert polynomial assures us that the condition is satisfied, because saying that this intersection consists of only a finite number of points is equivalent to saying it is given by  $P_0$  with some multiplicity. Indeed  $u_i$  is a cone over  $P_0$  for each  $i$ , so the intersection contains  $P_0$  only or otherwise it contains a line. Now we can compute the equation of the plane curve  $C_X$  as follows.

```
i: F1curves = gens F1cones;
i: G4mat = F1curves*matrix{{1},{0},{0}};
i: G4 = det G4mat
i: G3mat = F1curves*matrix{{0},{1},{0}}
i: G3 = 1/2*(det G3mat)
i: G2mat = F1curves*matrix{{0},{0},{1}}
i: G2 = det G2mat
i: F1 = det F1mat
i: F1 == G2*x_3^2+2*G3*x_3+G4
o = true
i: CX = G3^2-G2*G4;
```

In translating the instructions given to *Macaulay 2* into our notation the polynomials  $G_2$ ,  $G_3$  and  $G_4$  will become  $u_2$ ,  $u_3$  and  $u_4$ . Thus the three homogeneous forms above, which passed all the tests in *Macaulay 2*, were the following

$$\begin{aligned}
 u_2 &= 19x_0^2 - 33x_0x_1 + 50x_1^2 - 13x_0x_2 + 50x_1x_2 - 15x_2^2 \\
 u_3 &= -2x_0^2x_1 - 35x_0x_1^2 - 18x_0^2x_2 - 8x_0x_1x_2 - 36x_1^2x_2 - 4x_0x_2^2 + 45x_1x_2^2 \\
 u_4 &= -38x_0^2x_1^2 - 32x_0^2x_1x_2 - 32x_0x_1^2x_2 - 6x_0^2x_2^2 - 38x_0x_1x_2^2 + 2x_1^2x_2^2
 \end{aligned}$$

Now we are left with a plane curve, which we know has five ordinary nodes, and we want to compute its normalisation. In order to do so we must blow up the plane in five points, and for this we follow Griffiths and Harris [22, §4.4] and consider the linear system of cubic curves passing through the points.

```
i: R = ZZ/101[x_0,x_1,x_2]
i: P1 = ideal(x_0,x_1);
i: P2 = ideal(x_0,x_2);
i: P3 = ideal(x_1,x_2);
i: P4 = ideal(x_0-x_1,x_0-x_2);
i: P5 = ideal(x_1-2*x_0,x_2-3*x_0);
i: canC = basis(3,intersect(P1,P2,P3,P4,P5));
i: canC = canC ** R;
i: canC = super canC;
           1      5
o: Matrix R <--- R
```

Quoting from the online documentation of *Macaulay 2*: “Some explanation regarding the `basis` command is needed here. `canC` is a matrix whose target is the ideal of the intersection of these five points, and whose source is a free module over the coefficient ring. For our purposes, there are two problems with this. The first is that we want a map where both the source and target have the base ring  $R$ . This can be accomplished by tensoring with  $R$ . The second problem is that the image of a basis element is not obviously in the ideal: it is represented in terms of the generators of  $I$ . This can be alleviated by applying `super`: this takes a homomorphism  $f: M \rightarrow N$ , where  $N$  is a submodule of a quotient module  $F/I$ , and returns the homomorphism  $f: M \rightarrow F/I$ .” Now we construct the embedding of the curve  $CX$  in  $\mathbb{P}^4$  as follows.

```
i: S = (coefficientRing R)[y_0..y_4]
i: TC = R/ideal CX
i: ff = map(TC,S,substitute(canC, TC));
i: cancurve = mingens ker ff;
           1      3
o: Matrix S <--- S
```

The kernel of `ff` is the ideal of the curve  $\widetilde{C}_X$ , which we expect from Petri's Theorem (3.11) to be generated by three quadrics. Indeed the three generators we have found are quadrics.

```
i: Q1 = cancurve * matrix{{1},{0},{0}};
```

```
i: det Q1
```

2

```
o = y y + y - y y - 34y y + 33y y
     1 2    2    0 3    2 3    2 4
```

```
i: Q2 = cancurve * matrix{{0},{1},{0}};
```

```
i: det Q2
```

2

```
o = y y - 2y - 33y y - y y + 35y y
     0 2    2    2 3    1 4    2 4
```

```
i: Q3 = cancurve * matrix{{0},{0},{1}};
```

```
i: det Q3
```

2

2

2

```
o = y - 48y y + 5y + ... + 17y y - 19y
     0    0 1    1    3 4    4
```

In order to check that the double cover of  $\widetilde{C}_X$  constructed in Proposition 2.8 is not trivial we must show that the surface  $\Sigma$  is irreducible. This amounts to proving that the following ideal in  $\mathbb{K}[x, y][\zeta_0, \zeta_1, \zeta_2]$ , homogeneous in  $\zeta_0, \zeta_1, \zeta_2$ ,

$$(-\zeta_2^2 + \zeta_1^2 G_2 + 2\zeta_1 \zeta_0 G_3 + \zeta_0^2 G_4, G_3^2 - G_2 G_4)$$

is prime. But in fact we are working here over the open subset  $D_h(x_i)$  of  $\mathbb{P}^2$ , and  $G_d$  is the polynomial obtained from  $u_d$  by multiplication by  $1/x_i^d$ , so that actually we are to check that three (very similar) ideals are prime. First we compute the equations of  $u_d$  ( $d=2,3,4$ ) over the open affine subset  $D_h(x_i)$  by instructing for instance `G20 = substitute(G2, {x_0=>1})`, and similarly for `G30` and `G40` with the obvious association of  $u_2$  over  $D_h(x_0)$  with `G20`. Then we define the ideal to test

```
i: R = ZZ/101[x_1, x_2, z_0..z_2];
```

```
i: G20 = 50*x_1^2+50*x_1*x_2-15*x_2^2-33*x_1-13*x_2+19
```

```
i: G30 = -36*x_1^2*x_2+ ... -18*x_2
```

```
i: G40 = 2*x_1^2*x_2^2- ... -6*x_2^2
i: I = ideal(G30^2+G20*G40,
            -z_2^2+z_1^2*G20+2*z_1*z_0*G30+z_0^2*G40);
```

However we are not going to check primality directly, but instead we work inside the 5-dimensional affine space and cut the threefold with a codimension two linear space  $H$ . This is enough because if the threefold was reducible, say given by the union  $Z_1 \cup Z_2$ , when cutting with  $H$  one would see the union of  $(Z_1 \cap H)$  and  $(Z_2 \cap H)$ . So if what we get is irreducible one of those must be empty. But if  $Z_2 \cap H$  is empty,  $Z_2$  must be.

```
i4 : I = ideal(G30^2+G20*G40,
              -z_2^2+z_1^2*G20+2*z_1*z_0*G30+z_0^2*G40,
              z_1-1, z_2-1);
i5 : decompose I
o6 : List
i7 : toString o6
o7 = {ideal(z_2-1, z_1-1,
           x_1^4*x_2^2- ... +22*x_2^2,
           x_1^2*x_2^2*z_0^2- ... +9,
           x_1^4*x_2*z_0^2+ ... -19,
           -6*x_1^5*x_2*z_0+ ... +25,
           x_1*x_2^5*z_0^2+ ... -37,
           x_1^3*x_2^4*z_0- ... +29,
           x_2^6*z_0^3+ ... -50)}
```

The function `decompose` returns the list of the irreducible components of an ideal, thus the output is showing that  $I$  is irreducible. Clearly we are documenting here only one of three necessary computations, but the computations on the other two affine pieces are identical.

## 4.2 The dimension of the family of surfaces

In this section we deal with the family of quartic hypersurfaces we described in Proposition 3.5. We want to compute the differential of the projection from

$\mathbb{P}(I_4) \times \mathbb{P}^3$  to  $\mathbb{P}(I_4)$ , restricted to the subvariety  $B_0$  of the product given by the equations

$$\{(F, \mathfrak{p}) \mid F \in \mathfrak{p}^2\} = \left\{ (F, P) \mid F(P) = \frac{\partial F}{\partial x_i}(P) = 0 \right\}$$

We first need to compute a basis for the degree four part of the ideal, and this is achieved as follows.

```
i: R = ZZ/101[x_0..x_3];
i: P0 = ideal(x_0,x_1,x_2);
i: ...
i: P4 = ideal(x_0-x_1,x_0-x_2,x_0-x_3);
i: Surfs = basis(4,intersect(P0^2,P1^2,P2^2,P3^2,P4^2));
i: Surfs = Surfs ** R;
i: Surfs = super Surfs;
           1      15
o: Matrix R <---- R
```

And we also have to write our central fibre over this basis, which is given by typing simply `F1 // Surfs`.

```
o: matrix {{-33}, {17}, {19}, {-16}, {13}, {-36}, {-4}, {2},
           {-38}, {-46}, {-6}, {-32}, {-1}, {-32}, {-38}};
```

Now we wish to work over an affine open subset, which means that we shall have fourteen coordinates coming from the above basis and three coming from projective space. The following code produces the equations we are interested in over the relevant open set.

```
i: A = ZZ/101[t_1..t_14];
i: R = A[x_0..x_3];
i: MatSurf = Surfs * matrix{{-33},{t_1}..{t_14}};
i: MatSurf = substitute(MatSurf,{x_0=>1})
```

Note that we are choosing an open subset where  $x_0 \neq 0$ , therefore we now pass to a polynomial ring in three variables  $x_1, x_2, x_3$ . Besides we construct a matrix containing all the equations we need, four in total.

```
i: A = ZZ/101[t_1..t_14];
```

```

i: R = A[x_1..x_3];
i: MatSurf = as above
i: MatDiff = diff MatSurf;
i: MatFam = MatSurf|MatDiff;

```

Now we operate over an open affine subset, and again we need to change the ideal for the number of variables to be correct. We compute this time the Jacobian matrix of the four equations, or in other words the tangent space of  $B_0$  (embedded in  $\mathbb{A}^{17}$ ) at the point corresponding to our surface  $\mathbb{X}$ .

```

i: A = ZZ/101[t_1..t_14,x_1..x_3];
i: MatFam = as above
i: FamDiff = jacobian MatFam
i: FamJac = substitute(FamDiff,{t_1=>17, t_2=>19,
    t_3=>-16, t_4=>13, t_5=>-36, t_6=>-4, t_7=>2,
    t_8=>-38, t_9=>-46, t_10=>-6, t_11=>-32, t_12=>-1,
    t_13=>-32, t_14=>-38, x_1=>2, x_2=>3, x_3=>4})

```

Finally we compute the intersection of the ideal of definition of the tangent space with the subalgebra given by the first fourteen variables.

```

i: R = ZZ/101[t_1..t_17];
i: FamJac = as above
i: TgSpId = FamJac * matrix{{t_1}..{t_17}};
           4      1
o: Matrix R <--- R
i: TSI = ideal TgSpId;
i: MatPrj = matrix{{t_1..t_14}};
i: S = (coefficientRing R)[y_1..y_14];
i: TC = R/TSI;
i: ff = map(TC,S,substitute(MatPrj, TC));
o: RingMap TC <--- S
i: mingens ker ff
           1      1
o: Matrix S <--- S

```

### 4.3 The Family of Plane Curves

Starting from the family of quartic hypersurfaces we described in Proposition 3.5 we compute here the infinitesimal deformations of the standard test example  $\mathbb{X}$  and the associated family of deformations of sextic plane curves. We wish to work on an open affine subset of  $B_0$  centred at  $\mathbb{X}$ , so we first fix coordinates  $a, b, c, t_1 \dots t_{14}$ . We use the basis `surfs` for quartic hypersurfaces with nodes at five fixed points that we computed above.

```
i: A = ZZ/101[a,b,c,t_1..t_14];
i: R = A[x_0..x_3];
i: Surfs = as above
i: MatSurf = matrix {{-33}, {17+t_1}, {19+t_2},
                    {-16+t_3}, {13+t_4}, {-36+t_5}, {-4+t_6},
                    {2+t_7}, {-38+t_8}, {-46+t_9}, {-6+t_10},
                    {-32+t_11}, {-1+t_12}, {-32+t_13}, {-38+t_14}}
i: Surface = det(Surfs*MatSurf);
i: DffSurf = diff Surface;
i: Eqtn1 = substitute(Surface,
                    {x_0=>1, x_1=>2+a, x_2=>3+b, x_3=>4+c});
i: Eqtn2 = substitute(DffSurf,
                    {x_0=>1, x_1=>2+a, x_2=>3+b, x_3=>4+c});
```

In order to define coordinates over  $B_0$  we need to solve the linear system in  $t_1 \dots t_{14}$  with respect to  $a, b, c$ ; we can do it by computing a Gröbner basis for the ideal of the equations (function `gb`).

```
i1 : A = ZZ/101[a,b,c];
i2 : K = frac A;
i3 : R = K[t_1..t_14];
i4 : Eqtn1 = as above
i5 : Eqtn2 = as above
i6 : EqSys = ideal(Eqtn1)+ideal(Eqtn2);
i7 : Eqns = generators gb EqSys;
      1      4
o7 : Matrix R <--- R
```

At this point some adjustments are needed. The solution of the system allows us to replace four variables with polynomials, and we compute these four. In this case the variables happen to be  $t_5, t_3, t_2, t_1$  so we name the polynomials  $G_5, G_3, G_2, G_1$ . After this operation we will be left with an asymmetric set of variables, so we rename them.

```
i3 : R = K[t_1..t_14,s_1..s_10];
i4 : G1 = as above
i5 : H1 = substitute(G1, {t_4=>s_1, t_6=>s_2,
      t_7=>s_3, t_8=>s_4, t_9=>s_5, t_10=>s_6,
      t_11=>s_7, t_12=>s_8, t_13=>s_9, t_14=>s_10})
```

Now we substitute the solution of the linear system inside the expression of the family.

```
i1 : A = ZZ/101[a,b,c];
i2 : K = frac A;
i3 : B = K[s_1..s_10];
i4 : R = B[x_0..x_3];
i5 : H1 = as above
i6 : H2 = as above
i7 : H3 = as above
i8 : H5 = as above
i9 : Surfs = as above
i10 : MatSurf = matrix {{-33}, {17+H1}, {19+H2},
      {-16+H3}, {13+s_1}, {-36+H5}, {-4+s_2},
      {2+s_3}, {-38+s_4}, {-46+s_5}, {-6+s_6},
      {-32+s_7}, {-1+s_8}, {-32+s_9}, {-38+s_10}};
i11 : Surface = det(Surfs*MatSurf);
```

The coefficients of the quartic polynomial surface are rational functions on  $a, b, c$ . In order to deal with an infinitesimal family we then clear denominators.

```
i1 : A = ZZ/101[a,b,c];
i2 : K = frac A;
i3 : B = K[s_1..s_10];
i4 : R = B[x_0..x_3];
```

```

i5 : Surface = as above
i6 : Surface = (a^3*b*c- ... -12*a+4*b-5*c-4)*Surface
i7 : Surface = (b^2+5*b+6)*Surface
i8 : Surface = (c+4)*Surface
i9 : Surface = (a-b-1)*Surface
i10 : Surface = 38/12*Surface

```

Finally we can read the equation infinitesimally around the point  $\mathbb{X}$ , just by taking the quotient for the maximal ideal squared.

```

i1 : A = ZZ/101[a,b,c,s_1..s_10];
i2 : JJ = ideal(a,b,c,s_1..s_10);
i3 : B = A/JJ^2
i4 : R = B[x_0..x_3];
i5 : Surface = as above

```

We are now ready to compute the equation for the family of curves, but again *Macaulay 2* doesn't allow us to work directly over the infinitesimal ring. So first we make the computation over the fraction field and then we clear the terms of higher order with a quotient.

```

i1 : A = ZZ/101[a,b,c,s_1..s_10];
i2 : K = frac A;
i3 : R = K[x_0..x_3];
i4 : surface = as above
i5 : CoeffListF1 = coefficients({3},surface)
i6 : CoeffF1 = CoeffListF1_{1}
i7 : F1cones = ideal(CoeffF1)
i8 : F1curves = gens F1cones
i9 : G4mat = F1curves*matrix{{1},{0},{0}}
i10 : G4 = det G4mat
i12 : G3mat = F1curves*matrix{{0},{1},{0}}
i13 : G3 = 1/2*(det G3mat)
i15 : G2mat = F1curves*matrix{{0},{0},{1}}
i16 : G2 = det G2mat
i18 : surface == G2*x_3^2+2*G3*x_3+G4
o18 = true

```

```
i19 : curves = G3^2-G2*G4
```

In what follows all *Macaulay 2* does is read the polynomial we input, which we take from above, and then write it back leaving out all the terms of higher order.

```
i1 : A = ZZ/101[a,b,c,s_1..s_10];
i2 : JJ = ideal(a,b,c,s_1..s_10);
i3 : B = A/JJ^2
i4 : R = B[x_0..x_2];
i5 : curves =
```

## 4.4 The Rank of a big Matrix

With reference to Section 3.3, and in particular to Theorem 3.14, we want to compute the matrix  $M_{\mathcal{F}}$  where  $\mathcal{F}$  is the family of quartic surfaces we described in Section 4.2 or more precisely for our current purposes the family curves of sextic plane curves defined in the previous section. We start by computing the part of  $M_{\mathcal{F}}$  which doesn't depend on the family, but only on the central fibre. So, looking at the action of  $\mathfrak{gl}(3)$  we need to write the coefficients of the three quadratic forms already encountered in Section 4.1.

```
i: R = ZZ/101[y_0..y_4];
i: F1 = y_1*y_2+y_2^2-y_0*y_3-34*y_2*y_3+33*y_2*y_4;
i: F2 = y_0*y_2-2*y_2^2-33*y_2*y_3-y_1*y_4+35*y_2*y_4;
i: F3 = y_0^2-48*y_0*y_1+ ... +17*y_3*y_4-19*y_4^2;
i: MM = basis(2,R)
i7 : MM1 = transpose(F1//MM)
i9 : MM2 = transpose(F2//MM)
i10 : MM3 = transpose(F3//MM)
```

Given the vectors  $MM1$ ,  $MM2$  and  $MM3$  one can write down the corresponding  $9 \times 45$  matrix by hand, which is exactly what we did. Let it be  $MM9$ . As for the action of  $\mathfrak{sl}(5)$  we start by fixing a basis for the vector space of traceless  $5 \times 5$  matrices, denoting  $D24$  the matrix with a unique nonzero entry (which will be a 1) in position  $_{24}$  and by  $DD3$  the diagonal matrix  $\text{diag}(1, 0, 0, -1, 0)$ . Then

denoting by  $Q2$  the symmetric matrix associated to the polynomial  $F2$  above we do the following.

```

i: R = QQ[y_0..y_4];
i: MM = basis(2,R)
i: X = matrix{{y_0..y_4}}
i: Y = transpose X
i: F01P1 = det (X*(Q1*D01+D10*Q1)*Y)
i: F01P2 = det (X*(Q2*D01+D10*Q2)*Y)
i: F01P3 = det (X*(Q3*D01+D10*Q3)*Y)
i: C01P1 = transpose (F01P1 // MM)
i: C01P2 = transpose (F01P2 // MM)
i: C01P3 = transpose (F01P3 // MM)
i: C01 = C01P1|C01P2|C01P3
i: F02P1 = det (X*(Q1*D02+D20*Q1)*Y)
i: F02P2 = det (X*(Q2*D02+D20*Q2)*Y)
i: F02P3 = det (X*(Q3*D02+D20*Q3)*Y)
i: C02P1 = transpose (F02P1 // MM)
i: C02P2 = transpose (F02P2 // MM)
i: C02P3 = transpose (F02P3 // MM)
i: C02 = C02P1|C02P2|C02P3
i: slfive = C01||C02
i: ...
i: F24P1 = det (X*(Q1*DD4+DD4*Q1)*Y)
i: F24P2 = det (X*(Q2*DD4+DD4*Q2)*Y)
i: F24P3 = det (X*(Q3*DD4+DD4*Q3)*Y)
i: C24P1 = transpose (F24P1 // MM)
i: C24P2 = transpose (F24P2 // MM)
i: C24P3 = transpose (F24P3 // MM)
i: C24 = C24P1|C24P2|C24P3
i: slfive = slfive||C24
i: MatFam = slfive||MM9
          33      45
o: Matrix R  <--- R
i: rank MatFam

```

```
o = 33
```

For each first order deformation of the family curves mentioned above, we now need to compute the normalisation. But first we need to extract each equation from the equation of the family.

```
i: A = ZZ/101[a,b,c,s_1..s_10];
i: R = A[x_0..x_2];
i: curves = as above
i: C01 = substitute(curves,{a=>0, b=>0, c=>0, s_2=>0,
      s_3=>0, s_4=>0, s_5=>0, s_6=>0, s_7=>0,
      s_8=>0, s_9=>0, s_10=>0})
i: ...
i: C10 = substitute(curves,{a=>0, b=>0, c=>0, ..., s_9=>0})
i: C11 = substitute(curves,{a=>0, b=>0, s_1=>0, ..., s_10=>0})
i: C12 = substitute(curves,{a=>0, c=>0, s_1=>0, ..., s_10=>0})
i: C13 = substitute(curves,{b=>0, c=>0, s_1=>0, ..., s_10=>0})
```

In Section 4.1 we have already encountered the algorithm for the computation of the normalisation of a nodal plane curve of this type. Observe that it relies on a basis of plane cubics through the five points we want to blow up. It is clear then that the set of deformations above consists of two disjoint parts: those deformations with nodes at five fixed points, and those with a non-fixed node. The former is the biggest, containing eleven deformations out of thirteen, indeed from the geometry of the problem one sees immediately that it contains also the parameter  $c$ . Thus we distinguish two cases, and start from the most common one. We denote the parameter for each 1-dimensional family we consider by  $e$ , meaning successively  $e=c, e=s_1, \dots, e=s_{10}$

```
i: A = ZZ/101[e]/ideal e^2;
i: K = frac A
i: R = K[x_0,x_1,x_2];
i: C = this is the deformation we are considering
```

The following set of commands computes the pull-back of the family above, in other words it computes the corresponding deformation of canonical curves.

```
i: canC = as in Section 4.1
```

```

i: S = (coefficientRing R)[y_0..y_4];
i: TC = R/ideal C;
i: ff = map(TC,S,substitute(canC, TC));
i: cancurve = mingens ker ff
i: F1 = det( cancurve * matrix{{1},{0},{0}} )
i: F2 = det( cancurve * matrix{{0},{1},{0}} )
i: F3 = det( cancurve * matrix{{0},{0},{1}} )

```

Now we need to normalise the expressions, but *Macaulay 2* doesn't allow us to do it over this ring. We clear denominators and check that the central fiber coincides with our chosen one.

```

i: A = ZZ/101[e];
i: K = frac A
i: R = K[y_0..y_4];
i: Q1 = y_1*y_2+y_2^2-y_0*y_3-34*y_2*y_3+33*y_2*y_4
i: Q2 = y_0*y_2-2*y_2^2-33*y_2*y_3-y_1*y_4+35*y_2*y_4
i: Q3 = y_0^2-48*y_0*y_1+ ... +17*y_3*y_4-19*y_4^2
i: T1 = F1-Q1
i: T2 = F2-Q2
i: T3 = F3-Q3
i: MM = basis(2,R)
i: RP1 = transpose (T1 // MM)
i: RP2 = transpose (T2 // MM)
i: RP3 = transpose (T3 // MM)
i: Row01 = RP1|RP2|RP3

```

Lastly we attach the row just computed to the matrix we already have.

```

i: MatFam = MatFam||Row01

```

In the case of the parameters  $a, b$  the difference is of course in how we compute the normalisation. So we leave the rest untouched and go back to that computation. The following set of commands computes the pull-back of the family above, in other words it computes the corresponding deformation of canonical curves.

```

i: P1 = ideal(x_0,x_1);

```

```

i: P2 = ideal(x_0,x_2);
i: P3 = ideal(x_1,x_2);
i: P4 = ideal(x_0-x_1,x_0-x_2);
i: P5 = ideal(x_1-(2+a)*x_0,x_2-3*x_0);

```

or

```

i: P5 = ideal(x_1-2*x_0,x_2-(3+b)*x_0);
i: canC = basis(3,intersect(P1,P2,P3,P4,P5));
i: canC = canC ** R;
i: canC = super canC;
i: C = this is the deformation we are considering
i: S = (coefficientRing R)[y_0..y_4];
i: TC = R/ideal C;
i: ff = map(TC,S,substitute(canC, TC))
i: cancurve = mingens ker ff
i: F1 = det( cancurve * matrix{{1},{0},{0}} )
i: F2 = det( cancurve * matrix{{0},{1},{0}} )
i: F3 = det( cancurve * matrix{{0},{0},{1}} )

```

After this lengthy set of computations we were left with a  $46 \times 45$  matrix, whose entries are actually integers and whose rank we want to be maximal for our result to hold. The name we actually gave to this matrix for *Macaulay 2* to handle at this stage was an emblematic hope. We conclude this thesis with the answer we received, as it appeared on the screen one day, leaving the reader to imagine what happened next around that same screen...

```

          46      45
o28 : Matrix R  <--- R
i29 : rank hope
o29 = 45

```

# Bibliography

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths and J. Harris, *Geometry of Algebraic Curves. Vol. I*, *Grundlehren der Mathematischen Wissenschaften*, vol. 267, Springer-Verlag, New York, 1985.
- [2] E. Arbarello and E. Sernesi, *The equation of a plane curve*, *Duke Math. J.* **46** (1979) 469–485.
- [3] A. Beauville, *Prym varieties and the Schottky problem*, *Invent. Math.* **41** (1977) 149–196.
- [4] A. Beauville, *Prym varieties: a survey*, in *Theta functions—Bowdoin 1987*, Part 1 (Brunswick, ME, 1987), *Proc. Sympos. Pure Math.*, vol. 49, Amer. Math. Soc., Providence, RI, 1989, 607–620.
- [5] F. Catanese, *On the rationality of certain moduli spaces related to curves of genus 4*, in *Algebraic geometry (Ann Arbor, Mich., 1981)*, *Lecture Notes in Math.*, vol. 1008, Springer, Berlin, 1983, 30–50.
- [6] M. C. Chang and Z. Ran, *Unirationality of the moduli spaces of curves of genus 11, 13 (and 12)*, *Invent. Math.* **76** (1984) 41–54.
- [7] H. Clemens, *A scrapbook of Complex Curve Theory*, Plenum Press, New York, 1980. The University Series in Mathematics.
- [8] H. Clemens, *Double solids*, *Adv. in Math.* **47** (1983) 107–230.
- [9] H. Clemens, *The quartic double solid revisited*, in *Complex geometry and Lie theory (Sundance, UT, 1989)*, *Proc. Sympos. Pure Math.*, vol. 53, Amer. Math. Soc., Providence, RI, 1991, 89–101.

- [10] O. Debarre, *Tores et Variétés Abéliennes Complexes*, *Cours Spécialisés*, vol. 6, Société Mathématique de France, Paris, 1999.
- [11] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*, *Inst. Hautes Études Sci. Publ. Math.* (1969) 75–109.
- [12] R. Donagi, *The unirationality of  $\mathcal{A}_5$* , *Ann. of Math. (2)* **119** (1984) 269–307.
- [13] R. Donagi, *The fibers of the Prym map*, in *Curves, Jacobians, and abelian varieties* (Amherst, MA, 1990), *Contemp. Math.*, vol. 136, Amer. Math. Soc., Providence, RI, 1992, 55–125.
- [14] R. Donagi and R. C. Smith, *The structure of the Prym map*, *Acta Math.* **146** (1981) 25–102.
- [15] D. Eisenbud, *Commutative Algebra with a view toward Algebraic Geometry*, *Graduate Texts in Mathematics*, vol. 150, Springer-Verlag, New York, 1995.
- [16] D. Eisenbud, D. R. Grayson and M. Stillman, eds., *Computations in algebraic geometry with Macaulay 2*, *Algorithms and Computation in Mathematics*, vol. 8, Springer-Verlag, Berlin, 2002.
- [17] D. Eisenbud and J. Harris, *The Kodaira dimension of the moduli space of curves of genus  $\geq 23$* , *Invent. Math.* **90** (1987) 359–387.
- [18] D. Eisenbud and J. Harris, *The Geometry of Schemes*, *Graduate Texts in Mathematics*, vol. 197, Springer-Verlag, New York, 2000.
- [19] E. Freitag, *Siegelsche Modulfunktionen*, *Grundlehren der Mathematischen Wissenschaften*, vol. 254, Springer-Verlag, Berlin, 1983.
- [20] W. Fulton and J. Harris, *Representation Theory*, *Graduate Texts in Mathematics*, vol. 129, Springer-Verlag, New York, 1991.
- [21] D. R. Grayson and M. E. Stillman, *Macaulay 2, a software system for research in algebraic geometry* [online], version 0.9.2, available from <http://www.math.uiuc.edu/Macaulay2/> [Accessed 20 July 2005], Windows users may visit <http://www.comalg.org/m2win.shtml> [Accessed 20 July 2005], 2001.

- [22] P. Griffiths and J. Harris, *Principles of Algebraic Geometry, reprint of the 1978 original*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1994.
- [23] M. Grooten, Ordinary Double Solids [online], Master's thesis, Mathematics Department, University of Nijmegen, the Netherlands, available from <http://www.math.ru.nl/~grooten/maths.php> [Accessed 29 November 2005], 2001.
- [24] A. Grothendieck, *Techniques de construction et théorèmes d'existence en géométrie algébrique. IV. Les schémas de Hilbert*, in Séminaire Bourbaki, Vol. 6, Soc. Math. France, Paris, 1995, Exp. No. 221, 249–276.
- [25] S. Grushevsky and D. Lehavi,  $\mathcal{A}_6$  is of general type, math.AG/0512530 (preprint), 2005.
- [26] J. Harris and I. Morrison, *Moduli of Curves, Graduate Texts in Mathematics*, vol. 187, Springer-Verlag, New York, 1998.
- [27] J. Harris and D. Mumford, *On the Kodaira dimension of the moduli space of curves*, *Invent. Math.* **67** (1982) 23–88. With an appendix by William Fulton.
- [28] R. Hartshorne, *Algebraic Geometry, Graduate Texts in Mathematics*, vol. 52, Springer-Verlag, New York, 1977.
- [29] J. Kollár, K. E. Smith and A. Corti, *Rational and nearly rational varieties, Cambridge Studies in Advanced Mathematics*, vol. 92, Cambridge University Press, Cambridge, 2004.
- [30] B. Kreussler, *Small resolutions of double solids, branched over a 13-nodal quartic surface*, *Ann. Global Anal. Geom.* **7** (1989) 227–267.
- [31] B. Kreussler, *Another description of certain quartic double solids*, *Math. Nachr.* **212** (2000) 91–100.
- [32] S. Lang, *Abelian Varieties, reprint of the 1959 original*, Springer-Verlag, New York, 1983.

- [33] Q. Liu, *Algebraic Geometry and Arithmetic Curves*, *Oxford Graduate Texts in Mathematics*, vol. 6, Oxford Univ. Press, Oxford, 2002.
- [34] S. Mac Lane, *Categories for the Working Mathematician*, *Graduate Texts in Mathematics*, vol. 5, Springer-Verlag, New York, 2nd ed., 1998.
- [35] L. Masiewicki, *Universal properties of Prym varieties with an application to algebraic curves of genus five*, *Trans. Amer. Math. Soc.* **222** (1976) 221–240.
- [36] J. S. Milne, *Abelian Varieties* [online], available from <http://www.jmilne.org/math/> [Accessed 13 October 2005], under the Creative Commons License <http://creativecommons.org/licenses/by-nc-nd/2.0/>, 1998.
- [37] J. S. Milne, *Algebraic Geometry* [online], available from <http://www.jmilne.org/math/> [Accessed 6 July 2005], under the Creative Commons License <http://creativecommons.org/licenses/by-nc-nd/2.0/>, 2005.
- [38] S. Mori and S. Mukai, *The uniruledness of the moduli space of curves of genus 11*, in *Algebraic geometry (Tokyo/Kyoto, 1982)*, *Lecture Notes in Math.*, vol. 1016, Springer, Berlin, 1983, 334–353.
- [39] D. Mumford, *Abelian Varieties*, *Tata Institute of Fundamental Research Studies in Mathematics*, vol. 5, Oxford Univ. Press, published for the Tata Institute of Fundamental Research, Bombay, 1970.
- [40] D. Mumford, *Prym varieties. I*, in *Contributions to analysis (a collection of papers dedicated to Lipman Bers)*, Academic Press, New York, 1974, 325–350.
- [41] D. Mumford, *On the Kodaira dimension of the Siegel modular variety*, in *Algebraic geometry—open problems (Ravello, 1982)*, *Lecture Notes in Math.*, vol. 997, Springer, Berlin, 1983, 348–375.
- [42] D. Mumford, *The Red Book of Varieties and Schemes*, *Lecture Notes in Mathematics*, vol. 1358, Springer-Verlag, Berlin, 2nd ed., 1999.
- [43] M. Reid, *Undergraduate Algebraic Geometry*, *L.M.S. Student Texts*, vol. 12, Cambridge Univ. Press, Cambridge, 1988.

- [44] M. Reid, *Undergraduate Commutative Algebra*, *L.M.S. Student Texts*, vol. 29, Cambridge Univ. Press, Cambridge, 1995.
- [45] M. Reid, *Chapters on algebraic surfaces*, in *Complex algebraic geometry* (Park City, UT, 1993), *IAS/Park City Math. Ser.*, vol. 3, Amer. Math. Soc., Providence, RI, 1997, 3–159.
- [46] M. Reid, *Graded rings and varieties in weighted projective space* [online], available from <http://www.maths.warwick.ac.uk/~miles/surf/> [Accessed 6 July 2005], 2002.
- [47] B. Riemann, *Theorie der Abel'schen Functionen*, *J. Reine Angew. Math.* **54** (1857) 101–155.
- [48] E. Sernesi, *Unirationality of the variety of moduli of curves of genus twelve*, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **8** (1981) 405–439.
- [49] E. Sernesi, *Deformations of Algebraic Schemes*, *Grundlehren der Mathematischen Wissenschaften*, vol. 334, Springer-Verlag, Berlin, 2006.
- [50] I. R. Shafarevich, *Basic Algebraic Geometry 1: Varieties in Projective Space*, Springer-Verlag, Berlin, 2nd ed., 1994.
- [51] I. R. Shafarevich, *Basic Algebraic Geometry 2: Schemes and Complex Manifolds*, Springer-Verlag, Berlin, 2nd ed., 1994.
- [52] Y.-S. Tai, *On the Kodaira dimension of the moduli space of abelian varieties*, *Invent. Math.* **68** (1982) 425–439.
- [53] B. R. Tennison, *Sheaf Theory*, *London Math. Soc. Lecture Note Series*, vol. 20, Cambridge Univ. Press, Cambridge, 1975.
- [54] A. Verra, *A short proof of the unirationality of  $\mathcal{A}_5$* , *Nederl. Akad. Wetensch. Indag. Math.* **46** (1984) 339–355.
- [55] O. Zariski and P. Samuel, *Commutative Algebra. Vol. I*, The University Series in Higher Mathematics, D. Van Nostrand Company, Inc., Princeton, New Jersey, 1958.