

Spines in growth-fragmentation models


Alex Watson
University of Manchester

Paris–Bath IV
28 June 2016


A finite-activity model




A finite-activity model




A finite-activity model



A finite-activity model




A finite-activity model



- \mathbf{p} is picked from

$$\mathcal{P} = \{(p_1, p_2, \dots) : p_1 \geq p_2 \geq \dots, \sum_{i \geq 1} p_i = 1\}.$$

A finite-activity model




- \mathbf{p} is picked from

$$\mathcal{P} = \{(p_1, p_2, \dots) : p_1 \geq p_2 \geq \dots, \sum_{i \geq 1} p_i = 1\}.$$

- Mean measure:

$$\langle \mu_t, f \rangle = \int f(x) \mu_t(dx) = \mathbb{E} \left[\sum_{\substack{u \text{ particle} \\ \text{alive at } t}} f(\text{size}(u)) \right]$$

A finite-activity model



- \mathbf{p} is picked from

$$\mathcal{P} = \{(p_1, p_2, \dots) : p_1 \geq p_2 \geq \dots, \sum_{i \geq 1} p_i = 1\}.$$

- Mean measure:

$$\langle \mu_t, f \rangle = \int f(x) \mu_t(dx) = \mathbb{E} \left[\sum_{\substack{u \text{ particle} \\ \text{alive at } t}} f(\text{size}(u)) \right]$$

- $\nu(d\mathbf{p}) = \lambda K(d\mathbf{p})$

The pure fragmentation equation

$$\partial_t \langle \mu_t, f \rangle = \left\langle \mu_t, \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} f(x p_i) - f(x) \right\} \nu(d\mathbf{p}) \right\rangle,$$
$$f \in C_c^\infty(0, \infty),$$

$$\mu_0 = \delta_1$$

The growth-fragmentation equation

$$\partial_t \langle \mu_t, f \rangle = \left\langle \mu_t, \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} f(xp_i) - f(x) \right\} \nu(d\mathbf{p}) \right\rangle,$$

The growth-fragmentation equation

$$\begin{aligned} \partial_t \langle \mu_t, f \rangle &= \left\langle \mu_t, axf'(x) \right. \\ &\quad \left. + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} f(xp_i) - f(x) \right\} \nu(d\mathbf{p}) \right\rangle, \end{aligned}$$

The growth-fragmentation equation

$$\begin{aligned} \partial_t \langle \mu_t, f \rangle &= \left\langle \mu_t, axf'(x) \right. \\ &\quad \left. + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} f(xp_i) - f(x) + (1 - p_1)xf'(x) \right\} \nu(d\mathbf{p}) \right\rangle, \end{aligned}$$

The growth-fragmentation equation

$$\begin{aligned}\partial_t \langle \mu_t, f \rangle &= \left\langle \mu_t, a x f'(x) \right. \\ &\quad \left. + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} f(x p_i) - f(x) + (1 - p_1) x f'(x) \right\} \nu(d\mathbf{p}) \right\rangle,\end{aligned}$$

■ $a \in \mathbb{R}$

The growth-fragmentation equation

$$\begin{aligned} \partial_t \langle \mu_t, f \rangle &= \left\langle \mu_t, axf'(x) \right. \\ &\quad \left. + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} f(xp_i) - f(x) + (1 - p_1)xf'(x) \right\} \nu(d\mathbf{p}) \right\rangle, \end{aligned}$$

- $a \in \mathbb{R}$
- Require only $\int (1 - p_1)^2 \nu(d\mathbf{p}) < \infty$ (asymmetric children)

Questions for today

- Existence and representation
- Explore many-to-one theorem


Questions for today

- Existence and representation
- Explore many-to-one theorem

Fragmentation processes ($\nu < \infty$)

$$\mathcal{Z}(t) = \sum_{u \text{ fragments}} \delta_{\log(\text{size}(u))} \mathbb{1}_{\{u \text{ alive at time } t\}}$$

\mathcal{Z}




Fragmentation processes ($\nu < \infty$)

$$\mathcal{Z}(t) = \sum_{u \text{ fragments}} \delta_{\log(\text{size}(u))} \mathbb{1}_{\{u \text{ alive at time } t\}}$$

This is a compound Poisson process with immigration.

\mathcal{Z}




Compensated fragmentation processes, $\int (1 - p_1)^2 \nu(d\mathbf{p}) < \infty$

- We can build \mathcal{Z} in general
- Create a Lévy process whose Lévy measure is $\nu(\log p_1 \in \cdot)$
- At every jump of size z , sample from $\nu(d\mathbf{p} \mid \log p_1 = z)$ and immigrate new particles at relative positions $\log p_i$, $i \geq 2$.


Compensated fragmentation processes, $\int (1 - p_1)^2 \nu(d\mathbf{p}) < \infty$

- We can build \mathcal{Z} in general
- Create a Lévy process whose Lévy measure is $\nu(\log p_1 \in \cdot)$
- At every jump of size z , sample from $\nu(d\mathbf{p} \mid \log p_1 = z)$ and immigrate new particles at relative positions $\log p_i, i \geq 2$.




Compensated fragmentation processes, $\int (1 - p_1)^2 \nu(d\mathbf{p}) < \infty$

- We can build \mathcal{Z} in general
- Create a Lévy process whose Lévy measure is $\nu(\log p_1 \in \cdot)$
- At every jump of size z , sample from $\nu(d\mathbf{p} \mid \log p_1 = z)$ and immigrate new particles at relative positions $\log p_i, i \geq 2$.




Compensated fragmentation processes, $\int (1 - p_1)^2 \nu(d\mathbf{p}) < \infty$

- We can build \mathcal{Z} in general
- Create a Lévy process whose Lévy measure is $\nu(\log p_1 \in \cdot)$
- At every jump of size z , sample from $\nu(d\mathbf{p} \mid \log p_1 = z)$ and immigrate new particles at relative positions $\log p_i$, $i \geq 2$.



Compensated fragmentation processes, $\int (1 - p_1)^2 \nu(d\mathbf{p}) < \infty$

- We can build \mathcal{Z} in general
- Create a Lévy process whose Lévy measure is $\nu(\log p_1 \in \cdot)$
- At every jump of size z , sample from $\nu(d\mathbf{p} \mid \log p_1 = z)$ and immigrate new particles at relative positions $\log p_i$, $i \geq 2$.



(Aside: let's assume we have a consistent way to give the particles labels $u \in \mathcal{U}$. Write $\mathcal{Z}_u(t)$ for the position of particle u at time t .)

Solution of the equation

Define the **cumulant** κ : $\mathbb{E}[\sum_u e^{qZ_u(t)}] = e^{t\kappa(q)}$. It satisfies

$$\kappa(q) = aq + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} p_i^q - 1 + (1 - p_1)q \right\} \nu(d\mathbf{p}).$$

Solution of the equation

Define the cumulant κ : $\mathbb{E}[\sum_u e^{qZ_u(t)}] = e^{t\kappa(q)}$. It satisfies

$$\kappa(q) = aq + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} p_i^q - 1 + (1 - p_1)q \right\} \nu(d\mathbf{p}).$$

Theorem (Bertoin, W. '15)

Fix ω such that $\kappa(\omega) < \infty$.

- $\kappa(\cdot + \omega) - \kappa(\omega)$ is the Laplace exponent of a Lévy process; call it ξ .

Solution of the equation

Define the cumulant κ : $\mathbb{E}[\sum_u e^{qZ_u(t)}] = e^{t\kappa(q)}$. It satisfies

$$\kappa(q) = aq + \int_{\mathcal{P}} \left\{ \sum_{i \geq 1} p_i^q - 1 + (1 - p_1)q \right\} \nu(d\mathbf{p}).$$

Theorem (Bertoin, W. '15)

Fix ω such that $\kappa(\omega) < \infty$.

- $\kappa(\cdot + \omega) - \kappa(\omega)$ is the Laplace exponent of a Lévy process; call it ξ .
- Let

$$\langle \mu_t, f \rangle = e^{t\kappa(\omega)} \mathbb{E}[e^{-\omega\xi(t)} f(e^{\xi(t)})] = \mathbb{E}\left[\sum_u f(e^{Z_u(t)})\right].$$

This is the unique solution of the growth-fragmentation equation.

A martingale for the fragmentation

- Let

$$W(\omega, t) = e^{-t\kappa(\omega)} \sum_u e^{\omega Z_u(t)}, \quad t \geq 0.$$

The process W is the **additive martingale**.

A martingale for the fragmentation

- Let

$$W(\omega, t) = e^{-t\kappa(\omega)} \sum_u e^{\omega Z_u(t)}, \quad t \geq 0.$$

The process W is the additive martingale.

- It induces **new measure** \mathbb{P}_ω for \mathcal{Z} : if F_t is measurable with respect to paths up to time t ,

$$\mathbb{E}_\omega[F_t(\mathcal{Z})] = \mathbb{E}[F_t(\mathcal{Z})W(\omega, t)].$$

Construction of a spine process, I

Let ξ be the Lévy process with Laplace exponent $\kappa(\omega + \cdot) - \kappa(\omega)$.

- Lévy measure: $\Pi(dz) = \sum_{i \geq 1} e^{\omega z} \nu(\log p_i \in dz)$.

Construction of a spine process, I

Let ξ be the Lévy process with Laplace exponent $\kappa(\omega + \cdot) - \kappa(\omega)$.

- Lévy measure: $\Pi(dz) = \sum_{i \geq 1} e^{\omega z} \nu(\log p_i \in dz)$.
- Jump process: $M(ds, dz)$, a Poisson point process with intensity measure $ds \Pi(dz)$.

Construction of a spine process, II

We will define a *decoration* of the jump process M :

- $g_i(z) = \frac{e^{\omega z} \nu(\log p_i \in dz)}{\Pi(dz)}$, a Radon-Nikodym derivative

Construction of a spine process, II

We will define a *decoration* of the jump process M :

- $g_i(z) = \frac{e^{\omega z} \nu(\log p_i \in dz)}{\Pi(dz)}$, a Radon-Nikodym derivative
- $\mu(s, z, di, d\mathbf{p}) = g_i(z) \zeta(di) \nu(d\mathbf{p} \mid \log p_i = z)$,
where ζ is counting measure.

Construction of a spine process, II


We will define a *decoration* of the jump process M :

- $g_i(z) = \frac{e^{\omega z} \nu(\log p_i \in dz)}{\Pi(dz)}$, a Radon-Nikodym derivative
- $\mu(s, z, di, d\mathbf{p}) = g_i(z) \zeta(di) \nu(d\mathbf{p} \mid \log p_i = z)$,
where ζ is counting measure.
- Let $N(ds, dz, di, d\mathbf{p})$ be the μ -randomisation of M .

Construction of a spine process, II

We will define a *decoration* of the jump process M :


- $g_i(z) = \frac{e^{\omega z} \nu(\log p_i \in dz)}{\Pi(dz)}$, a Radon-Nikodym derivative
- $\mu(s, z, di, d\mathbf{p}) = g_i(z) \zeta(di) \nu(d\mathbf{p} \mid \log p_i = z)$, where ζ is counting measure.
- Let $N(ds, dz, di, d\mathbf{p})$ be the μ -randomisation of M .



Construction of a spine process, II

We will define a *decoration* of the jump process M :

- $g_i(z) = \frac{e^{\omega z} \nu(\log p_i \in dz)}{\Pi(dz)}$, a Radon-Nikodym derivative
- $\mu(s, z, di, d\mathbf{p}) = g_i(z) \zeta(di) \nu(d\mathbf{p} \mid \log p_i = z)$,
where ζ is counting measure.
- Let $N(ds, dz, di, d\mathbf{p})$ be the μ -randomisation of M .



Construction of a spine process, III

We construct a new process under \mathbb{P}_ω :

- Let $(\mathcal{Z}^{[s,j]})_{s \geq 0, j \geq 1}$ be a collection of independent compensated fragmentations under \mathbb{P} .

Construction of a spine process, III

We construct a new process under \mathbb{P}_ω :

- Let $(\mathcal{Z}^{[s,j]})_{s \geq 0, j \geq 1}$ be a collection of independent compensated fragmentations under \mathbb{P} .
- Define


$$\tilde{\mathcal{Z}}(t) = \delta_{\xi(t)} + \int N(ds, dz, di, d\mathbf{p}) \sum_{j \neq i} [\mathcal{Z}^{[s,j]}(t-s) + \xi(s-) + \log p_j].$$

Construction of a spine process, III

We construct a new process under \mathbb{P}_ω :

- Let $(\mathcal{Z}^{[s,j]})_{s \geq 0, j \geq 1}$ be a collection of independent compensated fragmentations under \mathbb{P} .
- Define

$$\tilde{\mathcal{Z}}(t) = \delta_{\xi(t)} + \int N(ds, dz, di, d\mathbf{p}) \sum_{j \neq i} [\mathcal{Z}^{[s,j]}(t-s) + \xi(s-) + \log p_j].$$




Construction of a spine process, III

We construct a new process under \mathbb{P}_ω :

- Let $(\mathcal{Z}^{[s,j]})_{s \geq 0, j \geq 1}$ be a collection of independent compensated fragmentations under \mathbb{P} .
- Define

$$\tilde{\mathcal{Z}}(t) = \delta_{\xi(t)} + \int N(ds, dz, di, d\mathbf{p}) \sum_{j \neq i} [\mathcal{Z}^{[s,j]}(t-s) + \xi(s-) + \log p_j].$$




Construction of a spine process, III

We construct a new process under \mathbb{P}_ω :

- Let $(\mathcal{Z}^{[s,j]})_{s \geq 0, j \geq 1}$ be a collection of independent compensated fragmentations under \mathbb{P} .
- Define

$$\tilde{\mathcal{Z}}(t) = \delta_{\xi(t)} + \int N(ds, dz, di, \mathbf{dp}) \sum_{j \neq i} [\mathcal{Z}^{[s,j]}(t-s) + \xi(s-) + \log p_j].$$



Construction of a spine process, III

We construct a new process under \mathbb{P}_ω :

- Let $(\mathcal{Z}^{[s,j]})_{s \geq 0, j \geq 1}$ be a collection of independent compensated fragmentations under \mathbb{P} .
- Define

$$\tilde{\mathcal{Z}}(t) = \delta_{\xi(t)} + \int N(ds, dz, di, d\mathbf{p}) \sum_{j \neq i} [\mathcal{Z}^{[s,j]}(t-s) + \xi(s-) + \log p_j].$$

We distinguish within $\tilde{\mathcal{Z}}$ the particle with position ξ : let U_t be such that $\tilde{\mathcal{Z}}_{U_t}(t) = \xi(t)$.

Many-to-one theorem

Theorem (W. '16)

For F_t measurable with respect to paths up to time t , and u any label,

$$\mathbb{E}_\omega[F_t(\tilde{\mathcal{Z}})\mathbb{1}_{\{U_t=u\}}] = e^{-t\kappa(\omega)}\mathbb{E}[F_t(\mathcal{Z})e^{\omega Z_u(t)}].$$

Many-to-one theorem

Theorem (W. '16)

For F_t measurable with respect to paths up to time t , and u any label,

$$\mathbb{E}_\omega[F_t(\tilde{\mathcal{Z}})\mathbb{1}_{\{U_t=u\}}] = e^{-t\kappa(\omega)}\mathbb{E}[F_t(\mathcal{Z})e^{\omega\mathcal{Z}_u(t)}].$$

Corollary

Summing over u ,

$$\mathbb{E}_\omega[F_t(\tilde{\mathcal{Z}})] = e^{-t\kappa(\omega)}\mathbb{E}[F_t(\mathcal{Z})W(\omega, t)] = \mathbb{E}_\omega[F_t(\mathcal{Z})].$$

That is, $\tilde{\mathcal{Z}} \stackrel{d}{=} \mathcal{Z}$ under \mathbb{P}_ω .

Many-to-one theorem

Theorem (W. '16)

For F_t measurable with respect to paths up to time t , and u any label,

$$\mathbb{E}_\omega[F_t(\tilde{\mathcal{Z}})\mathbb{1}_{\{U_t=u\}}] = e^{-t\kappa(\omega)}\mathbb{E}[F_t(\mathcal{Z})e^{\omega\mathcal{Z}_u(t)}].$$

Corollary

For Borel f ,


$$e^{t\kappa(\omega)}\mathbb{E}_\omega[e^{-\omega\tilde{\mathcal{Z}}_{U_t}(t)}f(\tilde{\mathcal{Z}}_{U_t}(t))] = \mathbb{E}\left[\sum_u f(\mathcal{Z}_u(t))\right].$$

Discussion of proof, I

We define *truncation* of the processes.

In the truncated process $\mathcal{Z}^{(b)}$, a child particle is kept only if:

- its displacement from the parent is less than $-b$, or
- it is the largest child




Discussion of proof, I

We define *truncation* of the processes.

In the truncated process $\mathcal{Z}^{(b)}$, a child particle is kept only if:

- its displacement from the parent is less than $-b$, or
- it is the largest child




Discussion of proof, I

We define *truncation* of the processes.

In the truncated process $\mathcal{Z}^{(b)}$, a child particle is kept only if:

- its displacement from the parent is less than $-b$, or
- it is the largest child




Discussion of proof, I

We define *truncation* of the processes.

In the truncated process $\mathcal{Z}^{(b)}$, a child particle is kept only if:

- its displacement from the parent is less than $-b$, or
- it is the largest child



Discussion of proof, II

The truncation operation can be applied to:

- \mathcal{Z} , under \mathbb{P} — let $\kappa^{(b)}$ be the cumulant of $\mathcal{Z}^{(b)}$
- $\tilde{\mathcal{Z}}$, under \mathbb{P}_ω

They are related by:

Lemma

Let

$$\zeta = \inf\{t \geq 0 : U_t \text{ not in } \tilde{\mathcal{Z}}^{(b)}\}.$$

Then,

$$\mathbb{E}_\omega [F_t(\tilde{\mathcal{Z}}^{(b)}) \mathbb{1}_{\{U_t=u\}} \mid \zeta > t] = e^{-t\kappa^{(b)}(\omega)} \mathbb{E} [F_t(\mathcal{Z}) e^{\omega \mathcal{Z}_u^{(b)}(t)}]$$

Further questions

- Derivative martingale

$$\partial W(\omega, t) = e^{-t\kappa(\omega)} \sum_u (\omega Z_u(t) - t\kappa'(\omega)) e^{\omega Z_u(t)}$$

- KPP equation
- 'Non-homogeneous' fragmentations (self-similar done by Bertoin–Budd–Curien–Kortchemski '16)

Further questions

- Derivative martingale

$$\partial W(\omega, t) = e^{-t\kappa(\omega)} \sum_u (\omega Z_u(t) - t\kappa'(\omega)) e^{\omega Z_u(t)}$$

- KPP equation
- ‘Non-homogeneous’ fragmentations (self-similar done by Bertoin–Budd–Curien–Kortchemski '16)



J. Bertoin

Compensated fragmentation processes and limits of dilated fragmentations



J. Bertoin, A. R. Watson

Probabilistic aspects of critical growth-fragmentation equations



A. R. Watson

Tilting of compensated fragmentations [in preparation]

Thank you!