

## Section A

1. Define subspaces  $U, W \leq \mathbb{R}^3$  by:

$$U = \text{span}\{(1, 1, 1), (1, 2, 0)\}, \quad W = \text{span}\{(1, 0, 2)\}.$$

Is the sum  $U + W$  direct? (You must justify your answer.) [4]

2. With  $U \leq \mathbb{R}^3$  as in question 1,  $v = (1, 2, 3)$  and  $w = (0, 2, 1)$ , is  $v \equiv w \pmod{U}$ ? (You must justify your answer.) [4]

3. For  $x, y \in \mathbb{C}^3$ , let

$$\langle x, y \rangle = x_1y_1 + x_2y_2 + \bar{x}_3y_3.$$

Is  $\langle \cdot, \cdot \rangle$  an inner product on  $\mathbb{C}^3$ ? (You must justify your answer.) [4]

4. Show that  $A \in M_{2 \times 2}(\mathbb{C})$  given by

$$\frac{1}{\sqrt{5}} \begin{pmatrix} 1 & 2i \\ 2i & 1 \end{pmatrix}$$

is a unitary matrix and compute its eigenvalues. [4]

5. With  $U \leq \mathbb{R}^3$  as in question 1, write down a non-zero element of  $\text{ann } U$ .

**Hint:** any  $\alpha \in (\mathbb{R}^3)^*$  is of the form  $\alpha(x) = \alpha_1x_1 + \alpha_2x_2 + \alpha_3x_3$ . [4]

6. Define a quadratic form  $Q$  on  $\mathbb{R}^2$  by

$$Q(x) = 4x_1^2 + 4x_1x_2 + x_2^2.$$

Compute the rank and signature of  $Q$ . [4]

### Section B

7. Let  $V$  be a vector space over a field  $\mathbb{F}$ .

- (a) Let  $V_1, \dots, V_k \leq V$  be subspaces of  $V$ .

Show that the sum  $V_1 + \dots + V_k$  is direct if and only if, whenever  $v_i \in V_i$ ,  $1 \leq i \leq k$ , satisfy

$$v_1 + \dots + v_k = 0$$

then each  $v_i = 0$ . [6]

- (b) Let  $u, w \in V$  and  $V_1, V_2 \leq V$  subspaces such that  $u + V_1 = w + V_2$ .

Prove that  $V_1 = V_2$ . [6]

- (c) Let  $\alpha \in V^*$  be non-zero. Prove that

$$\dim(V / \ker \alpha) = 1.$$

[6]

8. (a) Let  $a, b, c, d, e$  be positive real numbers. Show that

$$25 \leq (a + b + c + d + e)(1/a + 1/b + 1/c + 1/d + 1/e).$$

For what  $a, b, c, d, e$  do we get equality? [5]

- (b) Find the QR decomposition for the matrix  $A$  given by

$$A = \begin{pmatrix} 1 & 1 & 3 \\ -1 & 0 & 1 \\ 0 & -1 & -1 \end{pmatrix}.$$

[7]

- (c) Let  $U = \text{span}\{(1, -1, 0), (1, 0, -1)\} \leq \mathbb{R}^3$  and  $v = (2, 1, 2)$ .

Find the closest point of  $U$  to  $v$  (where we define distance in  $\mathbb{R}^3$  using the dot product). [6]

9. Let  $B$  be the symmetric bilinear form on  $\mathbb{R}^3$  defined by the matrix

$$A = \begin{pmatrix} -3 & 2 & -2 \\ 2 & -1 & 1 \\ -2 & 1 & -1 \end{pmatrix}.$$

Thus  $B(x, y) = \mathbf{x}^T A \mathbf{y}$ .

- (a) Find a diagonalising basis for  $B$ . [10]  
(b) Compute the rank and signature of  $B$ . [4]  
(c) Compute the radical of  $B$ . [4]