

Disconnecting the G_2 moduli space

Johannes Nordström

27 June 2017

Joint work in progress with
Diarmuid Crowley and Sebastian Goette

C-N, *New invariants of G_2 -structures*, *Geom. Topol.* 19 (2015)

C-G-N, *An analytic invariant of G_2 -manifolds*, arXiv:1505.02734

These slides available at
<http://people.bath.ac.uk/jl1pn20/disconnect.pdf>

The G_2 moduli space

Let M be a smooth closed 7-manifold admitting metrics with holonomy G_2 .
The moduli space

$$\mathcal{M} := \{\text{Holonomy } G_2 \text{ metrics on } M\} / \text{Diff}(M)$$

is an orbifold, locally homeomorphic to finite quotients of $H_{dR}^3(M)$.
So far little is known about the *global* properties of \mathcal{M} .

Main results:

Exhibit examples of closed G_2 -manifolds with \mathcal{M} disconnected, both

- by studying homotopies of G_2 -structures, and
- where the G_2 -structures are indistinguishable using homotopy theory

Outline:

1. Background and examples
2. Invariants of G_2 -structures
3. Constructions
4. Computation

1. Background and examples

The group G_2

$G_2 := \text{Aut } \mathbb{O}$, \mathbb{O} = octonions, normed division algebra of real dimension 8.

G_2 acts on $\text{Im } \mathbb{O} \cong \mathbb{R}^7$, preserving metric, orientation, cross product

$$a \times b := \text{Im}(ab), \text{ and}$$

$$\varphi_0(a, b, c) := \langle a \times b, c \rangle.$$

In terms of basis $e^1, \dots, e^7 \in (\mathbb{R}^7)^*$

$$\varphi_0 = e^{123} + e^{145} + e^{167} + e^{246} - e^{257} - e^{347} - e^{356} \in \Lambda^3(\mathbb{R}^7)^*.$$

Peculiar algebra facts:

- G_2 is not just contained in stabiliser of φ_0 in $GL(7, \mathbb{R})$, but equality holds.
- The $GL(7, \mathbb{R})$ -orbit of φ_0 is open in $\Lambda^3(\mathbb{R}^7)^*$.

G_2 -structures and holonomy

G_2 is an exceptional case in Berger's list of Riemannian holonomy groups. A metric with holonomy G_2 is always Ricci-flat.

Parallel tensor fields on Riemannian manifold $M \leftrightarrow$ invariants of $Hol(M)$.

A 3-form $\varphi \in \Omega^3(M^7)$ such that $(T_x M, \varphi) \cong (\mathbb{R}^7, \varphi_0)$ for all $x \in M$ defines a G_2 -structure. (This is an *open* condition on φ)

Because $G_2 \subset SO(7)$, this induces a metric and orientation.

$Hol(M) \subseteq G_2 \Leftrightarrow$ metric induced by some G_2 -structure φ such that $\nabla\varphi = 0$. Then call φ *torsion-free*. This is equivalent to the first-order non-linear PDE


$$d\varphi = d^*\varphi = 0.$$

[Bryant \(1985\)](#): Local examples

[Bryant-Salamon \(1987\)](#): Complete examples

[Joyce \(1994\)](#): Examples on closed manifolds

Two perspectives on G_2 -structures



The spin representation Δ of $Spin(7)$ is real of rank 8.
 $Spin(7)$ acts transitively on $S^7 \subset \Delta$ with stabiliser G_2 .

$$G_2 = \text{stabiliser in } GL(7, \mathbb{R}) \text{ of } \varphi_0 \in \Lambda^3(\mathbb{R}^7)^* = \text{stabiliser in } Spin(7) \text{ of a unit spinor } s_0$$

$$G_2\text{-structure on } M^7 \Leftrightarrow \text{positive } \varphi \in \Omega^3(M) \Leftrightarrow \begin{array}{l} \text{metric } g \\ + \text{ spin structure} \\ + \text{ unit spinor field } s \end{array}$$

$$\text{Holonomy } \subseteq G_2 \Leftrightarrow d\varphi = d^*\varphi = 0 \Leftrightarrow \nabla s = 0$$

Useful for

differential geometry

homotopy theory

Homotopies of G_2 -structures

Let M be a closed 7-dimensional spin manifold.

Given a metric g , the spinor bundle SM is a real vector bundle of rank 8. Two G_2 -structures inducing the same metric and spin structure are homotopic if the corresponding unit spinors can be connected by a path of non-vanishing spinors.

All metrics on M are homotopic, so if we fix the spin structure

$$\left\{ \begin{array}{l} \text{Homotopy classes of} \\ G_2\text{-structures on } M \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \text{Homotopy classes of non-} \\ \text{vanishing sections of } SM \end{array} \right\} \leftrightarrow \mathbb{Z}$$

by counting (with signs) the zeros of an interpolating section of a rank 8 bundle on $M \times [0, 1]$.

$\text{Diff}(M)$ can act by non-trivial translations.

Each component of the G_2 moduli space \mathcal{M} maps to a fixed class of G_2 -structures modulo homotopies *and* diffeomorphisms.

Classification of 2-connected manifolds

Let M be a closed smooth 7-manifold with $\pi_1(M) = \pi_2(M) = 0$ and $H^4(M)$ torsion-free. Remaining algebraic topology captured by $b_3(M)$.

Let $d(M) :=$ greatest integer dividing $\frac{1}{2}p_1(M) \in H^4(M)$
($d(M) := 0$ if $p_1(M) = 0$).

Theorem (Wilkins, 1972)

*Such M are classified up to homeomorphism by $(b_3(M), d(M)) \in \mathbb{N} \times 2\mathbb{N}$.
The number of inequivalent smooth structures on the topological manifold underlying M is*

$$\text{GCD}(28, \text{Numerator}\left(\frac{d(M)}{4}\right)).$$

Theorem (C-N)

The number of G_2 -structures up to homotopy+diffeomorphism on such M is

$$24 \text{ Numerator}\left(\frac{d(M)}{112}\right).$$

A 2-connected example

Example (C-G-N)

Let M be the unique smooth closed 2-connected 7-manifold with $H^4(M) = \mathbb{Z}^{97}$ and $d = 2$.

There are G_2 metrics g_1, g_2, g_3 on M such that

- A** the G_2 -structures φ_1, φ_2 associated to g_1 and g_2 are not equivalent under homotopies and diffeomorphisms; thus g_1 and g_2 are in different components of the G_2 moduli space \mathcal{M}
- B** the G_2 -structures φ_1 and φ_3 are homotopic, but nevertheless g_1 and g_3 lie in different components of \mathcal{M} .

So for this manifold, the moduli space \mathcal{M} has at least 3 connected components.

Ingredients

Invariants

- A** The G_2 -structures are distinguished by a homotopy and diffeomorphism invariant $\nu(\varphi) \in \mathbb{Z}/48\mathbb{Z}$.
- B** An analytic refinement $\widehat{\nu}(\varphi) \in \mathbb{Z}$ of $\nu(\varphi)$ is invariant under diffeomorphisms and under deformations through torsion-free G_2 -structures (but not under arbitrary homotopies), and can distinguish components of \mathcal{M} even when the G_2 -structures are homotopic.

Construction

The “twisted connected sum construction” of Kovalev and Corti-Haskins-N-Pacini produces large numbers of 2-connected G_2 -manifolds for which the invariants can be evaluated.

A more complicated version produces some 2-connected examples where $\widehat{\nu}$ takes a range of values.

2. Invariants of G_2 -structures

The homotopy invariant

Let X closed spin 8-manifold, and $n(X)$ the signed count of zeros of a transverse positive spinor field (\Leftrightarrow Euler class of rank 8 bundle S^+X).
Atiyah-Singer index theorem + $Spin(8)$ characteristic class computation

$$\rightsquigarrow -48 \operatorname{ind} D_X^+ = \chi(X) - 3\sigma(X) - 2n(X). \quad (*)$$

Let W be a compact spin 8-manifold with boundary M , s a transverse positive spinor field on W , and φ the G_2 -structure on M induced by $s|_M$.
Let $n(W, \varphi)$ be the signed count of zeros of s . (*) implies that

$$\nu(\varphi) := \chi(W) - 3\sigma(W) - 2n(W, \varphi) \pmod{48}$$

is independent of choice of coboundary W .

On a fixed M , ν takes the 24 values allowed by $\nu(\varphi) = \sum_{i=0}^3 b_i(M) \pmod{2}$.

If M is 2-connected with $H^4(M)$ torsion-free and d a divisor of 112, then ν distinguishes all classes.

Analytic invariant of G_2 -structures

Given a metric on a closed spin M^7 , define

$D =$ Dirac operator

$B : \Omega^{\text{ev}} \rightarrow \Omega^{\text{ev}} =$ odd signature operator, $(-1)^k(*d - d*)$ on Ω^{2k}

$h(D) = \dim \ker(D) \in \mathbb{Z}$

$\eta(D) := \eta(D, 0) \in \mathbb{R}$, defined by analytic continuation from

$$\eta(D, s) := \sum_{\lambda \in \text{Spec} D \setminus \{0\}} (\text{sign} \lambda) |\lambda|^{-s} \quad \text{for } \text{Re } s \gg 0.$$

For a G_2 -structure φ on M , define $MQ(\varphi) \in \mathbb{R}$ in terms of Mathai-Quillen current.

Definition

$$\hat{\nu}_0(\varphi) := -24\eta(D) + 3\eta(B) + 2MQ(\varphi) \in \mathbb{R}$$

$$\hat{\nu}(\varphi) := \hat{\nu}_0(\varphi) - 24h(D) \in \mathbb{R}$$

Analytic invariant as refinement

$$\widehat{\nu}_0(\varphi) := -24\eta(D) + 3\eta(B) + 2MQ(\varphi) \in \mathbb{R}$$

Reversing orientation changes the sign of $\widehat{\nu}_0$.

All terms are continuous in φ , except that the first jumps by 24 when an eigenvalue of D changes between zero and non-zero.

$$\widehat{\nu}(\varphi) := \widehat{\nu}_0(\varphi) - 24h(D) \in \mathbb{R}$$

$\widehat{\nu}$ is continuous in φ except for jumps by 48.

Theorem (C-G-N)

Let φ be G_2 -structure on a closed M^7 . Then

$$\nu(\varphi) = \widehat{\nu}(\varphi) \pmod{48}.$$

(In particular $\widehat{\nu}, \widehat{\nu}_0 \in \mathbb{Z}$.)

Analytic invariant as refinement

$$\widehat{\nu}(\varphi) := -24(\eta + h)(D) + 3\eta(B) + 2MQ(\varphi) \in \mathbb{R}$$

$$\nu(\varphi) := \chi(W) - 3\sigma(W) - 2n(W, \varphi) \in \mathbb{Z}/48\mathbb{Z}.$$

Proof.

For $\partial W = M$ with metric that is product on collar of M

$$\begin{array}{lll} \sigma(W) & = & \int_W L(\nabla) \quad - \quad \eta(B) \\ \text{ind } D_W^+ & = & \int_W \widehat{A}(\nabla) \quad - \quad \frac{1}{2}(\eta + h)(D) \\ n(W, \varphi) & = & \int_W e_+(\nabla) \quad - \quad MQ(\varphi) \end{array}$$

Chern-Weil term boundary correction

Linear combination of Chern-Weil terms gives $\int_W e(\nabla) = \chi(W)$ (essentially by characteristic class formula (*) used to show that ν is well-defined), so

$$\widehat{\nu}(\varphi) = \chi(W) - 3\sigma(W) - 2n(W, \varphi) + 48 \text{ind } D_W^+ \in \mathbb{Z} . \quad \square$$

Analytic invariant of torsion-free G_2 -structures

$$\widehat{\nu}_0(\varphi) := -24\eta(D) + 3\eta(B) + 2MQ(\varphi) \in \mathbb{Z}$$

$$\widehat{\nu}(\varphi) := \widehat{\nu}_0(\varphi) - 24h(D) \in \mathbb{Z}$$

For torsion-free φ

- $MQ(\varphi) = 0$
- $h(D) = 1 + b_1(M)$ (so 1 when $Hol = G_2$)
- $\eta(D)$ does not jump

Therefore $\widehat{\nu}_0$ and $\widehat{\nu}$ are constant on connected components of \mathcal{M} , and can distinguish components even when the associated G_2 -structures are homotopic.

Even if we are only interested in ν , it may be easier to evaluate the intrinsic formula for $\widehat{\nu}$ than to find a spin coboundary to compute ν .

Similarities with e.g. the use of Donnelly's analytic refinement of the Eells–Kuiper invariant by Kreck–Stolz and Goette–Kitchloo–Shankar.

3. Constructions

G_2 and $SU(3)$

The action of $SU(3)$ on $\mathbb{C}^3 \cong \mathbb{R}^6$ preserves

$$\omega_0 := \frac{i}{2}(dz^1 \wedge d\bar{z}^1 + dz^2 \wedge d\bar{z}^2 + dz^3 \wedge d\bar{z}^3) \in \Lambda^2(\mathbb{R}^6)^*$$

$$\Omega_0 := dz^1 \wedge dz^2 \wedge dz^3 \in \Lambda^3(\mathbb{R}^6)^* \otimes \mathbb{C}$$

On $\mathbb{R}^7 = \mathbb{R} \oplus \mathbb{C}^3$,

$$e^1 \wedge \omega_0 + \operatorname{Re} \Omega_0 \cong e^1 \wedge (e^{23} + e^{45} + e^{67}) + e^{246} - e^{257} - e^{347} - e^{356} = \varphi_0,$$

the 3-form preserved by G_2 .

The stabiliser in G_2 of a non-zero vector is $SU(3)$.

If X is a Calabi-Yau 3-fold (6-manifold with $\operatorname{Hol}(X) = SU(3)$) then $\operatorname{Hol}(S^1 \times X) = SU(3) \subset G_2$, so $S^1 \times X$ has a torsion-free G_2 -structure.

But we are more interested in manifolds with full holonomy G_2 .

Proposition (Joyce)


If M^7 is closed and $\operatorname{Hol}(M) \subseteq G_2$ then

$$\operatorname{Hol}(M) = G_2 \Leftrightarrow \pi_1(M) \text{ finite}$$

Twisted connected sums

Donaldson, Kovalev, Corti-Haskins-N-Pacini

- Construct simply-connected, complete Calabi-Yau 3-folds V , with “asymptotically cylindrical end” $\mathbb{R} \times S^1 \times K3$.
- $\text{Hol}(S^1 \times V) = \text{SU}(3) \subset G_2$, so $S^1 \times V$ has torsion-free G_2 -structure
- Find pairs of such V_{\pm} , with a diffeomorphism F of the cylindrical ends of $S^1 \times V_+$ and $S^1 \times V_-$ ensuring
 - Gluing G_2 -structures on the halves with “neck length” $T \gg 0$ defines φ_T on M with $\nabla\varphi_T$ exponentially small in T .
 - $M = S^1 \times V_+ \cup_F S^1 \times V_-$ is simply-connected (F is “twisted”)



- Perturb to φ_T so that $d\varphi_T = d^*\varphi_T = 0$. Then $\text{Hol}(M) = G_2$.

Matching

The ACyl end of $S^1 \times V_{\pm}$ is $\mathbb{R} \times S^1 \times S^1 \times K3_{\pm} \cong \mathbb{R} \times T_{\pm}^2 \times K3_{\pm}$.

Glue the cylindrical ends using a product isometry

$$F := (-1) \times m \times r : \mathbb{R} \times T_+^2 \times K3_+ \rightarrow \mathbb{R} \times T_-^2 \times K3_-,$$

where $m : T_+^2 \rightarrow T_-^2$ is the reflection $S^1 \times S^1 \rightarrow S^1 \times S^1$, $(u, v) \mapsto (v, u)$.

m swaps “internal” and “external” circles $\Rightarrow \pi_1 M = 0$ by van Kampen.

Matching problem: Find pairs V_+ and V_- such that there is an isometry $r : K3_+ \rightarrow K3_-$ making F an isomorphism of the ACyl G_2 -structures.

Kovalev (2003): Use Fano 3-folds to produce examples of pairs V_+ , V_- with solution to the matching problem.

Corti-Haskins-N-Pacini (2014): Millions of examples from weak Fano 3-folds. Topological type determined in many cases. Many gluings give same smooth manifold.

Invariants of twisted connected sums


Theorem (C-N)

Any twisted connected sum has $\nu = 24 \in \mathbb{Z}/48\mathbb{Z}$.

Theorem (C-G-N)

Any twisted connected sum has $\hat{\nu} = -24 \in \mathbb{Z}$.

Analytic computation reveals the result to be related to a geometric feature:
 $m : T_+^2 \rightarrow T_-^2$ aligns “external” circle tangents ∂_v at right angle.



Inevitable, because m is an isometry of rectangular tori, and is not allowed to align the external circles: otherwise M would have an S^1 factor.

Tori with symmetries


Warm-up question:

Let $a: S^1 \rightarrow S^1$ be the antipodal map $z \mapsto -z$.

Let $T^2 := S^1 \times S^1 / a \times a$ where the S^1 factors have circumference 1 and x .

For how many different x does T^2 have rotation symmetries other than ± 1 ?


$$x = 1, \sqrt{3}, \text{ or } \frac{1}{\sqrt{3}}$$



Isometries between tori

Consider a pair of tori that are either rectangular (metric product $S^1 \times S^1$) or quotient of a rectangular one by an involution ($S^1 \times S^1/a \times a$). For isometries between such tori, at what angles θ can the sides of the rectangles be aligned?

Can achieve $\theta = \frac{\pi}{4}$ with an involution on one side.




Isometries between tori

Consider a pair of tori that are either rectangular (metric product $S^1 \times S^1$) or quotient of a rectangular one by an involution ($S^1 \times S^1/a \times a$).

For isometries between such tori, at what angles θ can the sides of the rectangles be aligned?


With involutions on both sides, one can achieve $\theta = \frac{\pi}{3}$.



Isometries between tori

Consider a pair of tori that are either rectangular (metric product $S^1 \times S^1$) or quotient of a rectangular one by an involution ($S^1 \times S^1/a \times a$). For isometries between such tori, at what angles θ can the sides of the rectangles be aligned?

With involutions on both sides, one can achieve $\theta = \frac{\pi}{6}$.



Extra-twisted connected sums

Suppose V is an ACyl Calabi-Yau with an involution τ , that acts on the asymptotic cross-section $S^1 \times K3$ by $a \times \text{Id}_{K3}$.

Then $S^1 \times V / a \times \tau$ is an ACyl G_2 -manifold with cross-section

$$(S^1 \times S^1 / a \times a) \times K3 = T^2 \times K3.$$

Let M_{\pm} be a pair of ACyl G_2 -manifolds of this form, or of the form $S^1 \times V$.

Let $m : T_+^2 \rightarrow T_-^2$ be a reflection. Depending on the circumferences of the circles, m can align the external circle directions at angle $\theta = \frac{\pi}{3}, \frac{\pi}{4}$ or $\frac{\pi}{6}$.

θ -matching problem: Find pairs V_+ and V_- with involution, and with an isometry $r : K3_+ \rightarrow K3_-$ such that $(-1) \times m \times r$ is an isomorphism of the limits of the ACyl G_2 -structures of M_+ and M_- .

Can obtain some ACyl Calabi-Yau manifolds with involution, and solutions to the matching problem, from branched double covers of Fano 3-folds.

Examples from extra-twisted connected sums

For each $\theta \neq \frac{\pi}{2}$, a range of values of $\widehat{\nu}$ can be realised by θ -TCSs.

Claim A

For a certain $\frac{\pi}{4}$ -TCS M_2 , compute that $\pi_2 M_2 = 0$, $H^4(M_2) \cong \mathbb{Z}^{97}$, $d(M_2) = 2$, and $\nu(\varphi_2) = 36 \in \mathbb{Z}/48\mathbb{Z}$.

Among the millions of 2-connected ordinary TCS, find one that also has $H^4(M_1) = \mathbb{Z}^{97}$ and $d(M_1) = 2$. By the classification of 2-connected 7-manifolds, it is diffeomorphic to M_2 .

However, the ordinary TCS has $\nu(\varphi_1) = 24 \in \mathbb{Z}/48\mathbb{Z}$, so the G_2 -structures φ_1 and φ_2 are not homotopic (not even after changing the diffeomorphism that identifies M_1 and M_2). Hence the constructed G_2 -metrics g_1 and g_2 lie in different components of the G_2 moduli space on M .

Examples from extra-twisted connected sums

Claim B

For a certain $\frac{\pi}{6}$ -TCS M_3 , compute that $\pi_2 M_3 = 0$, $H^4(M_3) \cong \mathbb{Z}^{97}$, $d(M_3) = 2$, and $\widehat{\nu}(\varphi_3) = -72 \in \mathbb{Z}$.

M_3 is thus diffeomorphic to M_1 above.

On this manifold, there are precisely 24 $\text{Numerator}(\frac{d}{112}) = 24$ classes of G_2 -structures modulo homotopy and diffeomorphism, all distinguished by ν .

Since $\nu(\varphi_1) = \nu(\varphi_3) = 24 \in \mathbb{Z}/48\mathbb{Z}$, the diffeomorphism $M_1 \cong M_3$ can therefore be chosen so that the torsion-free G_2 -structures are homotopic.

However, the ordinary TCS has $\widehat{\nu}(\varphi_1) = -24$, so the two torsion-free G_2 -structures lie in different components of the G_2 moduli space.

$\frac{\pi}{3}$ -TCSs have 3-torsion in $H^4(M)$, making it harder to apply classification results to find different examples realising the same smooth manifold.

4. Computation

Limits of the eta invariants

$M_{\pm} := S^1 \times V_{\pm}$ or $S^1 \times V_{\pm}/a \times \tau$, with asymptotic limit $\mathbb{R} \times T_{\pm}^2 \times K3$.
 $m : T_+^2 \rightarrow T_-^2$ reflection, aligning external circle factors at angle $\theta \in (0, \frac{\pi}{2}]$.
Construct family of torsion-free G_2 -structures φ_T on M the result of gluing M_+ to M_- by $(-1) \times m \times r$ with “neck length” T .

Theorem

Let $\rho := \pi - 2\theta$. Then $\eta(D) \rightarrow \frac{\rho}{\pi}$ as $T \rightarrow \infty$.

Let $R_{\pm} : H^2(K3; \mathbb{R}) \rightarrow H^2(K3; \mathbb{R})$ be reflection in $\text{Im}(H^2(V_{\pm}) \rightarrow H^2(K3))$

Theorem

Define a unitary map $U : H^2(K3; \mathbb{C}) \rightarrow H^2(K3; \mathbb{C})$ by $e^{\pm i\rho} R_+ R_-$ on $H^{2,\pm}(K3; \mathbb{C})$. Then

$$\eta(B) \rightarrow \frac{1}{\pi} \sum_{\substack{\lambda \in \text{Spec } U \\ \lambda \neq -1}} \arg \lambda$$

as $T \rightarrow \infty$, where the branch of \arg takes values in $(-\pi, \pi)$.

Evaluating $\widehat{\nu}$

$U := e^{\pm i\rho} R_+ R_-$ on $H^{2,\pm}(K3; \mathbb{C})$. The theorems imply

$$\widehat{\nu}_0 = -24\eta(D) + 3\eta(B) = -24\frac{\rho}{\pi} + \frac{3}{\pi} \sum_{\substack{\lambda \in \text{Spec } U \\ \lambda \neq -1}} \arg \lambda.$$

If $\theta = \frac{\pi}{2}$ then $\rho = \pi - 2\theta = 0$, and U is the real orthogonal map $R_+ R_-$. Hence eigenvalues are ± 1 or occur in conjugate pairs, so $\sum \arg \lambda = 0$, and

$$\widehat{\nu}_0 = 0.$$

In general

$$\sum_{\substack{\lambda \in \text{Spec } U \\ \lambda \neq -1}} \arg \lambda = \sum \pm \rho + \sum_{\substack{\lambda \in \text{Spec } R_+ R_- \\ \lambda \neq -1}} \arg \lambda + \pi b = -16\rho + \pi b,$$

where $b \in \mathbb{Z}$ counts “half branch jumps” between λ and $e^{\pm i\rho} \lambda$. Then

$$\widehat{\nu}_0 = -72\frac{\rho}{\pi} + 3b.$$

Sketch proof of theorem for $\eta(B)$

Theorem

$$\eta(B) \rightarrow \frac{1}{\pi} \sum_{\substack{\lambda \in \text{Spec} U \\ \lambda \neq -1}} \arg \lambda$$

as $T \rightarrow \infty$, for $U := e^{\pm i\rho} R_+ R_-$ on $H^{2,\pm}(K3; \mathbb{C})$.

The proof relies on

Kirk-Lesch gluing formula:

$$\eta(B) \rightarrow \eta(B_+) + \eta(B_-) + \text{Maslov index}$$

as $T \rightarrow \infty$, for B_{\pm} the odd signature operators on manifolds with boundary.

Because M_{\pm} have an S^1 -factor they have an orientation-reversing isometry. Therefore B_{\pm} has spectral symmetry, so $\eta(B_{\pm}) = 0!$

Hence it remains only to evaluate the Maslov index.

The Maslov index

Consider $H^3(T^2 \times K3)$ as a complex vector space, with complex structure $*$.

$$\text{Maslov index} := \frac{1}{\pi} \sum_{\substack{\lambda \in \text{Spec}(-\tilde{R}_+ \tilde{R}_-) \\ \lambda \neq -1}} \arg \lambda$$

where \tilde{R}_\pm is reflection of $H^3(T^2 \times K3)$ in the image of $H^3(M_\pm)$.

Thus it suffices to prove that $-\tilde{R}_+ \tilde{R}_-$ has the same spectrum as U .

$$H^3(T^2 \times K3) \cong H^1(T^2) \otimes H^2(K3) \cong \mathbb{C} \otimes H^{2,+}(K3) \oplus \bar{\mathbb{C}} \otimes H^{2,-}(K3).$$

$\tilde{R}_\pm \cong R_{\partial_{u_\pm}} \otimes R_\pm$, where

$R_{\partial_{u_\pm}} : H^1(T^2) \rightarrow H^1(T^2)$ is reflection in internal circle direction of M_\pm

$R_\pm : H^2(K3) \rightarrow H^2(K3)$ is reflection in $\text{Im}(H^2(V_\pm) \rightarrow H^2(K3))$ as before.

$-R_{\partial_{u_+}} R_{\partial_{u_-}}$ is rotation by $\rho = \pi - 2\theta$. Therefore on $H^{2,\pm}(K3; \mathbb{C})$

$$-\tilde{R}_+ \tilde{R}_- \cong e^{\pm i\rho} R_+ R_-$$