

# **Complete cohomogeneity one solitons for $G_2$ Laplacian flow**

Johannes Nordström  
University of Bath

24 March 2025

Joint work with Mark Haskins and Rowan Juneman

arXiv:2112.09095 & arXiv:2501.05437

These slides available at  
<http://people.bath.ac.uk/jlpn20/g2sol4.pdf>

# Introduction

---

**Context:** Riemannian 7-manifolds with holonomy group  $G_2$ ,  
a special kind of Ricci-flat manifolds

Bryant's Laplacian flow: a cousin of Ricci flow for closed  $G_2$ -structures

$G_2$  solitons: self-similar solutions to Laplacian flow

We have found asymptotically conical  $G_2$  solitons of cohomogeneity one  
on  $\Lambda^2 \mathbb{C}P^2$  and  $\Lambda^2 S^4$ , of all three types (shrinker, expander and steady),  
as well as complete solitons with different end behaviours.

## Outline

1. Laplacian flow, solitons, and AC  $G_2$ -structures
2. Main existence results
3. Invariant soliton ODE and singular initial value problem
4. Forward-completeness

# 1. Laplacian flow, solitons, and AC $G_2$ -structures

## Riemannian holonomy $G_2$

---

$G_2 := \text{Aut } \mathbb{O}$ ,  $\mathbb{O} = \text{octonions, normed division algebra of real dimension 8.}$

$G_2$  acts on  $\text{Im } \mathbb{O} \cong \mathbb{R}^7$ , preserving metric, orientation, cross product

$G_2$  is the stabiliser in  $GL(7, \mathbb{R})$  of a stable  $\varphi_0 \in \Lambda^3(\mathbb{R}^7)^*$   
(i.e. the  $GL(7, \mathbb{R})$ -orbit of  $\varphi_0$  is open).

$\varphi \in \Omega^3(M^7)$  pointwise equivalent to  $\varphi_0$  defines a  $G_2$ -structure.

Because  $G_2 \subset SO(7)$ , such a  $\varphi$  induces a metric and orientation.

$Hol(M) \subseteq G_2 \Leftrightarrow$  metric induced by some  $G_2$ -structure  $\varphi$  such that  $\nabla \varphi = 0$ .  
Then call  $\varphi$  *torsion-free*. This is equivalent to the first-order non-linear PDE

$$d\varphi = d^*\varphi = 0.$$

Metrics with holonomy  $G_2$  are always Ricci-flat.

All known constructions of examples on closed manifolds ([Joyce 1994...](#))  
solve the elliptic PDE by gluing together pieces with dimensional reduction.

# Bryant's Laplacian flow

---

Solve

$$\frac{d\varphi_t}{dt} = \Delta_{\varphi_t} \varphi_t$$

with initial condition  $\varphi_0$  satisfying  $d\varphi_0 = 0$ . (Then  $d\varphi_t = 0$  for all  $t$ .)

Starting observations

- $d\varphi = dd^*\varphi = 0 \Rightarrow d^*\varphi = 0$ , so the stationary points are exactly the torsion-free  $G_2$ -structures.
- *Upward* gradient flow for  $\text{vol}(\varphi)$  restricted to the cohomology class of  $\varphi_0$  (the Hitchin functional)
- Induced metric evolves by

$$\frac{dg_t}{dt} = -2\text{Ric}(g_t) + \text{terms quadratic in the torsion } d^*\varphi_t$$

(while  $\text{Ric}(g_t)$  is linear in  $d^*\varphi_t$ )

# What do we know?

---

## Theorem (Bryant-Xu 2011, Lotay-Wei 2017)

- *Short-time existence and uniqueness.*
- *The stationary points are stable:*  
*initial conditions close to a torsion-free  $G_2$ -structure  $\varphi_0$  lead to flow defined for all time, with limit isomorphic to  $\varphi_0$ .*

Lotay-Wei also show that the Riemann curvature or gradient of torsion must blow up at a finite-time singularity, but little is known about the long-term behaviour in general.

Absent an analogue of Perelman's no-local-collapse theorem for Ricci flow it is not known how to obtain blow-up models for singularities.

Nevertheless, solitons for the flow should play a role in the eventual picture...

## $G_2$ soliton equations

$G_2$ -structure  $\varphi$ , vector field  $X$ , dilation constant  $\lambda \in \mathbb{R}$  satisfying

$$\begin{cases} d\varphi = 0, \\ \Delta_\varphi \varphi = \lambda \varphi + \mathcal{L}_X \varphi \end{cases}$$

$\Leftrightarrow$  self-similar solution of Laplacian flow

$$\varphi_t = k(t)^3 f_t^* \varphi$$

for  $k(t) = \frac{3 + 2\lambda t}{3}$  and diffeomorphisms  $f_t$  such that  $\frac{df_t}{dt} = k(t)^{-2} X$

$\lambda > 0$ : expanders (immortal solutions)

$\lambda = 0$ : steady solitons (eternal solutions)

$\lambda < 0$ : shrinkers (ancient solutions)

- Non-steady soliton  $\Rightarrow \varphi$  exact
- Scaling behaviour:  $(\varphi, X)$  is a  $\lambda$ -soliton  $\Leftrightarrow (k^3 \varphi, k^{-2} X)$  is a  $k^{-2} \lambda$ -soliton.

## Asymptotically conical solitons

---

Solitons for the Laplacian flow should play a role in the eventual picture...  
...but not compact ones.

- There can be no compact shrinkers, because the Laplacian flow is the upward gradient flow for volume.
- Any compact steady soliton must be static ( $\varphi$  torsion-free,  $X = 0$ )
- No known examples of compact expanders (or even exact  $G_2$ -structures)

A natural class of non-compact solitons that has been studied in Ricci flow and mean curvature flow is that of asymptotically conical ones.

- AC shrinkers provide models for the formation of an isolated conical finite-time singularity
- AC expanders provide models for resolving an isolated conical singularity
- A shrinker and an expander whose asymptotic cones match provide a model for “flowing through” a singularity.

# Asymptotic cones of Laplacian solitons

$SU(3)$ -structure on  $\Sigma^6$

$\leftrightarrow$  Hermitian 2-form  $\omega \in \Omega^2(\Sigma)$  and (normalised) real part  $\alpha \in \Omega^3(\Sigma)$   
of complex 3-form with respect to some almost complex structure  
 $\leftrightarrow$  conical  $G_2$ -structure  $\varphi_C$  on  $\mathbb{R}_{>0} \times \Sigma$ .

$$\varphi_C = r^2 dr \wedge \omega + r^3 \alpha \rightsquigarrow g_C = dr^2 + r^2 g_\Sigma$$

$\varphi_C$  is torsion-free if the  $SU(3)$ -structure  $(\omega, \alpha)$  is “nearly Kähler”.

For  $X = -\frac{\lambda r}{3} \frac{\partial}{\partial r}$

$$\mathcal{L}_X \varphi_C = -\lambda \varphi_C,$$

so if  $\varphi_C$  is torsion-free cone then  $(\varphi_C, X)$  is a “Gaussian”  $\lambda$ -soliton.

**For  $\lambda \neq 0$**   $(\varphi_C, X)$  is a sensible asymptotic model for an AC  $\lambda$ -soliton  
even if merely  $d\varphi_C = 0$ , because  $\Delta_\varphi \varphi$  has lower order

**For  $\lambda = 0$**  the only reasonable asymptotic cones are static,  
i.e.  $\varphi_C$  torsion-free and  $X = 0$ .

## Bryant-Salamon AC $G_2$ metrics

Where to look for Laplacian solitons  $(M, \varphi)$  that are asymptotically conical, i.e.  $M \setminus (\text{compact set}) \cong \mathbb{R}_+ \times \Sigma^6$  and, for some “asymptotic rate”  $\nu < 0$ ,

$$\varphi = r^2 dr \wedge \omega + r^3 \alpha + O(r^\nu)?$$

Try spaces with well-known AC torsion-free  $G_2$ -structures (ideally exact).

**Bryant-Salamon (1987)** found the first examples of complete  $G_2$  metrics.

These examples are AC, and moreover they have a cohomogeneity 1 action by a group  $G$ , i.e. the generic orbit  $\Sigma$  has dimension 6.

$M$	$G$	$\Sigma$	$\nu$
$\Lambda^2 S^4$	$Sp(2)$	$\mathbb{C}P^3$	-4
$\Lambda^2 \mathbb{C}P^2$	$SU(3)$	$SU(3)/T^2$	-4
$S^3 \times \mathbb{R}^4$	$SU(2)^3$	$S^3 \times S^3$	-3

Remark: In the last two cases,  $\Sigma$  has a  $\mathbb{Z}_3$  of automorphisms that do not extend to diffeomorphisms of  $M$

$\rightsquigarrow G_2$  conifold transitions: 3 topologically distinct ways to glue in zero section to resolve conical singularity  $\mathbb{R}_{>0} \times \Sigma$ .

## Invariant $G_2$ -structures on $\Lambda_-^2 S^4$ and $\Lambda_-^2 \mathbb{C}P^2$

---

$Sp(2)$ -invariant  $G_2$ -structures  $\varphi$  on  $\Lambda_-^2 S^4 \setminus$  zero section  $\cong \mathbb{R}_{>0} \times \mathbb{C}P^3$

$$\leadsto f_1, f_2 : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$$

$\leftrightarrow$  scale of base and  $S^2$  fibres of  $\mathbb{C}P^3 \rightarrow S^4$ .

$SU(3)$ -invariant  $G_2$ -structures on  $\Lambda_-^2 \mathbb{C}P^2 \setminus$  zero section  $\cong \mathbb{R}_{>0} \times SU(3)/T^2$

$$\leadsto f_1, f_2, f_3 : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$$

$\leftrightarrow$  scale of  $S^2$  fibres of three different fibrations  $SU(3)/T^2 \rightarrow \mathbb{C}P^2$ .

Cones:  $G_2$ -structure  $\varphi_C$  defined by  $f_i = c_i r$ .

Closed cones:  $d\varphi_C = 0 \Leftrightarrow$  scale-normalisation of  $c$

Unique holonomy  $G_2$  cone:  $f_1 = f_2 = f_3 = \frac{r}{2}$

The Bryant-Salamon  $G_2$  metrics on  $\Lambda_-^2 S^4$  and  $\Lambda_-^2 \mathbb{C}P^2$  are defined by the same  $f_1$  and  $f_2$ , with  $f_3 = f_2$  in the latter case. Each  $\frac{f_i}{r} \rightarrow \frac{1}{2}$  as  $r \rightarrow \infty$ .

## Anti-self-dual bundle construction

---

Multiplication by  $-1$  on the fibres of  $\Lambda^2_- S^4$  is an isometry of any  $Sp(2)$ -invariant  $G_2$ -structure.

Multiplication by  $-1$  on the fibres of  $\Lambda^2_- \mathbb{C}P^2$  is an isometry of the  $SU(3)$ -invariant  $G_2$ -structure defined by  $(f_1, f_2, f_3) \Leftrightarrow f_2 = f_3$

The equations encoding the torsion-free condition for  $SU(3) \times \mathbb{Z}_2$ -invariant and  $Sp(2)$ -invariant  $G_2$ -structures are identical.

Indeed, Bryant-Salamon derived the same equations for certain  $G_2$ -structures on  $\Lambda^2_- X$  for any positive Einstein self-dual  $X$  (but  $X = S^4$  or  $\mathbb{C}P^2$  are the only complete smooth possibilities).

Similarly, the ODEs for  $Sp(2)$ -invariant solitons can be regarded as a subsystem of the  $SU(3)$ -invariant soliton ODEs.

## 2. Main existence results

### Explicit asymptotically conical shrinkers

---

Shrinkers provide models for formation of singularities, and tend to be rare.

#### Theorem A

*There is a complete  $SU(3) \times \mathbb{Z}_2$ -invariant shrinking  $G_2$ -soliton on  $\Lambda^2_- \mathbb{C}P^2$ , and an  $Sp(2)$ -invariant one on  $\Lambda^2_- S^4$ , defined by  $\lambda = -1$  and*

$$f_1 = r, \quad f_2 = f_3 = \sqrt{\frac{9}{4} + \frac{r^2}{4}}, \quad X = \left( \frac{r}{3} + \frac{4r}{9+r^2} \right) \frac{\partial}{\partial r}.$$

This is asymptotically conical because  $\frac{f_i}{r} \rightarrow c_i$  (for  $c = (1, \frac{1}{2}, \frac{1}{2})$ ).

The asymptotic cone is characterised by  $\frac{c_1}{c_2} = \lim_{r \rightarrow \infty} \frac{f_1}{f_2} = 2$ .

The asymptotic rate is  $\nu = -2$  because  $\frac{f_i}{r} = c_i + O(r^{-2})$ .

Conjecture: This is the unique complete  $Sp(2)$ -invariant shrinker.

# Asymptotically conical expanders on $\Lambda_-^2 S^4$

---

Expanders provide models for how the flow can smooth out a singularity.

## Theorem B

*Every complete  $Sp(2)$ -invariant expanding  $G_2$ -soliton on  $\Lambda_-^2 S^4$  is AC with rate  $-2$ .*

*Up to scale, there is precisely a 1-parameter family of such expanders.*

*Their asymptotic limits are distinct, bijecting with  $(0, 1)$  by  $\frac{c_1}{c_2} = \lim_{r \rightarrow \infty} \frac{f_1}{f_2}$ .*

Keeping the scale fixed, the family can be parametrised by  $\lambda > 0$ .

Limit as  $\lambda \rightarrow 0$  is the Bryant-Salamon  $G_2$ -manifold (which has  $c_1 = c_2 = \frac{1}{2}$ ), considered as a static  $G_2$ -soliton.

## Remark

The asymptotic cone of the explicit AC shrinker on  $\Lambda_-^2 S^4$  ( $\frac{c_1}{c_2} = 2$ ) does *not* match the asymptotic cone of any AC  $Sp(2)$ -invariant expander ( $\frac{c_1}{c_2} < 1$ ).

## Asymptotically conical expanders on $\Lambda_-^2 \mathbb{C}P^2$

---

Theorem B also yields a corresponding 1-parameter family of  $SU(3) \times \mathbb{Z}_2$ -invariant expanders on  $\Lambda_-^2 \mathbb{C}P^2$ .

AC expander ends are stable, making it possible to perturb this 1-parameter family of  $SU(3) \times \mathbb{Z}_2$ -invariant solutions to obtain

### Theorem C

*Up to scale,  $\Lambda_-^2 \mathbb{C}P^2$  admits a 2-parameter family of  $SU(3)$ -invariant expanding  $G_2$ -solitons that are AC with rate  $-2$ .*

We do not expect every complete  $SU(3)$ -invariant expander to be AC.

$$f_1 = \frac{3}{\lambda r}, \quad f_2^2 = f_3^2 = r e^{\frac{\lambda r^2}{6}}$$

solves the soliton ODEs to leading order, and can be corrected to forward-complete solutions with doubly exponential volume growth.

Conjecture: The boundary of the 2-parameter family of  $SU(3)$ -invariant AC expanders corresponds to complete expanders with such ends.

## Flowing through conical singularity?

---

If a singularity forms modelled on the explicit AC shrinker on  $\Lambda_-^2 S^4$ , then no  $Sp(2)$ -invariant expander provides a model for how to smooth it out again.

Harder to control which closed  $SU(3)$ -invariant cones over  $SU(3)/T^2$  appear as asymptotic limits of complete  $SU(3)$ -invariant expanders on  $\Lambda_-^2 \mathbb{C}P^2$ , but numerics suggest:

the asymptotic cone of the shrinker on  $\Lambda_-^2 \mathbb{C}P^2$   
does match

the asymptotic cone of some  $SU(3)$ -invariant expander on  $\Lambda_-^2 \mathbb{C}P^2$

after applying an order 3 automorphism to  $SU(3)/T^2$  that does not extend to  $\Lambda_-^2 \mathbb{C}P^2$  (instead permuting 3 different  $S^2$ -fibrations  $SU(3)/T^2 \rightarrow \mathbb{C}P^2$ )

↔ potential model for “flowing through” the singularity, crushing a  $\mathbb{C}P^2$  and inflating it again in one of two topologically different ways.

This would realise a “ $G_2$  conifold transition”.

# Complete steady solitons

---

All known complete examples of steady Ricci solitons have sub-Euclidean volume growth. In contrast

## Theorem D

*There is precisely a 1-parameter family of  $SU(3)$ -invariant AC steady  $G_2$  solitons on  $\Lambda^2 \mathbb{C}P^2$ , all asymptotic with rate  $-1$  to the unique  $SU(3)$ -invariant torsion-free cone ( $f_i = \frac{r}{2} + O(1)$ ).*

One limit is again the static soliton on the Bryant-Salamon AC  $G_2$ -manifold.

The other limit is an explicit complete steady  $G_2$ -soliton:

$$f_1 = \sqrt{1 + e^{-r}}, \quad f_2 = \sqrt{1 + e^r}, \quad f_3 = 2 \sinh \frac{r}{2}, \quad X = \tanh \frac{r}{2} \frac{\partial}{\partial r}$$

Asymptotic geometry:

In one *other* fibration  $SU(3)/T^2 \rightarrow \mathbb{C}P^2$ , the  $S^2$  fibres have constant size, and the base is the sinh cone over  $\mathbb{C}P^2$ , i.e. the negative Einstein metric

$$dr^2 + (\sinh r)^2 g_{\mathbb{C}P^2} \quad \text{on} \quad \mathbb{R}_+ \times \mathbb{C}P^2.$$

### 3. Initial value problem for invariant solitons

#### Invariant $G_2$ -structures

---

$SU(3)$ -invariant  $G_2$ -structures  $\varphi$  on  $\mathbb{R}_+ \times SU(3)/T^2$  such that

- $\|\frac{\partial}{\partial r}\| = 1$ , and
- restriction to each slice  $SU(3)/T^2$  is closed

are parametrised by triples of functions  $f_1, f_2, f_3 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ .

$$\varphi = dr \wedge (f_1^2 \omega_1 + f_2^2 \omega_2 + f_3^2 \omega_3) + f_1 f_2 f_3 \alpha$$

for  $\omega_i \in \Omega^2(SU(3)/T^2)$  and  $\alpha \in \Omega^3(SU(3)/T^2)$   $SU(3)$ -invariant.

$f_i$  = scale of  $S^2$  fibres of one of three possible fibrations  $SU(3)/T^2 \rightarrow \mathbb{C}P^2$

$\text{Vol}(SU(3)/T^2)$  proportional to  $(f_1 f_2 f_3)^2$

Ones with  $f_2 = f_3$  have extra  $\mathbb{Z}_2$  symmetry, and also define analogous  $Sp(2)$ -invariant  $G_2$ -structures on  $\mathbb{R}_+ \times \mathbb{C}P^3$ .

(Then  $f_1$  = scale of  $S^2$  fibres of  $\mathbb{C}P^3 \rightarrow S^4$ , and  $f_2 = f_3$  a scale of base.)

## Closure and soliton ODE

---

$$d\varphi = 0 \Leftrightarrow 2 \frac{d}{dr} (f_1 f_2 f_3) = f_1^2 + f_2^2 + f_3^2$$

Cones  $\Leftrightarrow f_i = c_i r$  linear

Then  $d\varphi = 0 \Leftrightarrow 6c_1 c_2 c_3 = c_1^2 + c_2^2 + c_3^2$  is a scale-normalisation:

Unique choice of “cone angle” to make a closed cone for each homothety class on  $SU(3)/T^2$

$\leadsto$  **2-parameter family of closed  $SU(3)$ -invariant cones**

(1-parameter family with  $c_2 = c_3$ , giving  $Sp(2)$ -invariant closed cones)

The soliton condition for  $\varphi = f_1^2 \omega_1 + f_2^2 \omega_2 + f_3^2 \omega_3 + f_1 f_2 f_3 \alpha$  and  $X = u \frac{\partial}{\partial r}$  is naively a 2nd-order ODE system for  $(f_1, f_2, f_3, u)$  (with some constraints).

Useful to rewrite it as a 1st-order ODE in  $(f_1, f_2, f_3, \tau_1, \tau_2, \tau_3)$ , where  $\tau_i$  capture the torsion by  $d^* \varphi = \tau_1 \omega_1 + \tau_2 \omega_2 + \tau_3 \omega_3$ .

This is a system in 5 variables after taking into account that

$$d\varphi = 0 \Rightarrow \varphi \wedge d^* \varphi = 0 \Rightarrow \frac{\tau_1}{f_1^2} + \frac{\tau_2}{f_2^2} + \frac{\tau_3}{f_3^2} = 0.$$

## Smooth extension problem

---

Suppose that  $H \subset G$ , that  $H$  acts on a vector space  $V$ , and that the action is transitive on the unit sphere in  $V$ , with stabiliser  $K \subset H$ .

Then think of the vector bundle  $G \times_K V := (G \times V)/K \rightarrow G/K$  as

$$\text{zero section } G/K \quad \sqcup \quad \mathbb{R}_+ \times G/H.$$

To find complete structures on  $G \times_K V$ , the first step is to ask:

Which solutions on  $(0, \epsilon) \times G/H$  extend smoothly over  $G/K$  at  $r = 0$ ?

Applying methods **Eschenburg-Wang (2000)**

- Identify conditions on  $f_i : [0, \epsilon) \rightarrow \mathbb{R}_+$  to ensure smooth extension of  $\varphi = f_1^2 \omega_1 + f_2^2 \omega_2 + f_3^2 \omega_3 + f_1 f_2 f_3 \alpha$  from  $(0, \epsilon) \times SU(3)/T^2$  to  $\mathbb{C}P^2$ :
  - $f_1$  odd with  $f_1'(0) = 1$  (so that  $S^2$  fibres shrink to zero at right rate)
  - $f_2$  and  $f_3$  even with  $f_2(0) = f_3(0) = b = \sqrt[4]{\text{Vol}(\mathbb{C}P^2)} > 0$
- Then solve by power series.

# Solutions to the soliton initial value problem

## Proposition

For each  $\lambda \in \mathbb{R}$ , there is a 2-parameter family  $\varphi_{b,c}$  of solutions to the  $G_2$ -soliton equation with dilation constant  $\lambda$  defined for small  $r$  that extend smoothly to (a neighbourhood of zero section in)  $\Lambda^2_- \mathbb{C}P^2$ .

Two scale-invariant parameters:  $\lambda b^2$  and  $\frac{c}{b}$ .

↪ up to scale there are 2-parameter families of local expanders and shrinkers on  $\Lambda^2_- \mathbb{C}P^2$ , and 1-parameter family of local steady solitons.

The parameter  $b$  is  $\sqrt[4]{\text{Vol}(\mathbb{C}P^2)}$ , while  $c$  controls the leading term in  $f_2 - f_3$ .

The subfamily  $\varphi_{b,0}$  has  $f_2 = f_3$ , so

- has extra  $\mathbb{Z}_2$ -symmetry (multiplication by  $-1$  on fibres of  $\Lambda^2_- \mathbb{C}P^2$ )
- also defines solution near zero section of  $\Lambda^2_- S^4$ .

↪ up to scale there are 1-parameter families of local expanders and shrinkers on  $\Lambda^2_- S^4$ , and a unique local steady soliton.

The latter defines the static soliton on the Bryant-Salamon  $G_2$ -manifold.

Hence there are no non-trivial  $Sp(2)$ -invariant steady solitons on  $\Lambda^2_- S^4$ .

## 4. Forward-completeness

### Scale decoupling and AC ends for steady solitons

---

The steady case  $\lambda = 0$  has a very different character because the scale

$$g := \sqrt[3]{f_1 f_2 f_3} = \sqrt[6]{\text{vol}(\Sigma)}$$

essentially decouples from the homothety class

$$\left( \frac{f_1}{g}, \frac{f_2}{g}, \frac{f_3}{g} \right)$$

The latter evolves in a surface under a 2nd order autonomous ODE  
 $\Leftrightarrow$  1st order ODE in 4 parameters

Torsion-free cone  $c_1 = c_2 = c_3 = \frac{1}{2}$  is unique fixed point, and stable

$\Rightarrow$  Solutions with  $\frac{f_i}{g}$  bounded are asymptotic to the torsion-free cone.

Eigenvalues of linearisation at fixed point give rate  $-1$ .

Since  $\varphi_{b,0}$  is AC (static Bryant-Salamon),  $\varphi_{b,c}$  is AC too for  $c$  near 0.

# Trichotomy for $SU(3)$ -invariant steady ends

## Proposition

Any initial condition for an  $SU(3)$ -invariant steady soliton on  $\mathbb{R} \times SU(3)/T^2$  leads to one of the following behaviours forward in time (up to permuting  $f_i$ )

- (i) AC with rate  $-1$  to torsion-free cone
- (ii) Complete with exponential volume growth:  $f_1 \rightarrow \frac{1}{k}$ , while  $f_2 \sim f_3 \sim e^{kr}$ .
- (iii)  $f_1 = O(\sqrt{r_* - r})$ ,  $f_2, f_3 = O((r_* - r)^{-1/4})$  near finite extinction time  $r_*$ .

We can decide the type of each smoothly closing local solution  $\varphi_{b,c}$  thanks to spotting an explicit solution of type (ii) corresponding to  $\varphi_{\sqrt{2},3}$

$$f_1 = \sqrt{1 + e^{-r}}, \quad f_2 = \sqrt{1 + e^r}, \quad f_3 = 2 \sinh \frac{r}{2}, \quad u = \tanh \frac{r}{2}.$$

## Theorem D

For  $\lambda = 0$ , the local solution  $\varphi_{b,c}$  is (i) AC for  $\frac{c^2}{b^2} < \frac{9}{2}$ .

- (ii) Exponentially growing for  $\frac{c^2}{b^2} = \frac{9}{2}$ .
- (iii) Incomplete for  $\frac{c^2}{b^2} > \frac{9}{2}$ .

## Non-steady AC ends

---

The scale does not decouple for  $\lambda \neq 0$ . On the contrary, scaling up any point in the phase space makes  $\lambda$  terms more dominant, causing

### Proposition

Any  $SU(3)$ -invariant non-steady soliton with all ratios  $\frac{f_i}{f_j}$  bounded in forward time is AC with rate  $-2$ .

Schematically, because  $\lambda$  has dimensions of  $\text{length}^{-2}$ , the other factor  $S$  of those terms has dimensions of  $\text{length}^2$ , and satisfies an equation

$$\frac{dS}{dt} = -\lambda\alpha S + \beta$$

where  $\alpha$  and  $\beta$  have dimension of  $\text{length}$ , and  $\alpha > 0$  involves only  $f_i$  (not  $\frac{df_i}{dr}$ )

If  $\frac{\beta}{\alpha} \rightarrow m$  as  $r \rightarrow \infty$  then  $S \rightarrow \frac{m}{\lambda}$ .

$S$  converging despite having dimension  $\text{length}^2 \rightsquigarrow$  AC rate is  $-2$

This behaviour is **stable** for  $\lambda > 0$

**unstable** for  $\lambda < 0$

# AC end solutions

---

## Proposition

For each  $\lambda \neq 0$  and  $(c_1, c_2, c_3)$  such that  $c_1^2 + c_2^2 + c_3^2 = 6c_1c_2c_3$  there exists

- a unique AC end solution if  $\lambda < 0$
- a 2-parameter family of AC end solutions if  $\lambda > 0$

asymptotic to the corresponding closed cone (i.e.  $\frac{f_i}{r} \rightarrow c_i$ ).

Letting  $u = r^{-2}$ , the sign of  $\lambda$  becomes significant in an ODE analogous to

$$\frac{dx}{du} = \lambda \frac{x}{u^2} + a \frac{x}{u} + b \quad \text{with } x \rightarrow 0 \text{ as } u \rightarrow 0.$$

Setting  $a = b = 0$ , the general solution for  $\lambda \neq 0$  is  $\exp\left(\frac{-\lambda}{u}\right)$ .

But  $\exp\left(\frac{-\lambda}{u}\right) \rightarrow 0$  as  $u \rightarrow 0$  only for  $\lambda > 0$ .

**Haskins-Khan-Payne (2022)** prove rigidity of AC ends for gradient shrinking  $G_2$  solitons without cohomogeneity one assumption

## AC shrinkers

---

Heuristic for  $\lambda < 0$ :

Invariant shrinkers on  $\mathbb{R}_+ \times SU(3)/T^2$  are flow lines in 5-dim phase space.

In 4-dimensional space of flow lines

- 2-dimensional submanifold extends across zero section  $\mathbb{C}P^2 \subset \Lambda_-^2 \mathbb{C}P^2$
- 2-dimensional submanifold has AC behaviour

Expect isolated intersections  $\rightsquigarrow$  *finitely* many AC shrinkers on  $\Lambda_-^2 \mathbb{C}P^2$ .

### Theorem A

For  $\lambda = -1$ ,  $\varphi_{\frac{3}{2}, 0}$  is the explicit solution

$$f_1 = r, \quad f_2^2 = f_3^2 = \frac{9}{4} + \frac{r^2}{4}, \quad u = \frac{r}{3} + \frac{4r}{9 + r^2}.$$

$\rightsquigarrow$   $SU(3) \times \mathbb{Z}_2$ -invariant shrinker on  $\Lambda_-^2 \mathbb{C}P^2$  and  $Sp(2)$ -invariant shrinker on  $\Lambda_-^2 S^4$ , AC with rate  $-2$  to cone with  $c_1 = 1$ ,  $c_2 = c_3 = \frac{1}{2}$ .

Conjecture: Unique complete shrinker with this symmetry.

# Trichotomy for $Sp(2)$ -invariant shrinker ends

---

In the  $Sp(2)$ -invariant case, the AC end solutions are a 1-dimensional submanifold of a 2-dimensional space of end solutions.

We can characterise the other possible end behaviours.

## Theorem

*Any initial condition for an  $Sp(2)$ -invariant shrinker on  $\mathbb{R} \times \mathbb{C}P^3$  leads to one of the following behaviours forward in time*

- (i) AC with rate  $-2$  to a closed cone
- (ii) Complete, exponential growth:  $f_1 \sim 4e^{\mu r}$  and  $f_2^2 \sim \frac{1}{\mu} e^{\mu r}$  for  $\mu = \sqrt{\frac{-\lambda}{18}}$ .
- (iii)  $f_1 = O(\sqrt{r_* - r})$ ,  $f_2 = O((r_* - r)^{-1/4})$  near finite extinction time  $r_*$ .

The end in (ii) is modelled on an explicit shrinking soliton on  $\mathbb{R} \times$  Iwasawa manifold, found by [Fowdar \(2022\)](#).

There is an essentially unique such end solution, appearing as a boundary point for the curve of AC ends solutions.

The incomplete end behaviour (iii) is the only stable type.

# Trichotomy for $Sp(2)$ -invariant expander ends

In the expander case,

$$f_1 = \frac{3}{\lambda r}, \quad f_2 = A\sqrt{r} e^{\frac{\lambda r^2}{12}}$$

can be uniquely corrected to a forward-complete end solution for each  $A > 0$ .

This 1-parameter family of end solutions with doubly exponential growth forms a “wall” between two different stable types of forward-evolution.

## Theorem

Any initial condition for an  $Sp(2)$ -invariant expander on  $\mathbb{R} \times \mathbb{C}P^3$  leads to one of the following behaviours forward in time

- (i) AC with rate  $-2$  to a closed cone
- (ii) Complete with doubly exponential growth:  $f_1 \sim \frac{3}{\lambda r}$ , while  $f_2 \sim \sqrt{r} e^{\frac{\lambda r^2}{12}}$ .
- (iii)  $f_1 = O(\sqrt{r_* - r})$ ,  $f_2 = O((r_* - r)^{-1/4})$  near finite extinction time  $r_*$ .

All smoothly-closing  $Sp(2)$ -invariant solutions  $\varphi_{b,0}$  on  $\Lambda_-^2 S^4$  fall in case (i), but we expect an analogous transition to be relevant for smoothly-closing  $SU(3)$ -invariant expanders  $\varphi_{b,c}$  on  $\Lambda_-^2 \mathbb{C}P^2$ .

# $Sp(2)$ -invariant AC expanders on $\Lambda_-^2 S^4$

## Theorem B

- (i) Every  $Sp(2)$ -invariant local expander  $\varphi_{b,0}$  defined near the zero section of  $\Lambda_-^2 S^4$  is AC.
- (ii) Moreover  $\mathcal{L} : b \mapsto \lim_{r \rightarrow \infty} \frac{f_1}{f_2}$  is a continuous bijection  $(0, \infty) \rightarrow (0, 1)$   
(so each closed  $Sp(2)$ -invariant cone  $f_i = c_i t$  such that  $c_1 < c_2$  is the asymptotic cone of unique complete expander).

Elementary techniques suffice for (i) and non-decreasingness of  $\mathcal{L}$ , checking that certain inequalities are preserved.

$\mathcal{L}$  strictly increasing builds on detailed understanding of AC ends with given asymptotic cone.

$b \rightarrow 0$  limit equivalent up to scale to keeping  $b$  fixed and letting  $\lambda \rightarrow 0$ .

This limit is the Bryant-Salamon manifold (with  $c_1 = c_2 = \frac{1}{2}$ ), so  $\mathcal{L}(b) \rightarrow 1$

## 5. Twistor bundles and rescaling limits of ODEs

### Rescalings of the $Sp(2)$ -invariant soliton ODE

---

Recall variables in  $Sp(2)$ -invariant soliton ODEs:

$f_2 \leftrightarrow$  scale of the base of  $\Lambda^2 S^4 \rightarrow S^4$

$f_1 \leftrightarrow$  scale of  $S^2$  fibres when foliating by sphere bundles

Changing variable  $f_2$  to  $\epsilon f_2 \Leftrightarrow$  rescaling “reference metric” on base of  $\Lambda^2 S^4$

Taking  $\epsilon \rightarrow 0$

$\rightsquigarrow$  ODE for solitons on  $\Lambda^2 \mathbb{R}^4$  invariant under  $\text{Isom}(\mathbb{R}^4) = SO(4) \ltimes \mathbb{R}^4$ .

This limit ODE is easier to analyse due to additional scale invariance and a conserved quantity.

The expander system has an explicit solution  $f_1 = \frac{3}{\lambda r}$ ,  $f_2^2 = r e^{\frac{\lambda r^2}{6}}$ , incomplete at  $r = 0$  but helps understand transition in original system.

Limit of  $\varphi_{b,0}$  as  $b \rightarrow \infty \rightsquigarrow$  AC solution in limit system  $\rightsquigarrow \mathcal{L}(b) \rightarrow 0$

Taking  $\epsilon = \sqrt{-1}$

$\rightsquigarrow$  ODE for solitons on  $\Lambda^2 H^4$  invariant under  $\text{Isom}(H^4) = SO(4, 1)$ .

## Twistor space interpretation

---

For  $X^4$  oriented, consider on the sphere bundle in  $\Lambda_-^2 X$  (twistor space)

- volume form  $\omega_1$  on fibres
- tautological 2-form  $\omega_2$

For  $X$  self-dual Einstein, the conditions for the  $\text{Isom}(X)$ -invariant

$$\varphi = dr \wedge (f_1^2 \omega_1 + f_2^2 \omega_2) + f_1 f_2^2 d\omega_1 \in \Omega^3(\Lambda_-^2 X)$$

to be a soliton depend only on the scalar curvature  $\kappa$ . So

$\kappa = 1 \rightsquigarrow \text{Sp}(2)$ -invariant ODE on  $\Lambda_-^2 S^4$ ;

in general get the  $\epsilon = \sqrt{\kappa}$  rescaling.

In particular, the  $\epsilon = 0$  limit can be interpreted as ODE for warped product solitons on  $\mathbb{R} \times S^2 \times Y^4$ , with  $Y$  hyper-Kähler.

The steady case was considered by [Ball \(2022\)](#).

## Rescalings of the $SU(3)$ -invariant soliton ODE

---

In a similar way, rescaling  $f_2$  and  $f_3$  in the  $SU(3)$ -invariant soliton ODEs is related to considering  $\Lambda^2_- X$  for self-dual Kähler-Einstein  $X^4$  for different  $\kappa$ .

(In particular, one could consider  $SU(2, 1)$ -invariant solitons of  $\Lambda^2_- \mathbb{C}H^2$ .)

$f_2 = f_3 \Leftrightarrow$  invariance under multiplying fibres of  $\Lambda^2_- X$  by  $-1$ , recovering ODE from the case not requiring Kähler.

The limit of rescalings of the explicit exponentially growing solution  $f_1 = \sqrt{1 + e^{-r}}$ ,  $f_2 = \sqrt{1 + e^r}$ ,  $f_3 = 2 \sinh \frac{r}{2}$  give an explicit solution

$$f_1 = 1, f_2 = f_3 = e^r$$

on  $\mathbb{R}_{>0} \times S^2 \times Y$  for  $Y$  hyperKähler

$\rightsquigarrow$  complete steady  $G_2$  soliton that is a metric product of  $S^2$  and  $dr^2 + e^r g_Y$

## Another rescaling of $Sp(2)$ -invariant soliton ODE

---

Changing variables  $f_1 \rightarrow \epsilon^2 f_1$ ,  $f_2 \rightarrow \epsilon f_2$  and taking  $\epsilon \rightarrow 0$

$\Leftrightarrow$  rescaling both fibres and base of  $S^2$ -fibration  $\mathbb{C}P^3 \rightarrow S^4$  of link of  $\Lambda^2_- S^4$

$\rightsquigarrow$  soliton ODE on  $\mathbb{R} \times \mathbb{C} \times \mathbb{C}^2$ , invariant under complex Heisenberg group  $\mathcal{H}_3$ .

Same as soliton ODE for certain  $G_2$ -structures on  $\mathbb{R} \times$  suitable  $T^2$ -bundle over hyper-Kähler  $Y^4$ .

This system can be reduced to one in two variables.

Fowdar's explicit shrinker corresponds to a fixed point.

We find another complete shrinker with one AC end and one cusp end modelled on Fowdar's.

The proof of the trichotomy for shrinkers relies on analysing this limit system.

