

Differential Topology

Example Sheet 2

1. Let $0 \rightarrow A^* \xrightarrow{f} B^* \xrightarrow{g} C^* \rightarrow 0$ be a short exact sequence of cochain complexes. Show that the sequence

$$\dots \rightarrow H^k(B^*) \xrightarrow{g} H^k(C^*) \xrightarrow{\delta} H^{k+1}(A^*) \rightarrow \dots$$

in the Snake lemma is exact at $H^k(C^*)$.

2. The Euler characteristic of a manifold M^n is the alternating sum of its Betti numbers (if finite): $\chi(M) = \sum_{i=0}^n (-1)^i b_i(M)$.

- (a) Let U and V be open subsets of a manifold M . Show that (if the terms are well-defined)

$$\chi(U) + \chi(V) = \chi(U \cup V) + \chi(U \cap V).$$

- (b) Let M_1^n, M_2^n be manifolds. Show that

$$\chi(M_1 \# M_2) = \chi(M_1) + \chi(M_2) - \begin{cases} 0 & \text{if } n \text{ is odd} \\ 2 & \text{if } n \text{ is even} \end{cases}$$

- (c) Let M^m, N^n be manifolds. Show that $\chi(M \times N) = \chi(M)\chi(N)$.

3. Let U, V open connected in \mathbb{R}^n such that $\mathbb{R}^n = U \cup V$. Show that $U \cap V$ is connected.
4. Show that $b_k(T^n) = \binom{n}{k}$ for all $0 \leq k \leq n$.
5. (a) Show that $S^2 \times S^4$ is not homotopy equivalent to $\mathbb{C}P^3$.
 (b) Are the punctured manifolds $(S^2 \times S^4) \setminus \{pt\}$ and $\mathbb{C}P^3 \setminus \{pt\}$ homotopy equivalent to any closed manifolds?
6. Let M_1^n, M_2^n be closed connected manifolds of equal dimension. Show that $b_k(M_1 \# M_2) = b_k(M_1) + b_k(M_2)$ for $0 < k \leq n - 2$. What is $b_{n-1}(M_1 \# M_2)$ when
- (a) M_1 and M_2 are both orientable?
 (b) M_1 is orientable and M_2 is not?
 (c) M_1 and M_2 are both nonorientable?
7. Show that there is an orientation-reversing diffeomorphism $\mathbb{C}P^n \rightarrow \mathbb{C}P^n$ if and only if n is odd.
8. Find a pair of connected oriented manifolds M, N that do not have orientation-reversing self-diffeomorphisms such that $M \# N$ does.

9. Let G^n be a compact Lie group, *i.e.* a smooth compact manifold of dimension n with a group structure such that the multiplication and inverse maps are smooth. For $g \in G$, let $L_g : G \rightarrow G$ be the left multiplication by g , *i.e.* $L_g : h \mapsto gh$.

- (a) Show that there is a left-invariant orientation form on G , *i.e.* a non-zero $\omega \in \Omega^n(G)$ such that $L_g^* \omega = \omega$ for any $g \in G$. Show that for any $f : G \rightarrow \mathbb{R}$ (or $f : G \rightarrow V$ for some vector space V)

$$\int_G f \omega = \int_G (f \circ L_g) \omega.$$

- (b) Let M^m be a smooth manifold with a smooth G -action, *i.e.* a smooth map $G \times M \rightarrow M$, $(g, x) \mapsto gx$ such that $g(hx) = (gh)x$. Let $\Omega_G^k(M) = \{\alpha \in \Omega^k(M) : g^* \alpha = \alpha \text{ for all } g \in G\}$. If G is connected, show that there is a projection $A : \Omega^k(M) \rightarrow \Omega_G^k(M)$ such that $[A(\alpha)] = [\alpha] \in H^k(M)$ for any closed $\alpha \in \Omega^k(M)$.

- (c) $U(n) \subset GL(n, \mathbb{C})$ is the subgroup that preserves the hermitian inner product $h_0 = \sum dz^j \otimes d\bar{z}^j$. Alternatively it can be characterised as preserving the real part g_0 of the hermitian product (which is just the Euclidean inner product), or the “Kähler form” $\omega_0 = g_0(i \cdot, \cdot) = \frac{i}{2} \sum dz^j \wedge d\bar{z}^j \in \Lambda_{\mathbb{R}}^2(\mathbb{C}^n)^*$. Show that the natural action of $U(n+1)$ on $\mathbb{C}P^n$ is transitive, with stabiliser isomorphic to $U(1) \times U(n)$. Describe the action of the stabiliser of a point on the tangent space at that point, and show that $\Omega_{U(n+1)}^2(\mathbb{C}P^n)$ is 1-dimensional (a certain normalised non-zero element ω is called the *Fubini-Study form*). Deduce that a generator $c \in H^2(\mathbb{C}P^n)$ satisfies $c^k \neq 0 \in H^{2k}(\mathbb{C}P^n)$ for $0 \leq k \leq n$. (*Hint*: Prove or assume that up to scaling ω_0 is the only real 2-form on \mathbb{C}^n invariant under $U(n)$.)

Questions and corrections to j.nordstrom@imperial.ac.uk.

January 31, 2013