

# Random Parking and Rubber Elasticity

Mathew Penrose  
(*University of Bath*)

Joint work with Antoine Gloria INRIA, Lille

Imperial College Stochastic Analysis Seminar  
January 2013

## Rubber Elasticity

Let  $d, n \in \mathbf{N}$  (e.g.  $d = n = 3$ ).

Suppose  $D \subset \mathbf{R}^d$  is a bounded domain.  $D$  represents a piece of rubber.

Let  $\mathcal{L} \subset \mathbf{R}^d$  be a locally finite point process.

$\mathcal{L} \cap D$  the locations of individual “molecules”. For  $x, y \in \mathcal{L}$  write  $x \sim y$  if they are Delaunay neighbours, let  $\mathcal{T}$  be the Delaunay triangulation.

Each  $u \in C(D, \mathbf{R}^n)$  represents a deformation of the rubber.

$u(x)$  is the location of  $x \in D$  under deformation.

We'll define a class of energy functionals  $F : C(D, \mathbf{R}^n) \rightarrow \mathbf{R}$ .

$F(u)$ , the energy of deformation  $u$ .

Let  $u_{\mathcal{L}}$  be affine on each  $T \in \mathcal{T}$  with  $u_{\mathcal{L}} \equiv u$  on  $\mathcal{L}$

## The energy functional

$$F_{\mathcal{L}}^D(u) = \sum_{x,y \in \mathcal{L}, x \sim y, [x,y] \subset D} |y-x|^d f\left(\frac{u(y)-u(x)}{|y-x|}\right) \\ + \sum_{T \in \mathcal{T}(\mathcal{L}), T \subset D} |T| g(\nabla u_{\mathcal{L}|T})$$

Assume we are given  $f \in C(\mathbb{R}^n, \mathbb{R}^+)$  the bond energy  
and  $g \in C(M^{n \times d}, \mathbb{R}^+)$  the cell energy (given).

Assume growth bounds on  $f, g$ : for some  $p > 1, C > 0$ ,

$$C^{-1} \leq \frac{f(z)}{|z|^p} \leq C, \quad |z| \geq 1$$

$$g(\Lambda) \leq C \|\Lambda\|^p, \quad \|\Lambda\| \geq 1.$$

## Desirable properties of $\mathcal{L}$ , our point process in $\mathbb{R}^d$

$\mathcal{L}$  is *stationary* if  $x + \mathcal{L} \stackrel{\mathcal{D}}{=} \mathcal{L}$ , for all  $x \in \mathbb{R}^d$ .

$\mathcal{L}$  is *isotropic* if  $R\mathcal{L} \stackrel{\mathcal{D}}{=} \mathcal{L}$  for all  $R \in SO_d$ .

$\mathcal{L}$  is in *general position* (or just *general*) if no  $d + 1$  points of  $\mathcal{L}$  lie in the same hyperplane, and no  $d + 2$  points are in the same hypersphere.

Let  $\mathcal{A}_{lf}$  be the class of locally finite point configurations in  $\mathbb{R}^d$ .

$\mathcal{L}$  is *ergodic* if for all  $A \subset \mathcal{A}_{lf}$  with  $T_x(A) = A$  for all  $x \in \mathbb{R}^d$  (where  $T_x$  is translation by  $x$ ) we have  $P[\mathcal{L} \in A] \in \{0, 1\}$ .

For  $0 < a < b$  let  $\mathcal{A}_{a,b} \subset \mathcal{A}_{lf}$  be the class of  $\xi$  such that

$$x, y \in \xi \implies |x - y| > a \quad (\text{hard core condition})$$

$$x \in \mathbb{R}^d \implies \xi \cap B(x, b) \neq \emptyset \quad (\text{no empty space condition})$$

$$B(x, b) := \{y : |y - x| \leq b\}.$$

Let  $\mathcal{A}_{a,\infty}$  be those  $\xi$  satisfying just the hard core condition.

## Gamma-convergence

If  $F_n$  and  $F$  are real-valued functions on some metric space  $X$ .

we say for  $x \in X$  that  $F_n \xrightarrow{\Gamma} F$  at  $x$  if

- (a) For all sequences  $x_n \rightarrow x$  we have  $\liminf F_n(x_n) \geq F(x)$ , and
- (b)  $\exists$  sequence  $x_n \rightarrow x$  with  $F_n(x_n) \rightarrow F(x)$ .

We say  $F_n \xrightarrow{\Gamma} F$  if  $F_n \xrightarrow{\Gamma} F$  at  $x$  for all  $x \in X$ .

Set  $Q_r = [-r/2, r/2]^d$ .

Recall we assume  $f$  (bond energy) and  $g$  (cell energy) satisfy growth bounds of order  $p > 1$ . We now state a

Homogenization result (Alicandro, Cicalese and Gloria 2011)

$$F_{\mathcal{L}}^D(u) = \sum_{x \sim y, [x,y] \subset D} |y-x|^d f\left(\frac{u(y)-u(x)}{|y-x|}\right) + \sum_{T \in \mathcal{T}(\mathcal{L}): T \subset D} |T|g(\nabla u_{\mathcal{L}}|_T)$$

Suppose  $0 < a < b$  and  $\mathcal{L}$  is stationary, ergodic, general and a.s. in  $\mathcal{A}_{a,b}$ .

Then a.s. as  $\varepsilon \downarrow 0$  we have  $F_{\varepsilon\mathcal{L}}^D \xrightarrow{\Gamma} F_{\text{hom}}^D$  on  $L^p(D, \mathbb{R}^n)$ , where

$$F_{\text{hom}}^D(u) = \begin{cases} \int_D W_{\text{hom}}(\nabla u(x)) dx, & u \in W^{1,p}(D, \mathbb{R}^n). \\ +\infty & \text{otherwise} \end{cases}$$

$$W_{\text{hom}}(\Lambda) = \lim_{r \rightarrow \infty} r^{-d} \inf \{ F_{\mathcal{L}}^{Q_r}(u) : u(x) = \Lambda \cdot x \text{ on } Q_r \setminus Q_{r-2b} \} \quad (1)$$

Idea - divide  $D$  into cubes of side  $\delta = \delta(\varepsilon)$  with  $\varepsilon \ll \delta \ll 1$ . Take  $u_\varepsilon = u$  (approx. affine) near boundary of each cube. Let  $u_\varepsilon$  optimise the energy inside each cube, subject to this constraint. Discuss (1) later.

## Existence of a stationary ergodic general $\mathcal{L}$ in $\mathcal{A}_{a,b}$ : Random parking.

Let  $\rho > 0$ . Let  $X_1, X_2, \dots$  be independent uniform random vectors in  $D$ .  $X_n$  is *accepted*, unless  $\exists m \leq n$  with  $X_m$  accepted and  $|X_n - X_m| \leq \rho$ .

Let  $\xi^D = \{\text{accepted } X_i\}$ . (random parking process on  $D$ ). It has the  $a$ -hardcore and  $b$ -no-empty space properties on  $D$  for any  $a < \rho < b$ .

$\xi^{Q_r}$  has weak limit  $\xi$  on  $\mathbb{R}^d$  (stationary ergodic, general, in  $\mathcal{A}_{a,b}$ ).

$\xi$  obtained from parking protocol for homogeneous Poisson point process  $\{(X_i, T_i)\}$  in  $\mathbb{R}^d \times \mathbb{R}^+$ , where  $T_i$  is arrival time of  $X_i$ .

Parking protocol on this Poisson process well-defined by a first passage percolation argument (Penrose 2001).

## Subadditivity

Suppose  $\mathcal{E}(S, R)$  is a real-valued energy functional defined for all locally finite  $S \subset \mathbb{R}^d$ , and rectangles  $R$ , e.g.  $p$ -weighted travelling salesman cost

$$\mathcal{E}_{TSP,p}(S, R) = \min \left\{ \sum_{i=1}^n |x_i - x_{i-1}|^p : S \cap R = \{x_1, \dots, x_n\}, x_0 = x_n \right\}$$

Known since BHH (1959) that there exists  $\beta$  such that as  $r \rightarrow \infty$ ,

$$r^{-d} \mathcal{E}_{TSP,1}(\mathcal{H}, Q_r) \rightarrow \beta$$

where  $\mathcal{H}$  is a homogeneous PPP on  $\mathbb{R}^d$ .

We shall describe generic properties of  $\mathcal{E}$  guaranteeing such convergence for  $\mathcal{E}_{TSP,p}$  and many other examples, e.g. the minimal matching and minimal spanning tree. (cf. Redmond and Yukich (1994), Yukich (1999)).



## Properties of $\mathcal{E}_{TSP,p}(\cdot)$ (and other choices of $\mathcal{E}(\cdot)$ with 'order' $p$ )

- Translation invariant:  $\mathcal{E}(x + S, x + R) = \mathcal{E}(S, R)$ , all  $x \in \mathbb{R}^d$ , all  $S, R$ .
- Almost subadditive:  $\mathcal{E}(S \cup T, R) \leq \mathcal{E}(S, R) + \mathcal{E}(T, R) + C(\text{diam}R)^p$ .
- Smooth:  $|\mathcal{E}(T, R) - \mathcal{E}(S, R)| \leq C(\text{diam}R)^p(\text{card}((S \Delta T) \cap R))^{1-p/d}$ .
- There is an approximate energy functional  $\tilde{\mathcal{E}}(S, R)$  defined for all rectangles  $R \subset \mathbb{R}^d$  with  $\tilde{\mathcal{E}}$  translation invariant, and
- Superadditive: if  $R_0 = \cup_{i=1}^n R_i$  (rectangles) then

$$\tilde{\mathcal{E}}(S, R) \geq \sum_{i=1}^m \tilde{\mathcal{E}}(S, R_i)$$

- Close to  $\mathcal{E}$  of order  $p$ :  $r^{-p}|\tilde{\mathcal{E}}(S, Q_r) - \mathcal{E}(S, Q_r)| = o(\text{card}(S \cap Q_r))$ .

For  $\mathcal{E}_{TSP,p}$  take  $\tilde{\mathcal{E}}(S, R)$  to be the TSP cost with 'free travel' outside  $R$ .

## General LLNs for $\mathcal{E}$

Suppose  $p \geq 1$  and  $\mathcal{E}$  is TI, almost subadditive, and smooth of order  $p$ . Suppose there exists  $\tilde{\mathcal{E}}(S, R)$  which is TI, superadditive and close to  $\mathcal{E}$  of order  $p$ . Then (Redmond/Yukich) there is a constant such that

$$r^{-d}\mathcal{E}(\mathcal{H}, Q_r) \rightarrow \gamma \text{ a.s.}$$

Moreover (Gloria-P. 2012), if  $\mathcal{L}$  is stationary and ergodic with  $\mathcal{L} \in \mathcal{A}_{a,\infty}$  for some  $a > 0$ , then there exists  $\gamma \in \mathbb{R}$  such that

$$r^{-d}\mathcal{E}(\mathcal{L}, Q_r) \rightarrow \gamma \text{ a.s.}$$

Proof via multiparameter subadditive ergodic theorem (Akcoğlu/Krengel).

Example: take  $\mathcal{E}(S, R) = \inf\{F_S^R(u) : u(x) = \Lambda x \text{ near } \partial R\}$ .

## Back to random parking

Results described so far show that if  $\xi$  is random parking on  $\mathbb{R}^d$  then for  $\mathcal{E}$  satisfying hypotheses of subadditivity etc., we have for some  $\gamma$  that

$$r^{-d} \mathcal{E}(\xi, Q_r) \rightarrow \gamma$$

e.g. with  $\mathcal{E}$  the  $p$ -weighted TSP on  $\xi \cap Q_R$   
or with  $\mathcal{E}$  the minimum of  $F_\xi^{Q_r}(u)$  given  $\Lambda$ -boundary conditions.

Would like to replace  $\xi$  by  $\xi^{Q_r}$  in the above results, since (i) any simulation studies would be on a finite region (ii) no physical reason for process generating  $\xi$  on  $D = Q_r$  to depend on input from outside  $D$   
Gloria-P. (2012): can indeed replace  $\xi$  by  $\xi^{Q_r}$  in the above.

Also: can extend the earlier homogenization result:

$$F_{\mathcal{L}}^D(u) = \sum_{x \sim y, [x,y] \subset D} |y-x|^d f\left(\frac{u(y)-u(x)}{|y-x|}\right) + \sum_{T \in \mathcal{T}(\mathcal{L}): T \subset D} |T| g(\nabla u_{\mathcal{L}}|_T)$$

Suppose  $\xi_{\rho}^D$  is the random parking process in  $D$  with hard-core parameter  $\rho$ . Then for  $\rho > 0$ , as  $\varepsilon \downarrow 0$  we have  $F_{\xi_{\varepsilon\rho}^D}^D \xrightarrow{\Gamma} F_{\text{hom}}^D$  on  $L^p(D, \mathbb{R}^n)$ , where

$$F_{\text{hom}}^D(u) = \begin{cases} \int_D W_{\text{hom}}(\nabla u(x)) dx, & u \in W^{1,p}(D, \mathbb{R}^n). \\ +\infty & \text{otherwise} \end{cases}$$

$$W_{\text{hom}}(\Lambda) = \lim_{r \rightarrow \infty} r^{-d} \inf \{ F_{\xi_{Q_r}^{Q_r}}^{Q_r}(u) : u(x) = \Lambda \cdot x \text{ on } Q_r \setminus Q_{r-2b} \}. \quad (2)$$

Proof relies heavily on exponential stabilization of random parking (Schreiber, P. and Yukich 2007)

## Exponential stabilization

Let  $\mathcal{H}$  be a homogeneous Poisson process in  $\mathbb{R}^d \times \mathbb{R}^+$ .

For  $x \in \mathbb{Z}^d$ , let  $Q_x = x + [0, 1]^d$ .

Let  $\xi_x = \xi(\mathcal{H}) \cap (Q_x \times [0, \infty))$ , where  $\xi(\cdot)$  is the set of accepted points for the random parking process with hard-core parameter 1.

Exponential stabilization says that for each  $x$  there exists  $R_x$  such that:

- $\xi_x$  is unaffected by changes to  $\mathcal{H}$  outside  $B_{R_x}(x) \times [0, \infty)$ .
- $P[R_x > t] \leq e^{-\alpha t}$ , some  $\alpha > 0$