

# MA40189 - Useful results for Markov chains

<https://people.bath.ac.uk/masss/ma40189.html>

Simon Shaw, s.shaw@bath.ac.uk

2021/22 Semester II

# Irreducible

## Definition

A Markov chain is said to be **irreducible** if for every  $i, j$  there exists  $k$  such that

$$P(X_{t+k} = j | X_t = i) > 0$$

that is, all states can be reached from any other state in a finite number of moves.

# Periodic/Aperiodic

## Definition

A state  $i$  is said to be **periodic** with period  $d_i$  if starting from state  $i$  the chain returns to it within a fixed number of steps  $d_i$  or a multiple of  $d_i$ .

$$d_i = \gcd\{t : P(X_t = i | X_0 = i) > 0\}$$

where  $\gcd$  is the greatest common divisor. If  $d_i = 1$  then the state  $i$  is said to be **aperiodic**.

Irreducible Markov chains have the property that all states have the same period. A Markov chain is called **aperiodic** if some (and hence all) states are aperiodic.

# Recurrent/Positive Recurrent

## Definition

Let  $\tau_{ii}$  be the time of the first return to state  $i$

$$\tau_{ii} = \min\{t > 0 : X_t = i \mid X_0 = i\}$$

A state  $i$  is **recurrent** if  $P(\tau_{ii} < \infty) = 1$  and **positive recurrent** if  $E(\tau_{ii}) < \infty$ .

Thus, a state  $i$  is recurrent if the chain will return to state  $i$  with probability 1 and positive recurrent if, with probability 1, it will return in a finite time. An irreducible Markov chain is positive recurrent if some (and hence all) states  $i$  are positive recurrent.

# Ergodic

## Definition

A state is said to be **ergodic** if it is aperiodic and positive recurrent. A Markov chain is ergodic if all of its states are ergodic.

# Stationary

## Definition

A distribution  $\pi$  is said to be a **stationary distribution** of a Markov chain with transition probabilities  $P_{ij} = P(X_t = j \mid X_{t-1} = i)$ , if

$$\sum_{i \in S} \pi_i P_{ij} = \pi_j \quad \forall j \in S$$

where  $S$  denotes the state space.

In matrix notation if  $P$  is the matrix of transition probabilities and  $\pi$  the vector with  $i$ th entry  $\pi_i$  then the stationary distribution satisfies

$$\pi = \pi P$$

# Existence and uniqueness

## Theorem

*Each irreducible and aperiodic Markov chain has a unique stationary distribution  $\pi$ .*

# Convergence

## Theorem

Let  $X_t$  be an irreducible and aperiodic Markov chain with stationary distribution  $\pi$  and arbitrary initial value  $X_0 = x_0$ . Then

$$P(X_t = x \mid X_0 = x_0) \rightarrow \pi(x)$$

as  $t \rightarrow \infty$ .



# Ergodic

## Theorem

Let  $X_t$  be an ergodic Markov chain with limiting distribution  $\pi$ . If  $E\{g(X) | X \sim \pi(x)\} < \infty$  then the sample mean converges to the expectation of  $g(X)$  under  $\pi$ ,

$$P \left\{ \frac{1}{N} \sum_{i=1}^N g(X_i) \rightarrow E\{g(X) | X \sim \pi(x)\} \right\} = 1.$$

## Consequence of these theorems

*If we can construct an ergodic Markov chain  $\theta_t$  which has the posterior distribution  $f(\theta | x)$  as the stationary distribution  $\pi(\theta)$  then, starting from an initial point  $\theta_0$ , if we run the Markov chain for long enough, we will sample from the posterior.*

- *for large  $t$ ,  $\theta_t \sim \pi(\theta) = f(\theta | x)$*
- *for each  $s > t$ ,  $\theta_s \sim \pi(\theta) = f(\theta | x)$*
- *the ergodic averages converge to the desired expectations under the target distribution.*